



planetmath.org

Math for the people, by the people.

uniqueness of additive inverse in a ring

Canonical name	UniquenessOfAdditiveInverseInARing
Date of creation	2013-03-22 14:13:54
Last modified on	2013-03-22 14:13:54
Owner	alozano (2414)
Last modified by	alozano (2414)
Numerical id	7
Author	alozano (2414)
Entry type	Theorem
Classification	msc 20-00
Classification	msc 16-00
Classification	msc 13-00
Related topic	UniquenessOfInverseForGroups

Lemma. *Let R be a ring, and let a be any element of R . There exists a unique element b of R such that $a+b = 0$, i.e. there is a unique <http://planetmath.org/Ringadditiveinverse> for a .*

Proof. Let a be an element of R . By definition of ring, there exists at least one <http://planetmath.org/Ringadditiveinverse> of a , call it b_1 , so that $a + b_1 = 0$. Now, suppose b_2 is another additive inverse of a , i.e. another element of R such that

$$a + b_2 = 0$$

where 0 is the <http://planetmath.org/Ringzero> element of R . Let us show that $b_1 = b_2$. Using properties for a ring and the above equations for b_1 and b_2 yields

$$\begin{aligned} b_1 &= b_1 + 0 \quad (\text{definition of zero}) \\ &= b_1 + (a + b_2) \quad (b_2 \text{ is an additive inverse of } a) \\ &= (b_1 + a) + b_2 \quad (\text{associativity in } R) \\ &= 0 + b_2 \quad (b_1 \text{ is an additive inverse of } a) \\ &= b_2 \quad (\text{definition of zero}). \end{aligned}$$

Therefore, there is a unique additive inverse for a . □