

## planetmath.org

Math for the people, by the people.

## prime residue class

Canonical name PrimeResidueClass
Date of creation 2013-03-22 15:43:12
Last modified on 2013-03-22 15:43:12

Owner pahio (2872) Last modified by pahio (2872)

Numerical id 18

Author pahio (2872)
Entry type Definition
Classification msc 20K01
Classification msc 13M99
Classification msc 11A07
Synonym prime class

Related topic MultiplicativeOrderOfAnIntegerModuloM

Related topic NonZeroDivisorsOfFiniteRing

Related topic GroupOfUnits
Related topic PrimitiveRoot
Related topic ResidueSystems
Related topic Klein4Group
Related topic EulerPhifunction

Related topic SummatoryFunctionOfArithmeticFunction

Defines residue class group

Let m be a positive integer. There are m residue classes  $a+m\mathbb{Z}$  modulo m. Such of them which have

$$\gcd(a, m) = 1,$$

are called the *prime residue classes* or *prime classes modulo* m, and they form an Abelian group with respect to the multiplication

$$(a+m\mathbb{Z})\cdot(b+m\mathbb{Z}) := ab+m\mathbb{Z}.$$

This group is called the residue class group modulo m. Its order is  $\varphi(m)$ , where  $\varphi$  means Euler's totient function. For example, the prime classes modulo 8 (i.e.  $1+8\mathbb{Z}$ ,  $3+8\mathbb{Z}$ ,  $5+8\mathbb{Z}$ ,  $7+8\mathbb{Z}$ ) form a group isomorphic to the Klein 4-group.

The prime classes are the units of the residue class ring  $\mathbb{Z}/m\mathbb{Z} = \mathbb{Z}_m$  consisting of all residue classes modulo m.

Analogically, in the http://planetmath.org/ExamplesOfRingOfIntegersOfANumberFieldrin R of integers of any algebraic number field, there are the residue classes and the prime residue classes modulo an ideal  $\mathfrak a$  of R. The number of all residue classes is  $N(\mathfrak a)$  and the number of the prime classes is also denoted by  $\varphi(\mathfrak a)$ . It may be proved that

$$\varphi(\mathfrak{a}) = \mathrm{N}(\mathfrak{a}) \prod_{\mathfrak{p} \mid \mathfrak{a}} \left( 1 - \frac{1}{\mathrm{N}(\mathfrak{p})} \right);$$

N is the absolute norm of ideal and  $\mathfrak{p}$  runs all distinct prime ideals dividing  $\mathfrak{a}$  (cf. the first formula in the entry "http://planetmath.org/EulerPhiFunctionEuler phi function"). Moreover, one has the result

$$\alpha^{\varphi(\mathfrak{a})} \equiv 1 \pmod{\mathfrak{a}}$$

for  $((a), \mathfrak{a}) = (1)$ , generalising the http://planetmath.org/EulerFermatTheoremEulerFermat theorem.