



Math for the people, by the people.

orthogonal group

Canonical name	OrthogonalGroup
Date of creation	2013-03-22 12:25:54
Last modified on	2013-03-22 12:25:54
Owner	djao (24)
Last modified by	djao (24)
Numerical id	6
Author	djao (24)
Entry type	Definition
Classification	msc 20G20
Defines	orthogonal transformation

Let  $Q$  be a non-degenerate symmetric bilinear form over the real vector space  $\mathbb{R}^n$ . A linear transformation  $T: V \rightarrow V$  is said to *preserve*  $Q$  if  $Q(Tx, Ty) = Q(x, y)$  for all vectors  $x, y \in V$ . The subgroup of the general linear group  $\text{GL}(V)$  consisting of all linear transformations that preserve  $Q$  is called the *orthogonal group* with respect to  $Q$ , and denoted  $\text{O}(n, Q)$ .

If  $Q$  is also positive definite (i.e.,  $Q$  is an inner product), then  $\text{O}(n, Q)$  is equivalent to the group of invertible linear transformations that preserve the standard inner product on  $\mathbb{R}^n$ , and in this case the group  $\text{O}(n, Q)$  is usually denoted  $\text{O}(n)$ .

Elements of  $\text{O}(n)$  are called *orthogonal transformations*. One can show that a linear transformation  $T$  is an orthogonal transformation if and only if  $T^{-1} = T^T$  (i.e., the inverse of  $T$  equals the transpose of  $T$ ).