



Math for the people, by the people.

alternative proof of condition on a near ring to be a ring

Canonical name	AlternativeProofOfConditionOnANearRingToBeARing
Date of creation	2013-03-22 17:20:06
Last modified on	2013-03-22 17:20:06
Owner	Wkbj79 (1863)
Last modified by	Wkbj79 (1863)
Numerical id	9
Author	Wkbj79 (1863)
Entry type	Proof
Classification	msc 20-00
Classification	msc 16-00
Classification	msc 13-00

Theorem 1. *Let $(R, +, \cdot)$ be a near ring with a multiplicative identity 1 such that the \cdot also left distributes over $+$; that is, $c \cdot (a + b) = c \cdot a + c \cdot b$. Then R is a ring.*

Proof. All that needs to be verified is commutativity of $+$.

Let $a, b \in R$. Consider the expression $(1 + 1)(a + b)$.

We have:

$$\begin{aligned} (1 + 1)(a + b) &= (1 + 1)a + (1 + 1)b && \text{by left distributivity} \\ &= 1a + 1a + 1b + 1b && \text{by right distributivity} \\ &= a + a + b + b && \text{since 1 is a multiplicative identity} \end{aligned}$$

On the other hand, we have:

$$\begin{aligned} (1 + 1)(a + b) &= 1(a + b) + 1(a + b) && \text{by right distributivity} \\ &= a + b + a + b && \text{since 1 is a multiplicative identity} \end{aligned}$$

Thus, $a + a + b + b = a + b + a + b$. Hence:

$$\begin{aligned} a + b &= 0 + (a + b) + 0 && \text{since 0 is an } \text{http://planetmath.org/AdditiveId} \\ &= (-a + a) + (a + b) + (b + -b) && \text{by definition of } \text{http://planetmath.org/AdditiveI} \\ &= -a + (a + a + b + b) + -b \\ &= -a + (a + b + a + b) + -b && \text{since } a + a \\ &= (-a + a) + (b + a) + (b + -b) \\ &= 0 + (b + a) + 0 \\ &= b + a \end{aligned}$$

□