



Math for the people, by the people.

alternating group is a normal subgroup of the symmetric group

Canonical name	AlternatingGroupIsANormalSubgroupOfTheSymmetricGroup
Date of creation	2013-03-22 13:42:32
Last modified on	2013-03-22 13:42:32
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	8
Author	CWoo (3771)
Entry type	Theorem
Classification	msc 20-00

Theorem 1. *The alternating group A_n is a normal subgroup of the symmetric group S_n*

Proof. Define the epimorphism $f : S_n \rightarrow \mathbb{Z}_2$ by $\sigma \mapsto 0$ if σ is an even permutation and $\sigma \mapsto 1$ if σ is an odd permutation. Hence, A_n is the kernel of f and so it is a normal subgroup of the domain S_n . Furthermore $S_n/A_n \cong \mathbb{Z}_2$ by the first isomorphism theorem. So by Lagrange's theorem

$$|S_n| = |A_n||S_n/A_n|.$$

Therefore, $|A_n| = n!/2$. That is, there are $n!/2$ many elements in A_n □

Remark. What we have shown in the theorem is that, in fact, A_n has index 2 in S_n . In general, if a subgroup H of G has index 2, then H is normal in G . (Since $[G : H] = 2$, there is an element $g \in G - H$, so that $gH \cap H = \emptyset$ and thus $gH = Hg$).