



Math for the people, by the people.

triangle groups

Canonical name	TriangleGroups
Date of creation	2013-03-22 14:25:07
Last modified on	2013-03-22 14:25:07
Owner	rmilson (146)
Last modified by	rmilson (146)
Numerical id	10
Author	rmilson (146)
Entry type	Definition
Classification	msc 20F05
Related topic	ExamplesOfGroups
Defines	von Dyck groups

Consider the following group presentation:

$$\Delta(l, m, n) = \langle a, b, c : a^2, b^2, c^2, (ab)^l, (bc)^n, (ca)^m \rangle$$

where $l, m, n \in \mathbb{N}$.

A group with this presentation corresponds to a triangle; roughly, the generators are reflections in its sides and its angles are $\pi/l, \pi/m, \pi/n$.

Denote by $D(l, m, n)$ the subgroup of <http://planetmath.org/Cosetindex> 2 in $\Delta(l, m, n)$, corresponding to preservation of of the triangle.

The $D(l, m, n)$ are defined by the following presentation:

$$D(l, m, n) = \langle x, y : x^l, y^m, (xy)^n \rangle$$

Note that $D(l, m, n) \cong D(m, l, n) \cong D(n, m, l)$, so $D(l, m, n)$ is of the l, m, n .

Arising from the geometrical nature of these groups,

$$1/l + 1/m + 1/n > 1$$

is called the *spherical case*,

$$1/l + 1/m + 1/n = 1$$

is called the *Euclidean case*, and

$$1/l + 1/m + 1/n < 1$$

is called the *hyperbolic case*

Groups either of the form $\Delta(l, m, n)$ or $D(l, m, n)$ are referred to as *triangle groups*; groups of the form $D(l, m, n)$ are sometimes referred to as *von Dyck groups*.