



planetmath.org

Math for the people, by the people.

injection can be extended to isomorphism

Canonical name	InjectionCanBeExtendedToIsomorphism
Date of creation	2013-03-22 18:56:50
Last modified on	2013-03-22 18:56:50
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	10
Author	pahio (2872)
Entry type	Theorem
Classification	msc 20A05
Classification	msc 03E20
Related topic	Restriction
Related topic	Cardinality

Theorem. If f is an injection from a set S into a group G , then there exist a group H containing S and a group isomorphism $\varphi: H \rightarrow G$ such that $\varphi|_S = f$.

Proof. Let M be a set such that $\text{card}(M) \geq \text{card}(G)$. Because $\text{card}(f(S)) = \text{card}(S)$, we have $\text{card}(M \setminus S) \geq \text{card}(G \setminus f(S))$, and therefore there exists an injection

$$\psi: G \setminus f(S) \rightarrow M \setminus S$$

(provided that $G \setminus f(S) \neq \emptyset$; otherwise the mapping $f: S \rightarrow G$ would be a bijection). Define

$$H := S \cup \psi(G \setminus f(S)),$$

$$\varphi(h) := \begin{cases} f(h) & \text{for } h \in S, \\ \psi^{-1}(h) & \text{for } h \in H \setminus S. \end{cases}$$

Then apparently, $\varphi: H \rightarrow G$ is a bijection and $\varphi|_S = f$. Moreover, define the binary operation “ $*$ ” of the set H by

$$h_1 * h_2 := \varphi^{-1}(\varphi(h_1) \cdot \varphi(h_2)). \quad (1)$$

We see first that

$$\begin{aligned} (h_1 * h_2) * h_3 &= \varphi^{-1}(\varphi(\varphi^{-1}(\varphi(h_1) \cdot \varphi(h_2))) \cdot \varphi(h_3)) \\ &= \varphi^{-1}((\varphi(h_1) \cdot \varphi(h_2)) \cdot \varphi(h_3)) \\ &= \varphi^{-1}(\varphi(h_1) \cdot (\varphi(h_2) \cdot \varphi(h_3))) \\ &= \varphi^{-1}(\varphi(h_1) \cdot \varphi(\varphi^{-1}(\varphi(h_2) \cdot \varphi(h_3)))) \\ &= h_1 * (h_2 * h_3). \end{aligned}$$

Secondly,

$$h * \varphi^{-1}(e) = \varphi^{-1}(\varphi(h) \cdot \varphi(\varphi^{-1}(e))) = \varphi^{-1}(\varphi(h)) = h,$$

whence $\varphi^{-1}(e)$ is the right identity element of H . Then,

$$h * \varphi^{-1}((\varphi(h))^{-1}) = \varphi^{-1}(\varphi(h) \cdot \varphi(\varphi^{-1}(\varphi(h)^{-1}))) = \varphi^{-1}(e),$$

and accordingly $\varphi^{-1}((\varphi(h))^{-1})$ is the right inverse of h in H . Consequently, $(H, *)$ is a group. The equation (1) implies that

$$\varphi(h_1 * h_2) = \varphi(h_1) \cdot \varphi(h_2),$$

whence φ is an isomorphism from H onto G . Q.E.D.