



Math for the people, by the people.

finite subgroup

Canonical name	FiniteSubgroup
Date of creation	2013-03-22 18:57:02
Last modified on	2013-03-22 18:57:02
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	5
Author	pahio (2872)
Entry type	Theorem
Classification	msc 20A05
Synonym	criterion for finite subgroup
Synonym	finite subgroup criterion

Theorem. A non-empty finite subset K of a group G is a subgroup of G if and only if

$$xy \in K \quad \text{for all } x, y \in K. \quad (1)$$

Proof. The condition (1) is apparently true if K is a subgroup. Conversely, suppose that a nonempty finite subset K of the group G satisfies (1). Let a and b be arbitrary elements of K . By (1), all powers of b belong to K . Because of the finiteness of K , there exist positive integers r, s such that

$$b^r = b^s, \quad r > s+1.$$

By (1),

$$K \ni b^{r-s-1} = b^{r-s}b^{-1} = eb^{-1} = b^{-1}.$$

Thus also $ab^{-1} \in K$, whence, by the theorem of the <http://planetmath.org/node/1045>parent entry, K is a subgroup of G .

Example. The multiplicative group G of all nonzero complex numbers has the finite multiplicative subset $\{1, -1, i, -i\}$, which has to be a subgroup of G .