

Will Kobe Bryant Make His Next Shot: Linear Discriminant Analysis and Logistic Regression using R

Paul Adams

Reannan McDaniel

Jeff Nguyen

Southern Methodist University

23 November 2019

Abstract:

This project investigates the correlation between multiple potential explanatory variables and Kobe Bryant's ability to make a shot while playing for the NBA team Los Angeles Lakers using data gathered from 1996-2015.

Exploratory Data Analysis

In preparation for analysis, derived data was needed to simplify overly complicated variables for the model. The over-complication is with respect to the availability of other predictor variables.

Next, we removed one-level factors. These will never change so are not useful to the model; including can cause issues with model sensitivity since linear trajectories will be down-weighted. Therefore, their significance will be lessened by the constant state of the additional parameters. While this is may not be significant, it is not conducive to model quality.


Outlier Check

A brief outlier check indicated a 2PT Field Goal was noted from the 3PT (3-point) range. Regardless of the actual score, the location matters more. Therefore, after confirming these values to be within 3PT-range, all shots will be encoded as 3-point once beyond 300 inches.

Addressing Multicollinearity: Correlation Plot for Visual Data Exploration

To address multicollinearity among quantitative predictor variables, we used a correlation matrix.

Correlation of Quantitative Predictor Variables, Before Variable Elimination



Post-Correlation Plot Variable Elimination

Following our correlation plot, we decided to eliminate some collinear terms. However, some of the collinearity is useful to capture the instances where the terms are unique. For example, combined_shot_type (factor variable) is collinear with shot_distance (quantitative variable), but it also accounts for the method Kobe may use to make a shot. For example, distance may be relatively the same between 10 and 11 feet, but the factor levels used to derive their short or far indications may differ. This difference could be whether Kobe makes a potentially more accurate heel-planted shot or if he is forced to lean forward and take a riskier shot at basket; the difference in distance may only be one foot, but the difference in technique could measure significant relative to the odds of success.

Addressing Multicollinearity: Correlation Matrix for Numerical Analysis

Following the removal of the most obvious collinear terms visually performing a correlation plot analysis, a correlation matrix for analyzing the remaining results. Collinear quantitative data was preliminarily removed following correlation plot analysis to desaturate the model to an extent that allows more distinction among significance measures for terms in the correlation matrix.

```
##           row      column      cor p
## 26      loc_y shot_distance 0.8181240 0
## 58      game_id      shot_id 0.7612322 0
## 56      recId      shot_id 0.6917176 0
## 44 shot_distance      shot_type 0.6686083 0
## 41      loc_y      shot_type 0.6066234 0
## 2       recId      game_id 0.5270915 0
## 104     attendance     avgnoisedb 0.5109168 0
```

Quadratic Discriminant Analysis

As requested within the requirements of this study, a Linear Discriminant Analysis must be assessed and provided. Discriminant analysis is an operation that compares a categorical response variable against measures of quantitative predictor variables. As a result, analysis for this section is performed on the numerical predictors, which include recId, game_event_id, game_id, loc_x, loc_y, minutes_remaining, seconds_remaining, shot_distance, shot_made_flag, shot_type, game_date, shot_id, attendance, arena_temp, avgnoisedb, controlling collinearity by eliminating a member of each collinear pair prior to model development.

Linear Discriminant Analysis requires a linear boundary between the predictor variables, respective of the response. If the boundary between predictors and response is not linear, Quadratic Discriminant Analysis must be used. Wilks' Lambda distribution is used to assess the nature of boundary linearity, which can be used for discriminant analysis. However, because of the large dimensions of the data set analyzed in this study, an approximation of Wilks' Lambda must be used. Bartlett's Test is an approximation of Wilks' Lambda that can be used for models with large dimensions by applying a measure against the Chi-Square distribution. Provided is a test statistic and p-value.

Bartlett's Test:

	Chi.Square.Statistic	Degrees.of.Freedom	Wilks..Lambda	p.value
Wilks' Lambda	1036.917	14	0.9511137	p < 0.0001

Bartlett's Test of this data set yielded a significant p-value, where $p < 0.0001$, indicating that the proportion of distribution beyond the derived test statistic is beyond that which could be explained by chance. Therefore, we must reject the null hypothesis that the boundary for analysis is linear; the boundary is non-linear. Thus, an analysis using Quadratic Discriminant Analysis is applied.

```
## mean.kobe.qda.posterior...1.. mean.kobe.qda.posterior...2..
## 1                0.542211                0.457789
## shot_made_flagg proportion
## 1                0    0.542211
## 2                1    0.457789
```

Logistic Model Development using Ordinary Least Squares

A preliminary, manual variable elimination process was performed during the analysis of multicollinear terms in preparation for model development. Below we perform logistic regression using Ordinary Least Squares (OLS). In preparation for the model development, a starting model and a finishing model must be developed to provide the scope of variable selection.

Forward Selection

Forward selection produced a model that produced an Akaike's Information Criterion score of 27,378.

Forward Selection Model:

shot_made_flagg = *shotdistance* + *attendance* + *combinedshottype* + *arenatemp* + *gameeventid* + *secondsremaining* + *shottype* + *gamedate* + *minutesremaining* + *locy* + *shotid*

Forward Selection - Akaike's Information Criterion for Logistic Regression:

```
## Warning in kable_styling(., position = "center"): Please specify format in
## kable. kableExtra can customize either HTML or LaTeX outputs. See https://
## haozhu233.github.io/kableExtra/ for details.
```

Akaikes.Information.Criterion..Foreward
27378.42

Backward Elimination

Backward elimination produced a model that produced an Akaike's Information Criterion score of 27,378.

Backward Elimination Model:

shotmade_flagg = *combinedshottype* + *gameeventid* + *locy* + *minutesremaining* + *secondsremaining* + *shotdistance* + *shottype* + *gamedate* + *shotid* + *attendance* + *arenatemp*

Backward Elmination - Akaike's Information Criterion for Logistic Regression:

Akaikes.Information.Criterion..Backward
27378.42

Stepwise Regression

Stepwise Regression produced a model that produced an Akaike's Information Criterion score of 27,378.

Stepwise Regression Model:

shotmadeflag = *combinedshottype* + *gameeventid* + *locy* + *minutesremaining* + *secondsremaining* + *shotdistance* + *shottype* + *gamedate* + *shotid* + *attendance* + *arenatemp*

Stepwise Regression - Akaike's Information Criterion for Logistic Regression:

```
## Warning in kable_styling(., position = "center"): Please specify format in
## kable. kableExtra can customize either HTML or LaTeX outputs. See https://
## haozhu233.github.io/kableExtra/ for details.
```

Akaikes.Information.Criterion..Stepwise
27378.42