

# Numerical Optimization

Zhentao Shi

February 28, 2018

## Numerical Optimization

Optimization is the key step to carry out econometric extremum estimation. A general optimization problem is formulated as

$$\min_{\theta \in \Theta} f(\theta) \quad \text{s.t.} \quad g(\theta) = 0, h(\theta) \leq 0,$$

where  $f(\cdot)$  is a criterion function,  $g(\theta) = 0$  is an equality constraint, and  $h(\theta) \leq 0$  is an inequality constraint.

Most established numerical optimization algorithms aim at finding a local minimum. However, there is no guarantee to locate the global minimum when multiple local minima exist.

Optimization without the equality and/or inequality constraints is called an *unconstrained* problem; otherwise it is called a *constrained* problem. The constraints can be incorporated into the criterion function via Lagrangian.

## Methods

There are many optimization algorithms in the field of operational research; they are variants of a small handful of main principles.

The fundamental idea for twice-differentiable objective function is the Newton's method. A necessary condition for optimization is  $s(\theta) = \partial f(\theta) / \partial \theta = 0$ .

At an initial trial value  $\theta_0$ , if  $s(\theta_0) \neq 0$ , the research is updated by

$$\theta_{t+1} = \theta_t - (H(\theta_t))^{-1} s(\theta_t)$$

for  $t = 0, 1, \dots$  where  $H(\theta) = \frac{\partial s(\theta)}{\partial \theta}$  is the Hessian matrix. The algorithm iterates until  $|\theta_{t+1} - \theta_t| < \epsilon$  (absolute criterion) and/or  $|\theta_{t+1} - \theta_t| / |\theta_t| < \epsilon$  (relative criterion), where  $\epsilon$  is a small positive number chosen as a tolerance level.

**Newton's Method.** Newton's method seeks the solution to  $s(\theta) = 0$ . Recall that the first-order condition is a necessary condition but not a sufficient condition. We still need to verify the second-order condition to identify whether a root to  $s(\theta)$  is associated to a minimizer or a maximizer, and we compare the values of the minima to decide a global minimum.

It is clear that Newton's method requires computation of the gradient  $s(\theta)$  and the Hessian  $H(\theta)$ . Newton's method converges at quadratic rate, which is fast.

**Quasi-Newton Method.** The most well-known quasi-Newton algorithm is **BFGS**. It avoids explicit calculation of the computationally expensive Hessian matrix. Instead, starting from an initial (inverse) Hessian, it updates the Hessian by an explicit formula motivated from the idea of quadratic approximation.

**Derivative-Free Method.** [Nelder-Mead](#) is a simplex method. It searches a local minimum by reflection, expansion and contraction.

## Implementation

R's optimization infrastructure has been constantly improving. [R Optimization Task View](#) gives a survey of the available CRAN packages.

For general-purpose nonlinear optimization, the package [optimx](#) (Nash 2014) effectively replaces R's default optimization commands. [optimx](#) provides a unified interface for various widely-used optimization algorithms. Moreover, it facilitates comparison among optimization algorithms.

## Example

We use [optimx](#) to solve pseudo Poisson maximum likelihood estimation (PPML). If  $y_i$  is a continuous random variable, it obviously does not follow a Poisson distribution, whose support consists of non-negative integers. However, if the conditional mean model

$$E[y_i|x_i] = \exp(x_i'\beta),$$

is satisfied, we can still use the Poisson regression to obtain a consistent estimator of the parameter  $\beta$  even if  $y_i$  does not follow a conditional Poisson distribution.

If  $Z$  follows a Poisson distribution with mean  $\lambda$ , the probability mass function

$$\Pr(Z = k) = \frac{e^{-\lambda}\lambda^k}{k!}, \text{ for } k = 0, 1, 2, \dots,$$

so that

$$\log \Pr(Y = y|x) = -\exp(x'\beta) + y \cdot x'\beta - \log k!$$

Since the last term is irrelevant to the parameter, the log-likelihood function is

$$\ell(\beta) = \log \Pr(\mathbf{y}|\mathbf{x}; \beta) = -\sum_{i=1}^n \exp(x_i'\beta) + \sum_{i=1}^n y_i x_i'\beta.$$

In addition, it is easy to write the gradient

$$s(\beta) = \frac{\partial \ell(\beta)}{\partial \beta} = -\sum_{i=1}^n \exp(x_i'\beta) x_i + \sum_{i=1}^n y_i x_i.$$

and verify that the Hessian

$$H(\beta) = \frac{\partial^2 \ell(\beta)}{\partial \beta \partial \beta'} = -\sum_{i=1}^n \exp(x_i'\beta) x_i x_i' - \sum_{i=1}^n \exp(x_i'\beta)$$

is negative definite. Therefore,  $\ell(\beta)$  is strictly concave in  $\beta$ .

In economics we have utility maximization and cost minimization. In statistics we have maximum likelihood estimation and minimal least squared estimation. In operational reserach, the default optimization is minimization, not maximization. To follow this convention, here we formulate the *negative* log-likelihood.

```

# Poisson likelihood
poisson.loglik = function( b ) {
  b = as.matrix( b )
  lambda = exp( X %*% b )
  ell = -sum( -lambda + y * log(lambda) )
  return(ell)
}

```

To implement optimization in R, it is recommended to write the criterion as a function of the parameter. Data can be fed inside or outside of the function. If the data is provided as additional arguments, these arguments must be explicit. (In contrast, in Matlab the parameter must be the sole argument for the function to be optimized, and data can only be injected through a nested function.)

```

# implement both BFGS and Nelder-Mead for comparison.

```

```

library(AER)
library(numDeriv)
library(optimx)

## prepare the data
data("RecreationDemand")
y = RecreationDemand$trips
X = with(RecreationDemand, cbind(1, income))

## estimation
b.init = c(0, 1) # initial value
b.hat = optimx(b.init, poisson.loglik, method = c("BFGS", "Nelder-Mead"), control = list(reltol = 1e-07, abstol = 1e-07))
print(b.hat)

```

```

##           p1          p2    value fevals gevals niter convcode
## BFGS      1.177411 -0.09994222 261.1141     99     21     NA        0
## Nelder-Mead 1.167261 -0.09703975 261.1317     53     NA     NA        0
##           kkt1 kkt2 xt看imes
## BFGS          TRUE TRUE    0.03
## Nelder-Mead FALSE TRUE    0.00

```

Given the conditional mean model, nonlinear least squares (NLS) is also consistent in theory. NLS minimizes

$$\sum_{i=1}^n (y_i - \exp(x_i \beta))^2$$

A natural question is: why do we prefer PPML to NLS? My argument is that, PPML's optimization for the linear index is globally convex, while NLS is not. It implies that the numerical optimization of PPML is easier and more robust than that of NLS. I leave the derivation of the non-convexity of NLS as an exercise.

In practice no algorithm suits all problems. Simulation, where the true parameter is known, is helpful to check the accuracy of one's optimization routine before applying to an empirical problem,

where the true parameter is unknown. Contour plot is a useful tool to visualize the function surface/manifold in a low dimension.


### Example

```
x.grid = seq(0, 1.8, 0.02)
x.length = length(x.grid)
y.grid = seq(-0.5, 0.2, 0.01)
y.length = length(y.grid)

z.contour = matrix(0, nrow = x.length, ncol = y.length)

for (i in 1:x.length) {
  for (j in 1:y.length) {
    z.contour[i, j] = poisson.loglik(c(x.grid[i], y.grid[j]))
  }
}

contour(x.grid, y.grid, z.contour, 20)
```



For problems that demand more accuracy, third-party standalone solvers can be invoked via interfaces to R. For example, we can access [NLOpt](#) through the packages [nloptr](#). However, standalone solvers usually have to be compiled and configured. These steps are often not as straightforward as installing commercial Windows software.

NLopt offers an [extensive list of algorithms](#).

## Example

We first carry out the Nelder-Mead algorithm in NLOPT.

```
library(nloptr)
## optimization with Nloptr

opts = list(algorithm = "NLOPT_LN_NELDERMEAD", xtol_rel = 1e-07, maxeval = 500)

res_NM = nloptr(x0 = b.init, eval_f = poisson.loglik, opts = opts)
print(res_NM)

##
## Call:
## nloptr(x0 = b.init, eval_f = poisson.loglik, opts = opts)
##
##
## Minimization using NLopt version 2.4.2
##
## NLopt solver status: 4 ( NLOPT_XTOL_REACHED: Optimization stopped because
## xtol_rel or xtol_abs (above) was reached. )
##
## Number of Iterations.....: 118
## Termination conditions:  xtol_rel: 1e-07 maxeval: 500
## Number of inequality constraints:  0
## Number of equality constraints:    0
## Optimal value of objective function: 261.114078295329
## Optimal value of controls: 1.177397 -0.09993984

# 'SLSQP' is indeed the BFGS algorithm in NLopt, though 'BFGS' doesn't
# appear in the name
opts = list(algorithm = "NLOPT_LD_SLSQP", xtol_rel = 1e-07)
```

To invoke BFGS in NLOPT, we must code up the gradient  $s(\beta)$ , as in the function `poisson.log.grad()` below.

```
poisson.loglik.grad = function(b) {
  b = as.matrix(b)
  lambda = exp(X %*% b)
  ell = -colSums(-as.vector(lambda) * X + y * X)
  return(ell)
}
```

We compare the analytical gradient with the numerical gradient to make sure the function is correct.

```
# check the numerical gradient and the analytical gradient
b = c(0, 0.5)
grad(poisson.loglik, b)

## [1] 6542.46 45825.40
```

```
poisson.loglik.grad(b)
```

```
##           income  
## 6542.46 45825.40
```

With the function of gradient, we are ready for BFGS.

```
res_BFGS = nloptr(x0 = b.init, eval_f = poisson.loglik, eval_grad_f = poisson.loglik.grad,  
  opts = opts)  
print(res_BFGS)
```

```
##  
## Call:  
##  
## nloptr(x0 = b.init, eval_f = poisson.loglik, eval_grad_f = poisson.loglik.grad,  
##      opts = opts)  
##  
##  
## Minimization using NLOpt version 2.4.2  
##  
## NLOpt solver status: 4 ( NLOPT_XTOL_REACHED: Optimization stopped because  
## xtol_rel or xtol_abs (above) was reached. )  
##  
## Number of Iterations.....: 38  
## Termination conditions:  xtol_rel: 1e-07  
## Number of inequality constraints:  0  
## Number of equality constraints:    0  
## Optimal value of objective function: 261.114078295329  
## Optimal value of controls: 1.177397 -0.09993984
```

## Constrained Optimization

- `optimx` can handle simple box-constrained problems.
- `constrOptim` can handle linear constrained problems.
- Some algorithms in `nloptr`, for example, `NLOPT_LD_SLSQP`, can handle nonlinear constrained problems.
- `Rdonlp2` is an alternative for general nonlinear constrained problems. `Rdonlp2` is a package offered by `Rmetric` project. It can be installed by `install.packages("Rdonlp2", repos="http://R-Forge.R-project.org")`

## Convex Optimization

If a function is convex in its argument, then a local minimum is a global minimum. Convex optimization is particularly important in high-dimensional problems. The readers are referred to Boyd and Vandenberghe (2004) for an accessible comprehensive treatment. They claim that “convex optimization is technology; all other optimizations are arts.” This is true to some extent.

### Example

- linear regression model MLE
- Lasso (Su, Shi, and Phillips 2016)
- Relaxed empirical likelihood (Shi 2016).

A class of common convex optimization can be reliably implemented in R. [Rmosek](#) is an interface in R to access [Mosek](#). [Mosek](#) is a high-quality commercial solver dedicated to convex optimization. It offers free academic licenses. ([Rtools](#) is a prerequisite to install [Rmosek](#).)

```
require(Rmosek)
lo1 <- list()
lo1$sense <- "max"
lo1$c <- c(3, 1, 5, 1)
lo1$A <- Matrix(c(3, 1, 2, 0, 2, 1, 3, 1, 0, 2, 0, 3), nrow = 3, byrow = TRUE,
  sparse = TRUE)
lo1$bc <- rbind(blc = c(30, 15, -Inf), buc = c(30, Inf, 25))
lo1$bx <- rbind(blx = c(0, 0, 0, 0), bux = c(Inf, 10, Inf, Inf))
r <- mosek(lo1)
```

A survey paper can be found here with [some recent development in convex optimization in econometrics](#).

## References

- Boyd, Stephen, and Lieven Vandenberghe. 2004. *Convex Optimization*. Cambridge university press.
- Nash, John C. 2014. "On Best Practice Optimization Methods in R." *Journal of Statistical Software* 60 (2): 1–14.
- Shi, Zhentao. 2016. "Econometric Estimation with High-Dimensional Moment Equalities." *Journal of Econometrics* 195 (1). Elsevier: 104–19.
- Su, Liangjun, Zhentao Shi, and Peter CB Phillips. 2016. "Identifying Latent Structures in Panel Data." *Econometrica* 84 (6). Wiley Online Library: 2215–64.