

# Spontaneous Droplet Jump with Electro-bouncing

Erin S. Schmidt<sup>1†</sup>, and M. W. Weislogel<sup>1</sup>

<sup>1</sup>Department of Mechanical Engineering, Portland State University, Portland, OR 97211, USA

(Received xx; revised xx; accepted xx)

We investigate the dynamics of water droplet jumps from superhydrophobic surfaces in the presence of an electric field during a step reduction in gravity level. In the brief free-fall environment of a drop tower, when a strong non-homogeneous electric field (with a measured strength between 0.39 and 2.36 kV/cm) is imposed, body forces acting on the jumped droplets are primarily supplied by polarization stress and Coulombic attraction instead of gravity. The droplet charge, measured to be on the order of  $2.3 \cdot (10^{-11})$  C, originates by electro-osmosis of charged species at the (PTFE coated) hydrophobic surface interface. This electric body force leads to a droplet bouncing behavior similar to well-known phenomena in 1-g, though occurring for larger drops  $\sim 0.1$  mL for a given range of impact Weber numbers,  $We < 20$ . In 1-g, for  $We > 0.4$ , impact recoil behavior on a super-hydrophobic surface is normally dominated by damping from contact line hysteresis and by air-layer interactions. However, in the strong electric field, the droplet bounce dynamics additionally include electrohydrodynamic effects on wettability and Cassie-Wenzel transition. This is qualitatively discussed in terms of coefficients of restitution and trends in contact time.

## 1. Introduction

† Email address for correspondence: esch2@pdx.edu