

Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials

Loïc Foissy

*Fédération de Recherche Mathématique du Nord Pas de Calais FR 2956
Laboratoire de Mathématiques Pures et Appliquées Joseph Liouville
Université du Littoral Côte d'Opale-Centre Universitaire de la Mi-Voir
50, rue Ferdinand Buisson, CS 80699, 62228 Calais Cedex, France*

Email: foissy@lmpa.univ-littoral.fr

Abstract

To any poset or quasi-poset is attached a lattice polytope, whose Ehrhart polynomial we study from a Hopf-algebraic point of view. We use for this two interacting bialgebras on quasi-posets. The Ehrhart polynomial defines a Hopf algebra morphism taking its values in $\mathbb{Q}[X]$; we deduce from the interacting bialgebras an algebraic proof of the duality principle, a generalization and a new proof of a result on B-series due to Whright and Zhao, using a monoid of characters on quasi-posets, and a generalization of Faulhaber's formula.

We also give non-commutative versions of these results: polynomials are replaced by packed words. We obtain in particular a non-commutative duality principle.

Keywords. Ehrhart polynomials; Quasi-posets; Characters monoids; Interacting bialgebras

AMS classification. 16T30; 06A11

Contents

1	Bialgebras in cointeraction	5
1.1	Definition	5
1.2	Monoids actions	6
1.3	Polynomial morphisms	8
2	Examples from quasi-posets	13
2.1	Definition	13
2.2	First coproduct	14
2.3	Second coproduct	16
2.4	Cointeractions	20
3	Ehrhart polynomials	22
3.1	Definition	22
3.2	Recursive computation of ehr and ehr^{str}	24
3.3	Characterization of quasi-posets by packed words	26
3.4	Link with linear extensions	27

4	Characters associated to ehr and ehr^{str}	28
4.1	The character λ	29
4.2	The character α^{str}	31
4.3	The character α and the duality principle	32
4.4	A link with Bernoulli numbers	34
5	Noncommutative version	35
5.1	Reminders on packed words	35
5.2	Hopf algebra morphisms in WQSym	35
5.3	The non-commutative duality principle	38
5.4	Compatibility with the other product and coproduct	40
5.5	Restriction to posets	44

Introduction

Let P be a lattice polytope, that is to say that all its vertices are in $\mathbb{Z}^n$. The Ehrhart polynomial $ehr_P^{cl}(X)$ is the unique polynomial such that, for all $k \geq 1$, $ehr_P^{cl}(k)$ is the number of points in $\mathbb{Z}^n \cap kP$, where kP is the image of P by the homothety of center 0 and ratio k . For example, if S is the square $[0, 1]^n$ and T is the triangle of vertices $(0, 0)$, $(1, 0)$ and $(1, 1)$:

$$ehr_S^{cl}(X) = (X + 1)^2, \quad ehr_T^{cl}(X) = \frac{(X + 1)(X + 2)}{2}.$$

These polynomials satisfy the reciprocity principle: for all $k \geq 1$, $(-1)^{\dim(P)} ehr^{cl}(-k)$ is the number of points of $\mathbb{Z}^n \cap k\dot{P}$, where $\dot{P}$ is the interior of P . For example:

$$ehr_S^{cl}(-X) = (X - 1)^2, \quad ehr_T^{cl}(-X) = \frac{(X - 1)(X - 2)}{2}.$$

We refer to [2] for general results on Ehrhart polynomials.

It turns out that these polynomials appear in the theory of B-series (B for Butcher [4]), as explained in [3, 6]. We now consider rooted trees:

$$\bullet; \quad \mathbf{1}; \quad \mathbf{V}, \mathbf{I}; \quad \mathbf{W}, \mathbf{V}, \mathbf{Y}, \mathbf{I}; \quad \mathbf{W}, \mathbf{V}, \mathbf{U}, \mathbf{V}, \mathbf{V}, \mathbf{V}, \mathbf{W}, \mathbf{Y}, \mathbf{I}, \mathbf{I}; \dots$$

If t is a rooted tree, we orient its edges from the root to the leaves. If i, j are two vertices of t , we shall write $i \xrightarrow{t} j$ if there is an edge from i to j in t .

To any rooted tree t , whose vertices are indexed by $1 \dots n$, we associate a lattice polytope $pol(t)$ in a following way:

$$pol(t) = \left\{ (x_1, \dots, x_n) \in [0, 1]^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i \leq x_j) \right\}$$

For example, if $t = \mathbf{1}$, indexed as $\mathbf{1}_1^2$, then $pol(t) = T$.

We can consider the Ehrhart polynomial $ehr_{pol(t)}^{cl}(X)$, which we shall simply denote by $ehr_t^{cl}(X)$: for all $k \geq 1$,

$$ehr_t^{cl}(k) = \# \left\{ (x_1, \dots, x_n) \in \{0, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i \leq x_j) \right\}.$$

Note that ehr_t^{cl} does not depend on the indexation of the vertices of t . By the duality principle:

$$(-1)^n ehr_t^{cl}(-k) = \# \left\{ (x_1, \dots, x_n) \in \{1, \dots, k-1\}^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i < x_j) \right\}.$$

A B-series is a formal series indexed by rooted trees, of the form:

$$\sum_t a_t \frac{t}{\text{aut}(t)} = a_{\bullet} \bullet + a_{\mathbf{1}} \mathbf{1} + a_{\mathbf{V}} \frac{\mathbf{V}}{2} + a_{\mathbf{1}^2} \mathbf{1}^2 + \dots,$$

where $\text{aut}(t)$ is the number of automorphisms of t . The following B-series is of special importance in numerical analysis:

$$E = \sum_t \frac{1}{t!} \frac{t}{\text{aut}(t)} = \bullet + \frac{1}{2} a_{\mathbf{1}} \mathbf{1} + \frac{1}{3} \frac{\mathbf{V}}{2} + \frac{1}{6} \mathbf{1}^2 + \dots,$$

where $t!$ is the tree factorial (see definition 33). This series is the formal solution of an ordinary differential equation, describing the flow of a vector field. The set of B-series is given a group structure by a substitution operation, which is dually represented by the contraction-extraction coproduct defined in [5]. The inverse of E is called the backward error analysis:

$$E^{-1} = \sum_t \lambda_t \frac{t}{\text{aut}(t)!}.$$

Wright and Zhao [19] proved that these coefficients λ_t are related to Ehrhart polynomials:

$$\lambda_t = (-1)^{|t|} \frac{d \, \text{ehr}_t^{\text{cl}}(X)}{dX} \Big|_{X=-1}.$$

We shall in this text study Ehrhart polynomial attached to quasi-posets in a combinatorial Hopf-algebraic way. A quasi-poset P is a pair $(A, \leq_P)$, where A is a finite set and $\leq_P$ is a reflexive and transitive relation on A . The isoclasses of quasi-posets are represented by their Hasse graphs:

$$1; \quad \bullet; \quad \bullet\bullet, \mathbf{1}, \bullet_2; \quad \dots, \bullet\mathbf{1}, \bullet\bullet_2, \mathbf{V}, \mathbf{\Lambda}, \mathbf{1}^2, \mathbf{1}_2, \bullet_3; \dots$$

In particular, rooted trees can be seen as quasi-posets. For any quasi-poset $P = (\{1, \dots, n\}, \leq_P)$, the polytope associated to P is:

$$\text{pol}(P) = \{(x_1, \dots, x_n) \in [0, 1]^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j) \implies (x_i \leq x_j)\}.$$

We put $\text{ehr}_P(X) = \text{ehr}_{\text{pol}(P)}^{\text{cl}}(X-1)$; note the translation by -1 , which will give us objects more suitable to our purpose. In other words, for all $k \geq 1$:

$$\text{ehr}_P(k) = \#\{(x_1, \dots, x_n) \in \{1, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j) \implies (x_i \leq x_j)\}.$$

We also define a polynomial $\text{ehr}_P^{\text{str}}(X)$ such that for all $k \geq 1$:

$$\text{ehr}_P^{\text{str}}(k) = \#\{(x_1, \dots, x_n) \in \{1, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j \text{ and not } j \leq_P i) \implies (x_i < x_j)\}.$$

See definition 21 and proposition 22 for more details. These polynomials can be inductively computed, with the help of the minimal elements of P (proposition 25).

We shall consider two products m and $\downarrow$, and two coproducts Δ and δ on the space $\mathcal{H}_{\mathbf{qp}}$ generated by isoclasses of quasi-posets. The coproduct Δ , defined in [10, 11] by restriction to open and closed sets of the topologies associated to quasi-posets, makes $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ a graded, connected Hopf algebra, and $(\mathcal{H}_{\mathbf{qp}}, \downarrow, \Delta)$ an infinitesimal bialgebra; the coproduct δ , defined in [8] by an extraction-contraction operation, makes $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ a bialgebra. Moreover, δ is also a right coaction of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ over $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$, and $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ becomes a Hopf algebra in the category of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ -comodules, which we summarize telling that $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$

are two bialgebras in cointeraction (definition 1). For example, the bialgebras $(\mathbb{K}[X], m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ where m is the usual product of $\mathbb{K}[X]$ and Δ, δ are the coproducts defined by

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X,$$

are two cointeracting bialgebras.

Ehrhart polynomials $ehr_P(X)$ and $ehr_P^{str}(X)$ can now be seen as maps from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$, and both are Hopf algebra morphisms from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$ (theorem 23); we shall prove in theorem 37 that ehr^{str} is the unique morphism from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$ compatible with both bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$ and $\mathbb{K}[X]$. Using the cointeraction between the two bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$, we show that the monoid $M_{\mathbf{qp}}$ of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ acts on the set $E_{\mathcal{H}_{\mathbf{qp}} \rightarrow \mathbb{K}[X]}$ of Hopf algebra morphisms from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $\mathbb{K}[X]$ (lemma 5). Moreover, there exists a particular homogeneous morphism $\phi_0 \in E_{\mathcal{H}_{\mathbf{qp}} \rightarrow \mathbb{K}[X]}$ such that for all quasi-poset P :

$$\phi_0(P) = \lambda_P X^{cl(P)} = \frac{\mu_P}{cl(P)!} X^{cl(P)},$$

where μ_P is the number of heap-orderings of P and $cl(P)$ is the number of equivalence classes of the equivalence associated to the quasi-order of P (proposition 32). This formula simplifies if P is a rooted tree: in this case,

$$\phi_0(P) = \frac{1}{P!} X^{|P|}.$$

We prove that there exist characters α and α^{str} in $M_{\mathbf{qp}}$, such that for any quasi-poset P :

$$ehr_P(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim} X^{cl(\sim)}, \quad ehr_P^{str}(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim}^{str} X^{cl(\sim)},$$

where the sum is over a certain family of equivalence relations $\sim$ on the set of vertices of V , $P|\sim$ is a restriction operation and $P/\sim$ is a contraction operation. Applied to corollas, this gives Faulhaber's formula. We prove that α^{str} is the inverse of the character λ associated to ϕ_0 (theorem 37), which is a generalization, as well as a Hopf-algebraic proof, of Wright and Zhao's result. We also give an algebraic proof of the duality principle (theorem 38), and we define a Hopf algebra automorphism $\theta : (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \longrightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ with the help of the cointeraction of the two bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$, satisfying $ehr^{str} \circ \theta = ehr$ (proposition 39).

We propose non-commutative versions of these results in the last section of the paper. Here, (isoclasses of) quasi-posets are replaced by quasi-posets on sets $[n] = \{1, \dots, n\}$, making a Hopf algebra $\mathcal{H}_{\mathbf{qp}}$, in cointeraction with $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$, and $\mathbb{K}[X]$ is replaced by the Hopf algebra of packed words **WQSym** [15]. We define two surjective Hopf algebra morphisms EHR and EHR^{str} from $\mathcal{H}_{\mathbf{qp}}$ to **WQSym** (proposition 41), generalizing ehr and ehr^{str} . The automorphism θ is generalized as a Hopf algebra automorphism $\Theta : \mathcal{H}_{\mathbf{qp}} \longrightarrow \mathcal{H}_{\mathbf{qp}}$, such that $EHR^{str} \circ \Theta = EHR$ (proposition 42), and we formulate a non-commutative duality principle

(theorem 45), and we obtain a commutative diagram of Hopf algebras:

$$\begin{array}{ccccc}
\mathcal{H}_{\mathbf{QP}} & & & & \\
\Theta \downarrow & \nearrow EHR & & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR^{str}} & \mathbf{WQSym} & & \\
\Psi \downarrow & & \Phi_{-1} \downarrow & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR} & \mathbf{WQSym} & & \\
& & & \nearrow H & \\
& & & \mathcal{H}_{\mathbf{qp}} & \nearrow ehr \\
& & & \theta \downarrow & \\
& & & \mathcal{H}_{\mathbf{qp}} & \xrightarrow{ehr^{str}} \mathbb{K}[X] \\
& & & \psi \downarrow & \searrow \phi_{-1} \\
& & & \mathcal{H}_{\mathbf{qp}} & \xrightarrow{ehr} \mathbb{K}[X]
\end{array}$$

The two triangles reflects the properties of morphisms Θ and θ , whereas the two squares are the duality principles.

Acknowledgment. The research leading these results was partially supported by the French National Research Agency under the reference ANR-12-BS01-0017.

Notations. We denote by $\mathbb{K}$ a commutative field of characteristic zero. All the objects (vector spaces, algebra, and so on) in this text are taken over $\mathbb{K}$.

1 Bialgebras in cointeraction

We give in this section some general results on bialgebras in cointeractions. They will be used in the sequel for quasi-posets, leading to Ehrhart polynomials. We shall use them on graphs in order to obtain chromatic polynomials in [9].

1.1 Definition

Definition 1 *Let A and B be two bialgebras. We shall say that A and B are in cointeraction if:*

- B coacts on A , via a map $\rho : \begin{cases} A & \longrightarrow A \otimes B \\ a & \longrightarrow \rho(a) = a_1 \otimes a_0. \end{cases}$
- A is a bialgebra in the category of B -comodules, that is to say:
 - $\rho(1_A) = 1_A \otimes 1_B$.
 - $m_{2,4}^3 \circ (\rho \otimes \rho) \circ \Delta_A = (\Delta_A \otimes Id) \circ \rho$, with:

$$m_{2,4}^3 : \begin{cases} A \otimes B \otimes A \otimes B & \longrightarrow A \otimes A \otimes B \\ a_1 \otimes b_1 \otimes a_2 \otimes b_2 & \longrightarrow a_1 \otimes a_2 \otimes b_1 b_2. \end{cases}$$

Equivalently, in Sweedler's notations, for all $a \in A$:

$$(a^{(1)})_1 \otimes (a^{(2)})_1 \otimes (a^{(1)})_0 (a^{(2)})_0 = (a_1)^{(1)} \otimes (a_1)^{(2)} \otimes a_0.$$

- For all $a, b \in A$, $\rho(ab) = \rho(a)\rho(b)$.
- For all $a \in A$, $(\varepsilon_A \otimes Id) \circ \rho(a) = \varepsilon_A(a)1_B$.

Examples of bialgebras in interaction can be found in [5] (for rooted trees) and in [14] (for various families of graphs). Another example is given by the algebra $\mathbb{K}[X]$, with its usual product m , and the two coproducts defined by:

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X.$$

The bialgebras $(\mathbb{K}[X], m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ are in cointeractions, via the coaction $\rho = \delta$. Identifying $\mathbb{K}[X] \otimes \mathbb{K}[X]$ and $\mathbb{K}[X, Y]$:

$$\Delta(P)(X, Y) = P(X + Y), \quad \delta(P)(X, Y) = P(XY).$$

Remark. If A and B are in cointeraction, the coaction of B on A is an algebra morphism.

Proposition 2 *Let A and B be two bialgebras in cointeraction. We assume that A is a Hopf algebra, with antipode S . Then S is a morphism of B -comodules, that is to say:*

$$\rho \circ S = (S \otimes Id) \circ \rho$$

Proof. We work in the space $End_{\mathbb{K}}(A, A \otimes B)$. As $A \otimes B$ is an algebra and A is a coalgebra, it is an algebra for the convolution product $\circledast$:

$$\forall f, g \in End_{\mathbb{K}}(A, A \otimes B), f \circledast g = m_{A \otimes B} \circ (f \otimes g) \circ \Delta_A.$$

Its unit is:

$$\eta : \begin{cases} A & \longrightarrow A \otimes B \\ a & \longrightarrow \varepsilon(a)1_A \otimes 1_B. \end{cases}$$

We consider three elements in this algebra, respectively ρ , $F_1 = (S \otimes Id) \circ \rho$ and $F_2 = \rho \circ S$. Firstly:

$$\begin{aligned} (F_1 \circledast \rho)(a) &= S((a^{(1)})_1)(a^{(2)})_1 \otimes (a^{(1)})_0(a^{(2)})_0 \\ &= S((a_1)^{(1)})(a_1)^{(2)} \otimes a_0 \\ &= \varepsilon_A(a_1)1_A \otimes a_0 \\ &= \varepsilon_A(a)1_A \otimes 1_B \\ &= \eta(a). \end{aligned}$$

Secondly:

$$\begin{aligned} (\rho \circledast F_2)(a) &= (a^{(1)})_1 S(a^{(2)})_1 \otimes (a^{(1)})_0 (S(a^{(2)}))_0 \\ &= \varepsilon_A(a)(1_A)_1 \otimes (1_A)_0 \\ &= \varepsilon_A(a)1_A \otimes 1_B \\ &= \eta(a). \end{aligned}$$

We obtain that $F_1 \circledast \rho = \rho \circledast F_2 = \eta$, so $F_1 = F_1 \circledast \eta = F_1 \circledast \rho \circledast F_2 = \eta \circledast F_2 = F_2$. $\square$

1.2 Monoids actions

Proposition 3 *Let A and B be two bialgebras in cointeraction, through the coaction ρ . We denote by M_A and M_B the monoids of characters of respectively A and B . Then B acts on A by monoid endomorphisms, via the map:*

$$\leftarrow : \begin{cases} M_A \times M_B & \longrightarrow M_A \\ (\phi, \lambda) & \longrightarrow \phi \leftarrow \lambda = (\phi \otimes \lambda) \circ \rho. \end{cases}$$

Proof. We denote by $*$ the convolution product of M_B and by $\star$ the convolution product of M_A . As $\rho : A \longrightarrow A \otimes B$ is an algebra morphism, $\leftarrow$ is well-defined. Let $\phi \in M_A$, $\lambda_1, \lambda_2 \in M_B$.

$$\begin{aligned} (\phi \leftarrow \lambda_1) \leftarrow \lambda_2 &= (\phi \otimes \lambda_1 \otimes \lambda_2) \circ (\rho \otimes Id) \circ \rho \\ &= (\phi \otimes \lambda_1 \otimes \lambda_2) \circ (Id \otimes \Delta_B) \circ \rho \\ &= \phi \leftarrow (\lambda_1 * \lambda_2). \end{aligned}$$

So $\leftarrow$ is an action. Let $\phi_1, \phi_2 \in M_A$, $\lambda \in M_B$. For all $a \in A$:

$$\begin{aligned} ((\phi_1 \star \phi_2) \leftarrow \lambda)(a) &= (\phi_1 \otimes \phi_2 \otimes \lambda) \circ (\Delta_A \otimes Id) \circ \rho(a) \\ &= (\phi_1 \otimes \phi_2 \otimes \lambda)((a_0)^{(1)} \otimes (a_0)^{(2)} \otimes a_1) \\ &= (\phi_1 \otimes \phi_2 \otimes \lambda)((a^{(1)})_0 \otimes (a^{(2)})_0 \otimes (a^{(1)})_1(a^{(2)})_1) \\ &= \phi_1((a^{(1)})_0) \lambda((a^{(1)})_1) \phi_2((a^{(2)})_0) \lambda((a^{(2)})_1) \\ &= (\phi_1 \leftarrow \lambda)(a^{(1)}) (\phi_2 \leftarrow \lambda)(a^{(2)}) \\ &= ((\phi_1 \leftarrow \lambda) \star (\phi_2 \leftarrow \lambda))(a). \end{aligned}$$

So $\leftarrow$ is an action by monoid endomorphisms. $\square$

Example. We take $A = (\mathbb{K}[X], m, \Delta)$, $B = (\mathbb{K}[X], m, \delta)$ and $\rho = \delta$. We consider the map:

$$ev : \begin{cases} \mathbb{K} & \longrightarrow & \mathbb{K}[X]^* \\ \lambda & \longrightarrow & \begin{cases} \mathbb{K}[X] & \longrightarrow & \mathbb{K} \\ P(X) & \longrightarrow & ev_\lambda(P) = P(\lambda). \end{cases} \end{cases}$$

Then ev is an isomorphism from $(\mathbb{K}, +)$ to $(M_A, \star)$ and from $(\mathbb{K}, \cdot)$ to $(M_B, *)$. Moreover, for all $\lambda, \mu \in \mathbb{K}$:

$$ev_\lambda \leftarrow ev_\mu = ev_{\lambda\mu}.$$

Proposition 4 *Let A and B be two bialgebras in cointeraction, through the coaction ρ .*

1. *Let H be a bialgebra. We denote by M_B the monoid of characters of B and by $E_{A \rightarrow H}$ the set of bialgebra morphisms from A to H . Then M_B acts on $E_{A \rightarrow H}$ via the map:*

$$\leftarrow : \begin{cases} E_{A \rightarrow H} \times M_B & \longrightarrow & E_{A \rightarrow H} \\ (\phi, \lambda) & \longrightarrow & \phi \leftarrow \lambda = (\phi \otimes \lambda) \circ \rho \end{cases}$$

2. *Let H_1 and H_2 be two bialgebras and let $\theta : H_1 \longrightarrow H_2$ be a bialgebra morphism. For all $\phi \in E_{A \leftarrow H_1}$, for all $\lambda \in M_B$, in $E_{A \leftarrow H_2}$:*

$$\theta \circ (\phi \leftarrow \lambda) = (\theta \circ \phi) \leftarrow \lambda.$$

3. *if $\lambda, \mu \in M_B$, in $E_{A \rightarrow A}$:*

$$(Id \leftarrow \lambda) \circ (Id \leftarrow \mu) = Id \leftarrow (\lambda * \mu).$$

The following map is an injective monoid morphism:

$$\begin{cases} (M_B, *) & \longrightarrow & (E_{A \rightarrow A}, \circ) \\ \lambda & \longrightarrow & Id \leftarrow \lambda. \end{cases}$$

Proof. 1. For all $\phi \in E_{A \leftarrow B}$, $\lambda \in M_B$, $\phi \leftarrow \lambda : A \longrightarrow H \otimes \mathbb{K} = H$. As ϕ , λ and ρ are algebra morphisms, by composition $\phi \leftarrow \lambda$ is an algebra morphism. Let $a \in A$.

$$\begin{aligned}
\Delta_H(\phi \leftarrow \lambda(a)) &= \Delta_H(\phi(a_0)\lambda(a_1)) \\
&= \lambda(a_1)\Delta_H \circ \phi(a_1) \\
&= \lambda(a_1)\phi(a_0)^{(1)} \otimes \phi(a_0)^{(2)} \\
&= \lambda(a_1)\phi((a_0)^{(1)}) \otimes \phi((a_0)^{(2)}) \\
&= \lambda((a^{(1)})_1(a^{(2)})_1)\phi((a^{(1)})_0) \otimes \phi((a^{(2)})_0) \\
&= \lambda((a^{(1)})_1)\lambda((a^{(2)})_1)\phi((a^{(1)})_0) \otimes \phi((a^{(2)})_0) \\
&= \phi((a^{(1)})_0)\lambda((a^{(1)})_1) \otimes \phi((a^{(2)})_0)\lambda((a^{(2)})_1) \\
&= \phi \leftarrow \lambda(a^{(1)}) \otimes \phi \leftarrow \lambda(a^{(2)}) \\
&= ((\phi \leftarrow \lambda) \otimes (\phi \leftarrow \lambda)) \circ \Delta_A(a).
\end{aligned}$$

So $\phi \leftarrow \lambda \in E_{A \rightarrow H}$.

Let $\phi \in E_{A \rightarrow H}$. For all $a \in A$, $\phi \leftarrow \eta \circ \varepsilon(a) = \phi(a_0)\varepsilon(a_1) = \phi(a)$. Let $\lambda, \mu \in M_B$.

$$(\phi \leftarrow \lambda) \leftarrow \mu = (\phi \otimes \lambda \otimes \mu) \circ (\rho \otimes Id) \circ \rho = (\phi \otimes \lambda \otimes \mu) \circ (Id \otimes \Delta_B) \circ \rho = \phi \leftarrow (\lambda * \mu).$$

So $\leftarrow$ is indeed an action of M_B on $E_{A \rightarrow H}$.

2. Let $a \in H$.

$$(\theta \circ \phi) \leftarrow \lambda(a) = \theta \circ \phi(a_1)\lambda(a_0) = \theta(\phi(a_1)\lambda(a_0)) = \theta(\phi \leftarrow \lambda(a)) = \theta \circ (\phi \leftarrow \lambda)(a).$$

So $(\theta \circ \phi) \leftarrow \lambda = \theta \circ (\phi \leftarrow \lambda)$.

3. Consequently, if $\lambda, \mu \in M_B$, in $E_{A \rightarrow A}$:

$$(Id \leftarrow \lambda) \circ (Id \leftarrow \lambda) = (Id \leftarrow \lambda) \leftarrow \mu = Id \leftarrow (\lambda * \mu).$$

If $Id \leftarrow \lambda = Id$, then, composing by ε' , we obtain $\varepsilon' * \lambda = \varepsilon'$, so $\lambda = \varepsilon'$. $\square$

Example. We take $A = (\mathbb{K}[X], m, \Delta)$, $B = (\mathbb{K}[X], m, \delta)$ and $\rho = \delta$. In $E_{A \rightarrow A}$, for any $\lambda \in \mathbb{K}$:

$$Id \leftarrow ev_\lambda(X) = ev_\lambda(X)X = \lambda X,$$

so for any $P \in \mathbb{K}[X]$, $(Id \leftarrow ev_\lambda)(P) = P(\lambda X)$. In this case, the monoids $(M_B, *)$ and $(E_{A \rightarrow A}, \circ)$ are isomorphic.

1.3 Polynomial morphisms

In this section, we deal with a family (A, m, Δ, δ) such that:

1. (A, m, Δ) is a graded, connected Hopf algebra. As a graded algebra, it is isomorphic to the symmetric algebra $S(V)$, where V is a graded subspace of A .
2. (A, m, δ) is a bialgebra.
3. (A, m, Δ) and (A, m, δ) are in cointeraction, through the coaction δ .
4. $V_1 = A_1$ has a basis $(g_i)_{i \in I}$ such that:

$$\forall i \in I, \delta(g_i) = g_i \otimes g_i.$$

We shall denote by J the set of sequences $\alpha = (\alpha_i)_{i \in I}$ with a finite support. For all $\alpha \in J$, we put $g_\alpha = \prod_{i \in I} g_i^{\alpha_i}$. These are group-like elements of (A, m, δ) .

5. For all $n \geq 2$, V_n can be decomposed as:

$$V_n = \bigoplus_{i \in I, \alpha \in J} V_n(g_i, g_\alpha),$$

such that for all $x \in V_n(g_i, g_\alpha)$:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha \in S(V_1 \oplus \dots \oplus V_{n-1})^{\otimes 2}.$$

The counit of (A, m, Δ) will be denoted by ε and the counit of (A, m, δ) by ε' . We denote by M_B the monoid of characters of (A, m, δ) .

Remark. If $x \in V_n(g_i, g_\alpha)$, as $\varepsilon'(g_i) = \varepsilon'(g_\alpha) = 1$, necessarily, $\varepsilon'(x) = 0$ and:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha \in \text{Ker}(\varepsilon')^{\otimes 2}.$$

Lemma 5 *Let $\lambda \in M_B$. It has an inverse in M_B if, and only if, for all $i \in I$, $\lambda(g_i) \neq 0$.*

Proof. $\implies$. Let μ be the inverse of λ in M_B . For all $i \in I$, $\lambda * \mu(g_i) = \lambda(g_i)\mu(g_i) = \varepsilon'(g_i) = 1$, so $\lambda(g_i) \neq 0$.

$\impliedby$. We define two characters $\mu, \nu \in M_B$ by inductively defining $\mu_n = \mu|_{V_n}$ and $\nu_n = \nu|_{V_n}$. For $n = 1$, we put $\mu_1(g_i) = \nu_1(g_i) = \lambda(g_i)^{-1}$. Let us assume that $g_1, \dots, g_{n-1}$ are already defined, with $n \geq 2$. If $x \in V_n(g_i, g_\alpha)$, we put:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha = \sum x'_k \otimes x''_k \in S(V_1 \oplus \dots \oplus V_{n-1})^{\otimes 2}.$$

Hence, for all k , $\mu(x'_k)$ and $\nu(x''_k)$ are defined. We put:

$$\begin{aligned} \mu_n(x) &= \prod_{i \in I} \frac{1}{\lambda(g_i)^{\alpha_i}} \left(\varepsilon'(x) - \mu(g_i)\lambda(x) - \sum \mu(x'_k)\lambda(x''_k) \right), \\ \nu_n(x) &= \frac{1}{\lambda(g_i)} \left(\varepsilon'(x) - \lambda(x)\nu(g_\alpha) - \sum \lambda(x'_k)\nu(x''_k) \right). \end{aligned}$$

Consequently, $\mu, \nu \in M_B$ and for all $x \in V$, $\mu * \lambda(x) = \lambda * \nu(x) = \varepsilon'(x)$, so $\mu * \lambda = \lambda * \nu = \varepsilon'$, and finally $\mu = \mu * (\lambda * \nu) = (\mu * \lambda) * \nu = \nu$, so λ is invertible in M_B . $\square$

Lemma 6 *Let C be a graded, connected Hopf algebra. We denote by $C_+ = \text{Ker}(\varepsilon_C)$ its augmentation ideal. Let $\lambda : C_+ \rightarrow \mathbb{K}$ be any linear map. There exists a unique coalgebra morphism $\phi : C \rightarrow \mathbb{K}[X]$ such that:*

$$\forall x \in C_+, \frac{d\phi(x)}{dX}(0) = \lambda(x).$$

Moreover:

1. ϕ is homogeneous if, and only if, for all $n \geq 2$, $\lambda(C_n) = (0)$.
2. ϕ is a Hopf algebra morphism if, and only if, for all $x, y \in C_+$, $\lambda(xy) = 0$.

Proof. Let $\pi : \mathbb{K}[X] \rightarrow \text{Vect}(X)$ be the canonical projection. For any $P \in \mathbb{K}[X]$:

$$\pi(P) = \frac{dP}{dX}(0)X.$$

Existence. We define $\phi|_{C_n}$ by induction on n . For $n = 0$, we put $\phi(1) = 1$. Let us assume that $\phi|_{C_0 \oplus \dots \oplus C_{n-1}}$ is defined such that for all $x \in C_0 \oplus \dots \oplus C_{n-1}$,

$$\Delta \circ \phi(x) = (\phi \otimes \phi) \circ \Delta(x), \quad \pi \circ \phi(x) = \lambda(x)X.$$

Let $x \in C_n$. As $\tilde{\Delta}(x) \in (C_1 \oplus \dots \oplus C_{n-1})^{\otimes 2}$, we can consider the element

$$y = (\phi \otimes \phi) \circ \tilde{\Delta}(x) \in \mathbb{K}[X]_+^{\otimes 2}.$$

We put:

$$y = \sum_{i,j \geq 1} a_{i,j} \frac{X^i}{i!} \otimes \frac{X^j}{j!}.$$

Moreover:

$$\begin{aligned} \tilde{\Delta} \otimes Id(y) &= ((\tilde{\Delta} \circ \phi) \otimes \phi) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes \phi \otimes \phi) \circ (\tilde{\Delta} \otimes Id) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes \phi \otimes \phi) \circ (Id \otimes \tilde{\Delta}) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes (\tilde{\Delta} \circ \phi)) \circ \tilde{\Delta}(x) \\ &= (Id \otimes \tilde{\Delta})(y). \end{aligned}$$

Hence:

$$\sum_{i,j,k \geq 1} a_{i+j,k} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \otimes \frac{X^k}{k!} = \sum_{i,j,k \geq 1} a_{ij+k} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \otimes \frac{X^k}{k!}.$$

For all $i, j, k \geq 1$, $a_{i+j,k} = a_{i,j+k}$, so there exist scalars a_n such that for all $i, j \geq 1$, $a_{i,j} = a_{i+j}$. We obtain that:

$$y = \sum_{n \geq 2} a_n \left(\sum_{i,j \geq 1, i+j=n} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \right) = \sum_{n \geq 2} a_n \tilde{\Delta} \left(\frac{X^n}{n!} \right).$$

We then put $\phi(x) = \sum_{n \geq 2} a_n \frac{X^n}{n!} + \lambda(x)X$.

We obtain in this way a coalgebra morphism such that $\pi \circ \phi(x) = \lambda(x)X$ for all $x \in C$.

Unicity. Let $\phi, \psi : C \longrightarrow \mathbb{K}[X]$ be coalgebra morphisms such that $\pi \circ \phi = \pi \circ \psi$. Let us prove that $\phi(x) = \psi(x)$ for all $x \in A_n$, $n \geq 0$, by induction on n . For $n = 0$, as the unique group-like element of $\mathbb{K}[X]$ is 1, $\phi(1) = \psi(1) = 1$. Let us assume the result at all ranks $k < n$. Then, if $x \in A_n$:

$$\tilde{\Delta} \circ \phi(x) = (\phi \otimes \phi) \circ \tilde{\Delta}(x) = (\psi \otimes \psi) \circ \tilde{\Delta}(x) = \tilde{\Delta} \circ \psi(x),$$

so $\phi(x) - \psi(x) \in \text{Ker}(\tilde{\Delta}) = \text{Vect}(X)$. Hence, $0 = \pi \circ \phi(x) - \pi \circ \psi(x) = \phi(x) - \psi(x)$.

1. $\implies$. If ϕ is homogeneous, then, for all $n \geq 2$, $\phi(C_n) \subseteq \text{Vect}(X^n)$, so:

$$\pi \circ \phi(C_n) = \lambda(C_n)X = (0).$$

1. $\Leftarrow$. Let us go back to the construction of ϕ is the *Existence* part. If $n \geq 1$, $x \in C_1$, then $\phi(x) = \lambda(x)X$ is homogeneous of degree 1. If $n \geq 2$, then, by homogeneity, $a_{i,j} = 0$ if $i + j \neq n$, so $a_k = 0$ if $n \neq k$, and $\phi(x) = a_n \frac{X^n}{n!}$: ϕ is homogeneous.

2. $\implies$. Let $x, y \in C_+$. Then $\phi(x), \phi(y) \in \mathbb{K}[X]_+ = X\mathbb{K}[X]$, so $\phi(xy) = \phi(x)\phi(y) \in X^2\mathbb{K}[X]$ and $\pi \circ \phi(xy) = \lambda(xy)X = 0$.

2. $\Leftarrow$. Let us consider $\phi_1 = m \circ (\phi \otimes \phi)$ and $\phi_2 = \phi \circ m$. By composition, they are both coalgebra morphisms from the graded bialgebra $C \otimes C$ to $\mathbb{K}[X]$. Moreover, if $x, y \in C_+$:

$$\begin{aligned} \pi \circ \phi_1(1 \otimes y) &= \pi \circ \phi(y), & \pi \circ \phi_2(1 \otimes y) &= \pi \circ \phi(y), \\ \pi \circ \phi_1(x \otimes 1) &= \pi \circ \phi(x), & \pi \circ \phi_2(x \otimes 1) &= \pi \circ \phi(x), \\ \pi \circ \phi_1(x \otimes y) &= \pi(\phi(x)\phi(y)) = 0, & \pi \circ \phi_2(x \otimes y) &= \lambda(xy)X = 0. \end{aligned}$$

So, for all $z \in (C \otimes C)_+ = (\mathbb{K}1 \otimes C_+) \oplus (C_+ \otimes \mathbb{K}1) \oplus (C_+ \otimes C_+)$, $\pi \circ \phi_1(z) = \pi \circ \phi_2(z)$. By the *Unicity* part, $\phi_1 = \phi_2$, so ϕ is an algebra morphism. $\square$

Remark. If $x \in V_1$, $\phi(x) = \lambda(x)X$.

Theorem 7 *Under the hypotheses 1–5, there exists a unique homogeneous Hopf algebra morphism $\phi_0 : (A, m, \Delta) \rightarrow (\mathbb{K}[X], m, \Delta)$ such that:*

$$\forall x \in A_1, \phi_0(x) = \varepsilon'(x)X.$$

Moreover, there exists a unique character $\lambda_0 \in M_B$, invertible in M_B , such that:

$$\forall n \geq 0, \forall x \in A_n, \phi_0(x) = \lambda_0(x)X^n.$$

Proof. Existence. Let $\lambda \in A^*$ such that $\lambda(x) = \varepsilon'(x)$ if $x \in A_1$ and $\lambda(x) = 0$ if $x \in A_n$, $n \geq 2$. We denote by ϕ_0 the unique coalgebra morphism such that $\frac{d\phi_0(x)}{dX}(0) = \lambda(x)$ for all $x \in A_+$. By the first point of lemma 6, ϕ_0 is homogeneous. As $A_+^2 \subseteq A_{\geq 2}$, by the second point of lemma 6, ϕ_0 is an algebra morphism.

Unicity. If ϕ is such a morphism, by the first point of lemma 6, for all $x \in A_n$, $n \geq 2$, $\pi \circ \phi(x) = 0$; hence, for all $x \in A_+$, $\pi \circ \phi_0 = \pi \circ \phi$. By the unicity in lemma 6, $\phi = \phi_0$.

By homogeneity of ϕ_0 , for all $x \in A_n$, there exists a unique scalar $\lambda_0(x) \in \mathbb{K}$ such that:

$$\phi_0(x) = \lambda_0(x)X^n.$$

If $x \in A_m$, $y \in A_n$, $xy \in A_{m+n}$ and then:

$$\phi(xy) = \lambda_0(xy)X^{n+m} = \phi(x)\phi(y) = \lambda_0(x)\lambda_0(y)X^{n+m},$$

so $\lambda_0 \in M_B$. For all $i \in I$, $\phi(g_i) = \varepsilon'(g_i)X = X$, so $\lambda_0(g_i) = 1$. By lemma 5, λ_0 is an invertible element of M_B . $\square$

Theorem 8 *Under the hypotheses 1–5, the following map is a bijection:*

$$\theta : \begin{cases} M_B & \longrightarrow E_{A \rightarrow \mathbb{K}[X]} \\ \lambda & \longrightarrow \phi_0 \leftarrow \lambda. \end{cases}$$

Moreover, if $\phi = \theta(\lambda)$, with $\lambda \in M_B$, then for all $x \in V$,

$$\lambda(x) = \frac{d\phi(x)}{dX}(0).$$

Proof. Let $\lambda \in M_B$, and $\phi = \phi_0 \leftarrow \lambda$. For all $i \in I$, $\phi(g_i) = X\lambda(g_i)$, so $\frac{d\phi(g_i)}{dX}(0) = \lambda(g_i)$. If $n \geq 2$ and $x \in V_n(g_i, g_\alpha)$, we put:

$$\delta(x) = g_i \otimes x + \sum x'_k \otimes x''_k,$$

with for all k , x'_k homogeneous of degree ≥ 2 or homogeneous of degree 1, with $\varepsilon'(x'_k) = 0$. For all k , $\pi \circ \phi_0(x'_k) = 0$. We obtain:

$$\pi \circ \phi(x) = \pi \left(\lambda(x)X + \sum \phi_0(x'_k)\lambda(x''_k) \right) = \lambda(x)X.$$

By the unicity in lemma 6, ϕ is injective. If $\psi \in E_{A \rightarrow \mathbb{K}[X]}$, we define $\lambda \in M_B$ by:

$$\forall x \in V, \lambda(x) = \frac{d\psi(x)}{dX}(0).$$

We put $\phi = \phi_0 \leftarrow \lambda$. Then for all $x \in V$, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0) = \lambda(x)$. If $x \in A_+^2$, by the second point of lemma 6, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0) = 0$. Finally, as V generates A , $A_+ = V + A_+^2$, and for all $x \in A_+$, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0)$. By the unicity in lemma 6, $\phi = \psi$, so θ is surjective. $\square$

Corollary 9 *Under the hypotheses 1–5, for any $\mu \in M_B$, there exists a unique Hopf algebra morphism $\phi : A \rightarrow \mathbb{K}[X]$, such that:*

$$\forall x \in A, \phi(x)(1) = \mu(1).$$

This morphism is: $\phi_0 \leftarrow (\lambda_0^{-1} * \mu)$.*

Proof. Let ϕ be a Hopf algebra morphism from A to $\mathbb{K}[X]$. By theorem 8, there exists $\lambda \in M_B$ such that $\phi = \phi_0 \leftarrow \lambda$. Let $x \in A$. We write $x = \sum x'_k \otimes x''_k$, with, for all k , x'_k homogeneous of degree n_k . Then:

$$\phi(x)(1) = \left(\sum \lambda_0(x'_k)X^{n_k}\lambda(x''_k) \right)_{|X=1} = \sum \lambda_0(x'_k)\lambda(x''_k) = \lambda_0 * \lambda(x).$$

So ϕ satisfies the required conditions if, and only if, $\lambda_0 * \lambda = \mu$, if, and only if, $\lambda = \lambda_0^{*-1} * \mu$, as λ_0 is invertible. So such a ϕ exists and is unique. $\square$

Corollary 10 *Under the hypotheses 1–5, there exists a unique morphism $\phi_1 : A \rightarrow \mathbb{K}[X]$ such that:*

1. ϕ_1 is a Hopf algebra morphism from (A, m, Δ) to $(\mathbb{K}[X], m, \Delta)$.
2. ϕ_1 is a bialgebra morphism from (A, m, δ) to $(\mathbb{K}[X], m, \delta)$.

Moreover, $\phi_1 = \phi_0 \leftarrow \lambda_0^{*-1}$ and, for all $x \in A$, $\phi_1(x)(1) = \varepsilon'(x)$.

Proof. *Unicity.* If such a ϕ_1 exists, then for all $x \in A$, $\phi_1(x)(1) = \varepsilon' \circ \phi_1(x) = \varepsilon'(x)$. By corollary 9, $\phi_1 = \phi_0 \leftarrow (\lambda_0^{*-1} * \varepsilon') = \phi_0 \leftarrow \lambda_0^{*-1}$.

Existence. Let ϕ_1 be the unique Hopf algebra morphism in $E_{A \rightarrow \mathbb{K}[X]}$ such that for all $x \in A$, $\phi_1(x)(1) = \varepsilon'(x)$. Recall that we identify $\mathbb{K}[X] \otimes \mathbb{K}[X]$ with $\mathbb{K}[X, Y]$; for all $P \in \mathbb{K}[X]$:

$$\Delta(P)(X, Y) = P(X + Y), \quad \delta(P)(X, Y) = P(XY).$$

Let us fix $x \in A$. Let us prove that for all $k, l \in \mathbb{N}^*$, $\delta \circ \phi_1(x)(k, l) = (\phi_1 \otimes \phi_1) \circ \Delta(x)(k, l)$ by induction on k . First, observe that $\delta \circ \phi_1(x)(k, l) = \phi_1(x)(kl)$. If $k = 1$, then:

$$(\phi_1 \otimes \phi_1) \circ \Delta(x)(1, l) = \phi_1((\varepsilon' \otimes Id) \circ \Delta(x))(l) = \phi_1(x)(l).$$

Let us assume the result at rank k . Then:

$$\begin{aligned}
(\phi_1 \otimes \phi_1) \circ \Delta(x)(k+1, l) &= (\Delta \otimes Id) \circ (\phi_1 \otimes \phi_1) \circ \delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1) \circ (\Delta \otimes Id) \circ \delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1) \circ m_{2,4}^3 \circ (\delta \otimes \delta) \circ \Delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1 \otimes \phi_1) \circ (\delta \otimes \delta) \circ \Delta(x)(k, l, 1, l) \\
&= (\phi_1 \otimes \phi_1) \circ \Delta(x)(kl, l) \\
&= \Delta \circ \phi_1(x)(kl, l) \\
&= \phi_1(x)(kl + l) \\
&= \phi_1(x)((k+1)l).
\end{aligned}$$

So the result is true for all $k, l \geq 1$. Hence, $(\phi_1 \otimes \phi_1) \circ \delta(x) = \delta \circ \phi_1(x)$: ϕ_1 is a bialgebra morphism from (A, m, δ) to $(\mathbb{K}[X], m, \delta)$. $\square$

2 Examples from quasi-posets

2.1 Definition

Definition 11 1. Let A be a set finite set. A quasi-order on A is a transitive, reflexive relation $\leq$ on A . If $\leq$ is a quasi-order on A , we shall say that $(A, \leq)$ is a quasi-poset. If P is a quasi-poset:

(a) Its isoclass is denoted by $[P]$.

(b) $\sim_P$ is defined by:

$$\forall a, b \in A, a \sim_P b \text{ if } (a \leq b \text{ and } b \leq a).$$

It is an equivalence on A .

(c) $\overline{A} = A / \sim_P$ is given an order by:

$$\forall a, b \in A, \overline{a} \leq \overline{b} \text{ if } a \leq b.$$

The poset $(\overline{A}, \leq)$ is denoted by $\overline{P}$.

(d) The cardinality of $\overline{P}$ is denoted by $cl(P)$.

2. Let $n \in \mathbb{N}$.

(a) The set of quasi-posets which underlying set is $[n] = \{1, \dots, n\}$ is denoted by $\mathbf{QP}(n)$.

(b) The set of posets which underlying set is $[n]$ is denoted by $\mathbf{P}(n)$.

(c) The set of isoclasses of quasi-posets of cardinality n is denoted by $\mathbf{qp}(n)$.

(d) The set of isoclasses of quasi-posets of cardinality n is denoted by $\mathbf{p}(n)$.

We put:

$$\begin{aligned}
\mathbf{QP} &= \bigsqcup_{n \geq 0} \mathbf{QP}(n), & \mathbf{P} &= \bigsqcup_{n \geq 0} \mathbf{P}(n), & \mathbf{qp} &= \bigsqcup_{n \geq 0} \mathbf{qp}(n), & \mathbf{p} &= \bigsqcup_{n \geq 0} \mathbf{p}(n), \\
\mathcal{H}_{\mathbf{QP}} &= Vect(\mathbf{QP}), & \mathcal{H}_{\mathbf{P}} &= Vect(\mathbf{P}), & \mathcal{H}_{\mathbf{qp}} &= Vect(\mathbf{qp}) & \mathcal{H}_{\mathbf{p}} &= Vect(\mathbf{p}).
\end{aligned}$$

As posets are quasi-posets, there are canonical injections from $\mathcal{H}_{\mathbf{P}}$ into $\mathcal{H}_{\mathbf{QP}}$ and from $\mathcal{H}_{\mathbf{p}}$ into $\mathcal{H}_{\mathbf{qp}}$. Moreover, the map $P \longrightarrow \overline{P}$ induces surjective maps from $\mathcal{H}_{\mathbf{QP}}$ to $\mathcal{H}_{\mathbf{P}}$ and from

$\mathcal{H}_{\mathbf{qp}}$ to $\mathcal{H}_{\mathbf{p}}$, both denoted by ξ . The map $P \longrightarrow [P]$ induces maps $\llbracket \cdot \rrbracket : \mathcal{H}_{\mathbf{QP}} \longrightarrow \mathcal{H}_{\mathbf{qp}}$ and $\llbracket \cdot \rrbracket : \mathcal{H}_{\mathbf{P}} \longrightarrow \mathcal{H}_{\mathbf{p}}$. The following diagram commutes:

$$\begin{array}{ccc}
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{qp}} \\
\uparrow & \searrow \xi & \uparrow \searrow \xi \\
& \mathcal{H}_{\mathbf{P}} & \xrightarrow{\sqcup} \mathcal{H}_{\mathbf{p}} \\
& \nearrow Id & \nearrow Id \\
\mathcal{H}_{\mathbf{P}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{p}}
\end{array} \quad (1)$$

We shall represent any element P of $\mathbf{QP}$ by the Hasse graph of $\overline{P}$, indicating on the vertices the elements of the corresponding equivalence class. For example, the elements of $\mathbf{QP}(n)$, $n \leq 3$, are:

$$1; \quad \cdot_1; \quad \cdot_1 \cdot_2, \mathbf{1}_1^2, \mathbf{1}_2^1, \cdot_{1,2}; \quad \cdot_1 \cdot_2 \cdot_3, \cdot_1 \mathbf{1}_2^3, \cdot_1 \mathbf{1}_3^2, \cdot_2 \mathbf{1}_1^3, \cdot_2 \mathbf{1}_3^1, \cdot_3 \mathbf{1}_1^2, \cdot_3 \mathbf{1}_2^1, \cdot_1 \cdot_2 \cdot_3, \cdot_2 \cdot_1 \cdot_3, \cdot_3 \cdot_1 \cdot_2,$$

$${}^2\mathbb{V}_1^3, {}^1\mathbb{V}_2^3, {}^1\mathbb{V}_3^2, {}_2\mathbb{A}_3^1, {}_1\mathbb{A}_3^2, {}_1\mathbb{A}_2^3, \mathbb{I}_1^3, \mathbb{I}_1^2, \mathbb{I}_2^3, \mathbb{I}_2^2, \mathbb{I}_3^2, \mathbb{I}_3^1, \mathbb{I}_1^{2,3}, \mathbb{I}_2^{2,3}, \mathbb{I}_3^{1,2}, \mathbb{I}_{2,3}^1, \mathbb{I}_{1,3}^2, \mathbb{I}_{1,2}^3, \bullet_{1,2,3}.$$

We shall represent any element $P \in \mathbf{qp}$ by the Hasse graph of $\overline{P}$, indicating on the vertices the cardinality of the corresponding equivalence class, if this cardinality is not equal to 1. For example, the elements of $\mathbf{qp}(n)$, $n \leq 3$, are:

$$1; \quad \bullet; \quad \bullet\bullet, \mathbf{1}, \bullet_2; \quad \bullet\bullet\bullet, \bullet\mathbf{1}, \bullet\bullet_2; \quad \vee, \wedge, \mathbf{1}, \mathbf{1}^2, \mathbf{1}_{2, \bullet 3}.$$

2.2 First coproduct

By Alexandroff's theorem [1, 18], finite quasi-posets are in bijection with finite topological spaces. Let us recall the definition of the topology attached to a quasi-poset.

Definition 12 1. Let $P = (A, \leq)$ be a quasi-poset. An open set of P is a subset O of A such that:

$$\forall i, j \in A, (i \in O \text{ and } i \leq j) \implies (j \in O).$$

The set of open sets of P (the topology associated to P) is denoted by $\text{top}(P)$.

2. Let $P = (A, \leq)$ be a quasi-poset and $B \subseteq A$. We denote by $P|_B$ the quasi-poset $(B, \leq|_B)$.

3. Let $P = (A, \leq_P)$ be a quasi-poset. We assume that A is also given a total order $\leq$: for example, A is a subset of $\mathbb{N}$. If the cardinality of A is n , there exists a unique increasing bijection f from $[n]$, with its usual order, to $(A, \leq)$. We denote by $\text{Std}(P)$ the quasi-poset, element of $\mathbf{QP}(n)$, defined by:

$$\forall i, j \in [n], i \leq_{Std(P)} j \iff f(i) \leq_P f(j).$$

Proposition 13 1. We define a product m on $\mathcal{H}_{\mathbf{QP}}$ in the following way: if $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, then $PQ = m(P, Q) \in \mathbf{QP}(k + l)$ and

$$\forall i, j \in [k+l], i \leq_{PQ} j \iff (1 \leq i, j \leq k \text{ and } i \leq_P j) \\ \text{or } (k+1 \leq i, j \leq k+l \text{ and } i-k \leq_Q j-k).$$

2. We define a second product $\downarrow$ on $\mathcal{H}_{\mathbf{QP}}$ in the following way: if $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, then $PQ = m(P, Q) \in \mathbf{QP}(k+l)$ and

$$\begin{aligned} \forall i, j \in [k+l], i \leq_{PQ} j \iff & (1 \leq i, j \leq k \text{ and } i \leq_P j) \\ & \text{or } (k+1 \leq i, j \leq k+l \text{ and } i-k \leq_Q j-k) \\ & \text{or } (1 \leq i \leq k < j \leq k+l). \end{aligned}$$

3. We define a coproduct Δ on $\mathcal{H}_{\mathbf{QP}}$ in the following way:

$$\forall P \in \mathbf{QP}(n), \Delta(P) = \sum_{O \in \text{top}(P)} \text{Std}(P|_{[n] \setminus O}) \otimes \text{Std}(P|_O).$$

Then $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ is a non-commutative, non-cocommutative Hopf algebra, and $(\mathcal{H}_{\mathbf{QP}}, \downarrow, \Delta)$ is an infinitesimal bialgebra.

Proof. See [10, 11]. □

Examples. If $\{a, b\} = \{1, 2\}$ and $\{i, j, k\} = \{1, 2, 3\}$:

$$\begin{aligned} \Delta(\bullet_1) &= \bullet_1 \otimes 1 + 1 \otimes \bullet_1, \\ \Delta(\mathbf{1}_a^b) &= \mathbf{1}_a^b \otimes 1 + 1 \otimes \mathbf{1}_a^b + \bullet_a \otimes \bullet_b, \\ \Delta({}^j\mathbf{V}_i^k) &= {}^j\mathbf{V}_i^k \otimes 1 + 1 \otimes {}^j\mathbf{V}_i^k + \mathbf{1}_i^j \otimes \bullet_k + \mathbf{1}_i^k \otimes \bullet_j + \bullet_i \otimes \bullet_j \bullet_k, \\ \Delta({}_j\mathbf{A}_k^i) &= {}_j\mathbf{A}_k^i \otimes 1 + 1 \otimes {}_j\mathbf{A}_k^i + \bullet_j \otimes \mathbf{1}_k^i + \bullet_k \otimes \mathbf{1}_j^i + \bullet_j \bullet_k \otimes \bullet_i, \\ \Delta(\mathbf{1}_i^k) &= \mathbf{1}_i^k \otimes 1 + 1 \otimes \mathbf{1}_i^k + \bullet_i \otimes \mathbf{1}_j^k + \mathbf{1}_i^j \otimes \bullet_k. \end{aligned}$$

Remark. This Hopf algebraic structure is compatible with the morphisms of (1), that is to say:

1. $\mathcal{H}_{\mathbf{P}}$ is a Hopf subalgebra of $\mathcal{H}_{\mathbf{QP}}$.
2. observe that:
 - If (P_1, P_2) and (Q_1, Q_2) are pairs of isomorphic quasi-posets, then $P_1 Q_1$ and $P_2 Q_2$ are isomorphic.
 - If P_1 and P_2 are isomorphic quasi-posets of $\mathbf{QP}(n)$, and if $\phi : [n] \rightarrow [n]$ is an isomorphism from P_1 to P_2 , then the topology associated to P_2 is the image by ϕ of the topology associated to P_1 and for any subset I of P_1 , $\phi|_I$ is an isomorphism from $(P_1)|_I$ to $(P_2)|_{\phi(I)}$.

Consequently, the surjective map $\lfloor \rfloor : \mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{qp}}$ is compatible with the product and the coproduct: $\mathcal{H}_{\mathbf{qp}}$ inherits a Hopf algebra structure. Its product is the disjoint union of quasi-posets. For any quasi-poset $P = (A, \leq_P)$:

$$\Delta(\lfloor P \rfloor) = \sum_{O \in \text{top}(P)} \lfloor P|_{A \setminus O} \rfloor \otimes \lfloor P|_O \rfloor.$$

3. $\mathcal{H}_{\mathbf{P}}$ is a Hopf subalgebra of $\mathcal{H}_{\mathbf{qp}}$.
4. All the morphisms in (1) are Hopf algebra morphisms.

Definition 14 1. We shall say that a finite quasi-poset $P = (A, \leq_P)$ is connected if its associated topology is connected.

2. For any finite quasi-poset P , we denote by $cc(P)$ the number of connected components of its associated topology.

It is well-known that P is connected if, and only if, the Hasse graph of $\overline{P}$ is connected. Any quasi-poset P can be decomposed as the disjoint union of its connected components; in an algebraic setting, $\mathcal{H}_{\mathbf{QP}}$ is generated as a polynomial algebra by the connected quasi-posets. This is not true in $\mathcal{H}_{\mathbf{QP}}$: for example, $\mathbf{!}_1^3 \cdot_2$ is both not connected and indecomposable in $\mathcal{H}_{\mathbf{QP}}$.

2.3 Second coproduct

Definition 15 Let $P = (A, \leq_P)$ be a quasi-poset and let $\sim$ be an equivalence on A .

1. We define a second quasi-order $\leq_{P|\sim}$ on A by the relation:

$$\forall x, y \in A, x \leq_{P|\sim} y \text{ if } (x \leq_P y \text{ and } x \sim y).$$

2. We define a third quasi-order $\leq_{P/\sim}$ on A as the transitive closure of the relation $\mathcal{R}$ defined by:

$$\forall x, y \in A, x \mathcal{R} y \text{ if } (x \leq_P y \text{ or } x \sim y).$$

3. We shall say that $\sim$ is P -compatible and we shall denote $\sim \triangleleft P$ if the two following conditions are satisfied:

- The restriction of P to any equivalence class of $\sim$ is connected.
- The equivalences $\sim_{P/\sim}$ and $\sim$ are equal. In other words:

$$\forall x, y \in A, (x \leq_{P/\sim} y \text{ and } y \leq_{P/\sim} x) \implies x \sim y;$$

note that the converse assertion trivially holds.

Remarks.

1. $P|\sim$ is the disjoint union of the restriction of $\leq_P$ to the equivalence classes of $\sim$.
2. Let $x, y \in P$. Then $x \leq_{P/\sim} y$ if there exist $x_1, x'_1, \dots, x_k, x'_k \in A$ such that:

$$x \leq_P x_1 \sim x'_1 \leq_P \dots \leq_P x_k \sim x'_k \leq_P y.$$

3. If $\sim \triangleleft P$, then:

- (a) The equivalence classes of $\sim_{P/\sim}$ are the equivalence classes of $\sim$ and are included in a connected component of P . This implies that the connected components of $P/\sim$ are the connected components of P . Consequently:

$$cl(P/\sim) = cl(\sim), \quad cc(P/\sim) = cc(P), \quad (2)$$

where $cl(\sim)$ is the number of equivalence classes of $\sim$.

- (b) If $x \sim_P y$ and $x \sim y$, then $x \sim_{P|\sim} y$: the equivalence classes of $\sim_{P|\sim}$ are the equivalence classes of $\sim_P$; the connected components of $P|\sim$ are the equivalence classes of $\sim$. Consequently:

$$cl(P|\sim) = cl(P), \quad cc(P|\sim) = cl(\sim). \quad (3)$$

Definition 16 Let $P \in \mathbf{QP}$. We shall say that P is discrete if $[\overline{P}] = \cdot^{cl(P)}$.

In other words, P is discrete if, and only if, $\sim_P = \leq_P$.

Definition 17 We define a second coproduct δ on $\mathcal{H}_{\mathbf{QP}}$ in the following way: for all $P \in \mathbf{QP}$,

$$\delta(P) = \sum_{\sim \triangleleft P} (P / \sim) \otimes (P | \sim).$$

Then $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ is a bialgebra. Its counit ε' is given by:

$$\forall P \in \mathbf{QP}, \varepsilon'(P) = \begin{cases} 1 & \text{if } P \text{ is discrete,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Firstly, let us prove the compatibility of δ and m . Let $P = (A, \leq_P)$ and $Q = (B, \leq_Q)$ be two elements of $\mathbf{QP}$. Let $\sim$ be an equivalence relation on P . We denote by $\sim'$ and $\sim''$ the restriction of $\sim$ to P and Q . Then:

- If $\sim \triangleleft PQ$, then as the equivalence classes of $\sim$ are connected, they are included in A or in B . Consequently, if $x \in A$ and $y \in B$, x and y are not equivalent for $\sim$. Moreover, $\sim' \triangleleft P$ and $\sim'' \triangleleft Q$, and:

$$PQ | \sim = (P | \sim')(Q | \sim''), \quad PQ / \sim = (P / \sim')(Q / \sim'').$$

- Conversely, if $\sim' \triangleleft P, \sim'' \triangleleft Q$ and for all $x \in A, y \in B$, x and y are not $\sim$ -equivalent, then $\sim \triangleleft PQ$.

Hence:

$$\begin{aligned} \delta(PQ) &= \sum_{\sim \triangleleft PQ} (PQ / \sim) \otimes (PQ | \sim) \\ &= \sum_{\sim' \triangleleft P, \sim'' \triangleleft Q} (P / \sim')(Q / \sim'') \otimes (P | \sim')(Q | \sim'') \\ &= \delta(P)\delta(Q). \end{aligned}$$

Let us now prove the coassociativity of δ . Let $P \in \mathbf{QP}$.

First step. We put:

$$A = \{(r, r') \mid r \triangleleft P, r' \triangleleft P/r\}, \quad B = \{(s, s') \mid s \triangleleft P, s' \triangleleft P|s\}.$$

We consider the maps:

$$F : \begin{cases} A & \longrightarrow B \\ (r, r') & \longrightarrow (r', r), \end{cases} \quad G : \begin{cases} B & \longrightarrow A \\ (s, s') & \longrightarrow (s', s). \end{cases}$$

F is well-defined: we put $(s, s') = (r', r)$. The equivalence classes of s are the equivalence classes of r' , so are P -connected. If $x \sim_{P/s} y$, there exist $x_1, x'_1, \dots, x_k, s'_k$ and $y_1, y'_1, \dots, y_l, y_l$ such that:

$$x \leq_P x_1 r' x'_1 \leq_P \dots \leq_P x_k r' x'_k \leq_P y, \quad y \leq_P y_1 r' y'_1 \leq_P \dots \leq_P y_l r' y'_l \leq_P x.$$

Hence:

$$x \leq_{P/r} x_1 r' x'_1 \leq_{P/r} \dots \leq_{P/r} x_k r' x'_k \leq_{P/r} y, \quad y \leq_{P/r} y_1 r' y'_1 \leq_{P/r} \dots \leq_{P/r} y_l r' y'_l \leq_{P/r} x.$$

So $x \sim_{P/r} y$. As $r' \triangleleft P/r, x \sim_P y: s \triangleleft P$.

Let us assume that $xs'y$. Then xry , so, as $r \triangleleft y$, there exists a path from x to y in the Hasse graph of P , made of vertices all r -equivalent to x and y . If x' and y' are two elements of this path, Then $x'r'y'$, so $x' \leq_{G/r} y'$ and finally $x' \leq_{(P/r)/r'} y'$. As $r' \triangleleft P/r, x'r'y'$, so xsy . So the elements of this path are all $P|s$ -equivalent: the equivalence classes of s' are $P|s$ -connected.

Let us assume that $x \sim_{(P|s)/s'} y$. There exist $x_1, x'_1, \dots, x_k, x'_k$ and $y_1, y'_1, \dots, y_l, y'_l$ such that:

$$x \leq_{P|r'} x_1 r x'_1 \leq_{P|r'} \dots \leq_{P|r'} x_k r x'_k \leq_{P|r'} y, \quad y \leq_{P|r'} y_1 r y'_1 \leq_{P|r'} \dots \leq_{P|r'} y_l r y'_l \leq_{P|r'} x.$$

Then:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y, \quad y \leq_P y_1 r y'_1 \leq_P \dots \leq_P y_l r y'_l \leq_P x,$$

So $x \leq_{P/r} y$ and $y \leq_{P/r} x$. As $r \triangleleft P$, xry , so $xs'y$: we obtain that $s' \triangleleft P | s$.

G is well-defined: let $(s, s') \in B$ and let us put $G(s, s') = (r, r')$. The equivalence classes of r are $P|s$ -connected, so are P -connected. Let us assume that $x \sim_{P/r} y$. There exist $x_1, x'_1, \dots, x_k, x'_k$ and $y_1, y'_1, \dots, y_l, y'_l$ such that:

$$x \leq_P x_1 s' x'_1 \leq_P \dots \leq_P x_k s' x'_k \leq_P y, \quad y \leq_P y_1 s' y'_1 \leq_P \dots \leq_P y_l s' y'_l \leq_P x.$$

As the equivalence classes of s' are $P|s$ -connected, all this elements are in the same connected component of $P|s$, so are s -equivalent:

$$x \leq_{P|s} x_1 s' x'_1 \leq_{P|s} \dots \leq_{P|s} x_k s' x'_k \leq_{P|s} y, \quad y \leq_{P|s} y_1 s' y'_1 \leq_{P|s} \dots \leq_{P|s} y_l s' y'_l \leq_{P|s} x.$$

Hence, $x \sim_{(P|s)/s'} y$, so as $s' \triangleleft P | s$, $xs'y$, so xry : $r \triangleleft P$.

The equivalence classes of r' are the equivalence classes of s , so are P -connected and therefore P/r -connected. Let us assume that $x \sim_{(P/r)/r'} y$. Note that if $x's'y'$, then x' and y' are in the same connected component of $P|s$, so $x'sy$. By the definition of $\leq_{P/s'}$ as a transitive closure, using this observation, we obtain:

$$x \leq_P x_1 s x'_1 \leq_P \dots \leq_P x_k s x'_k \leq_P y, \quad y \leq_P y_1 s y'_1 \leq_P \dots \leq_P y_l s y'_l \leq_P x.$$

So $x \sim_{P/s} y$. As $s \triangleleft P$, xsy , so $xr'y$: $r' \triangleleft P/r$.

Clearly, F and G are inverse bijections.

Second step. Let $(r, r') \in A$ and let $F(r, r') = (s, s')$. Note that if xry , then $x/ \sim_{P/r} y$, so $x/ \sim_{(P/r)/r'} y$, so $xr'y$ as $r' \triangleleft P/r$. Then:

$$\begin{aligned} \leq_{(P/r)/r'} &= \text{transitive closure of } ((xr'y) \text{ or } (x \leq_{P/r} y)) \\ &= \text{transitive closure of } ((xr'y) \text{ or } (x \leq_P y) \text{ or } (x \leq_r y)) \\ &= \text{transitive closure of } ((xr'y) \text{ or } (x \leq_P y)) \\ &= \text{transitive closure of } ((xsy) \text{ or } (x \leq_P y)) \\ &= \leq_{P/s}. \end{aligned}$$

So $P/s = (P/r)/r'$.

$$\begin{aligned} \leq_{(P|s)/s'} &= \text{transitive closure of } ((xs'y) \text{ or } (x \leq_{P|s} y)) \\ &= \text{transitive closure of } ((xry) \text{ or } (x \leq_{P|r'} y)) \\ &= \text{transitive closure of } ((xry) \text{ or } ((x \leq_P y) \text{ and } (xr'y))) \\ &= \text{transitive closure of } (((xry) \text{ or } (x \leq_P y)) \text{ and } ((sry) \text{ or } (sr'y))) \\ &= \text{transitive closure of } ((x \leq_{P/r} y) \text{ and } (sr'y)) \\ &= \leq_{(P/r)/r'}. \end{aligned}$$

So $(P|s)/s' = (P/r)|r'$. For all x, y :

$$\begin{aligned}
x \leq_{(P|s)/s'} y &\iff (x \leq_{P|s} y) \text{ and } (xs'y) \\
&\iff (x \leq_P y) \text{ and } xsy \text{ and } (xs'y) \\
&\iff (x \leq_P y) \text{ and } xr'y \text{ and } (xry) \\
&\iff (x \leq_P y) \text{ and } (xry) \\
&\iff x \leq_{P|r} y.
\end{aligned}$$

So $(P|s)|s' = P|r$. Finally:

$$\begin{aligned}
(\delta \otimes Id) \circ \delta(P) &= \sum_{(r,r') \in A} (P/r)/r' \otimes (P/r)|r' \otimes P|r \\
&= \sum_{(s,s') \in B} P/s \otimes (P|s)/s' \otimes (P|s)|s' \\
&= (Id \otimes \delta) \circ \delta(P).
\end{aligned}$$

So $\mathcal{H}_{\mathbf{QP}}$ is a bialgebra.

Let P be a quasi-poset. If P is discrete, then $\delta(P) = P \otimes P$, so $(\varepsilon' \otimes Id) \circ \delta(P) = (Id \otimes \varepsilon') \circ \Delta(P) = P$. If P is not discrete, there are three types of relations $\sim \triangleleft P$:

1. The equivalence classes of $\sim$ are the connected components of P : in this case, $P| \sim = P$ and $P/ \sim = P_1$ is discrete.
2. $\sim = \sim_P$: in this case, $P/ \sim = P$ and $P| \sim = P_2$ is discrete.
3. $\sim$ is not one of two preceding relations: in this case, nor $P/ \sim$, nor $P| \sim$ is discrete.

So:

$$\delta(P) - P_1 \otimes P - P \otimes P_2 \in Ker(\varepsilon') \otimes Ker(\varepsilon'),$$

which implies that $(\varepsilon' \otimes Id) \circ \delta(P) = (Id \otimes \varepsilon') \circ \Delta(P) = P$. □

Examples. If $\{a, b\} = \{1, 2\}$ and $\{i, j, k\} = \{1, 2, 3\}$:

$$\begin{aligned}
\delta(\cdot_1) &= \cdot_1 \otimes \cdot_1, \\
\delta(\mathbf{1}_a^b) &= \mathbf{1}_a^b \otimes \cdot_a \cdot_b + \cdot_{a,b} \otimes \mathbf{1}_a^b, \\
\delta({}^j\mathbf{V}_i^k) &= {}^j\mathbf{V}_i^k \otimes \cdot_i \cdot_j \cdot_k + \mathbf{1}_{i,j}^k \otimes \mathbf{1}_i^j \cdot_k + \mathbf{1}_{i,k}^j \otimes \mathbf{1}_i^k \cdot_j + \cdot_{i,j,k} \otimes {}^j\mathbf{V}_i^k, \\
\delta({}_j\mathbf{A}_i^k) &= {}_j\mathbf{A}_i^k \otimes \cdot_i \cdot_j \cdot_k + \mathbf{1}_k^{i,j} \otimes \mathbf{1}_j^i \cdot_k + \mathbf{1}_j^{i,k} \otimes \mathbf{1}_k^i \cdot_j + \cdot_{i,j,k} \otimes {}_j\mathbf{A}_i^k, \\
\delta(\mathbf{1}_i^k) &= \mathbf{1}_i^k \otimes \cdot_i \cdot_j \cdot_k + \mathbf{1}_{i,j}^k \otimes \mathbf{1}_i^j \cdot_k + \mathbf{1}_i^{j,k} \otimes \cdot_i \mathbf{1}_j^k + \cdot_{i,j,k} \otimes \mathbf{1}_i^k.
\end{aligned}$$

Remarks.

1. δ is the internal coproduct of [8].
2. $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ is not a Hopf algebra: for all $n \geq 1$, $\delta(\cdot_n) = \cdot_n \otimes \cdot_n$, and $\cdot_n$ has no inverse in $\mathcal{H}_{\mathbf{QP}}$.
3. This coproduct is also compatible with the map $\lfloor \rfloor$, so we obtain a bialgebra structure on $\mathcal{H}_{\mathbf{qp}}$ with the coproduct defined by:

$$\delta(\lfloor P \rfloor) = \sum_{\sim \triangleleft P} [P/ \sim] \otimes [P| \sim].$$

4. $\mathcal{H}_{\mathbf{P}}$ and $\mathcal{H}_{\mathbf{p}}$ are not stable under δ , as if P is a poset and $\sim \triangleleft P$, $P/\sim$ is not necessarily a poset (although $P|\sim$ is). However, there is a way to define a coproduct $\bar{\delta} = (\xi \otimes Id) \circ \delta$ on $\mathcal{H}_{\mathbf{p}}$:

$$\forall P \in \mathbf{P}(n), \bar{\delta}([P]) = \sum_{\sim \triangleleft P} \overline{[P/\sim]} \otimes [P|\sim].$$

$(\mathcal{H}_{\mathbf{p}}, m, \bar{\delta})$ is a quotient of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ through the map ξ .

2.4 Cointeractions

Theorem 18 *We consider the map:*

$$\rho = (Id \otimes [\sqcup]) \circ \delta : \begin{cases} \mathcal{H}_{\mathbf{QP}} & \longrightarrow \mathcal{H}_{\mathbf{QP}} \otimes \mathcal{H}_{\mathbf{qp}} \\ P \in \mathbf{QP} & \longrightarrow \sum_{\sim \triangleleft P} (P/\sim) \otimes [P|\sim]. \end{cases}$$

The bialgebras $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \bar{\delta})$ are in cointeraction via ρ .

Proof. By composition, ρ is an algebra morphism. Let us take $P \in \mathbf{QP}(n)$. We put:

$$A = \{(r, O) \mid r \triangleleft P, O \in \text{top}(P/r)\}, \quad B = \{(O, s, s') \mid O \in \text{top}(P), s \triangleleft P_{[n] \setminus O}, s' \triangleleft P_{|O}\}.$$

First step. We define a map $F : A \longrightarrow B$, sending (r, O) to (O, s, s') , by:

- xsy if xry and x, y are in the same connected component of $P_{|[n] \setminus O}$.
- $xs'y$ if xry and x, y are in the same connected component of $P_{|O}$.

Let us prove that F is well-defined. Let us take $x, y \in [n]$, with $x \in O$ and $x \leq_P y$. Then $x \leq_{P/r} y$. as O is an open set of P/r , $y \in O$: O is an open set of P . By definition, the equivalence classes of s' are the intersection of the equivalence classes of r and of the connected components of O . As O is a union of equivalence classes of r , they are $P_{|O}$ -connected. If $x \sim_{P_{|O}/s'} y$, then $x \sim_{P/r} y$ and x and y are in the same connected component of O . As $r \triangleleft r$, xry , so $xs'y$: $s' \triangleleft P_{|O}$. Similarly, $s \triangleleft P_{|[n] \setminus O}$.

Second step. We define a map $G : B \longrightarrow A$, sending (O, s, s') to (O, r, r') , by:

$$xry \text{ if } (x, y \notin O \text{ and } xsy) \text{ or } (x, y \in O \text{ and } xs'y).$$

Let us prove that G is well-defined. Let $x, y \in [N]$, with $x \in O$ and $x \leq_{P/r} y$. There exists $x_1, x'_1, \dots, x_k, x'_k$ such that:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y.$$

As O is an open set of P , $x_1 \in O$; by definition of r , $x'_1 \in O$. Iterating, we obtain that $x_2, x'_2, \dots, x_k, x'_k, y \in O$. So O is open in P/r .

Let us assume that xry . Then $x, y \in O$ or $x, y \notin O$. As $s \triangleleft P_{|[n] \setminus O}$ and $P_{|O}$, there exists a path from x to y in the Hasse graph of P formed by elements s - or s' - equivalent to x and y , so the equivalence classes of r are P -connected.

Let us assume that $x \sim_{P/r} y$. here exists $x_1, x'_1, \dots, x_k, s'_k$ and $y_1, y'_1, \dots, y_l, y_l$ such that:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y, \quad y \leq_P y_1 r y'_1 \leq_P \dots \leq_P y_l r y'_l \leq_P x.$$

If $x, y \in O$, then all these elements are in O , so $x \sim_{P_{|O}/s'} y$, and then $xs'y$, so xry . If $x, y \notin O$, as O is an open set, none of these elements is in O , so $x \sim_{P_{|[n] \setminus O}/s} y$, so xsy and finally xry : $r \triangleleft P$.

Third step. Let $(r, O) \in A$. We put $F(r, O) = (O, s, s')$ and $G(O, s, s') = (\tilde{r}, O)$. If xry , as O is an open set of P/r , both x and y are in O or both are not in O . Hence, xsy or $xs'y$, so $x\tilde{r}y$.

If $x\tilde{r}y$, then xsy or $xs'y$, so xry : $\tilde{r} = r$ and $G \circ F = Id_A$.

Let $(O, s, s') \in B$. We put $G(O, s, s') = (r, O)$ and $F(r, O) = (O, \tilde{s}, \tilde{s}')$. If xsy , then x and y are in the same connected component of $[n] \setminus O$ as $s \triangleleft P_{[n] \setminus O}$ and xry , so $x\tilde{s}y$. If $x\tilde{s}y$, then xry , so xsy : we obtain that $\tilde{s} = s$. Similarly, $\tilde{s}' = s'$, which proves that $F \circ G = Id_B$.

We proved that F and G are inverse bijections. Let $(r, O) \in A$ and $(O, s, s') = F(O, r)$.

$$\begin{aligned} \leq_{(P/r)_{|[n] \setminus O}} &= \text{transitive closure of } (xry \text{ and } x \leq_P y) \text{ restricted to } [n] \setminus O \\ &= \text{transitive closure of } (xry \text{ and } x \leq_{P_{[n] \setminus O}} y) \\ &= \text{transitive closure of } (xsy \text{ and } x \leq_{P_{[n] \setminus O}} y) \\ &= \leq_{P_{[n] \setminus [n]}/s} . \end{aligned}$$

So $(P/r)_{|[n] \setminus O} = P_{[n] \setminus O}/s$. Similarly, $(P/r)_O = P_O/s'$.

Let us now consider $P|_R$. Its connected components are the equivalence classes of r , that is to say the equivalence classes of s and s' ; for any such equivalence class I , $(P|_R)_I = P|_I$. So $P|_R$ is the disjoint union of $(P_{[n] \setminus O})|_s$ and $(P_O)|_{s'}$, and therefore is isomorphic to $Std(P_{[n] \setminus O})|_s Std((P_O)|_{s'})$, but not equal, because of the reindexation induced by the standardization. Hence, $[P|_R] = [(P_{[n] \setminus O})|_s] [(P_O)|_{s'}]$.

Finally:

$$\begin{aligned} (\Delta \otimes Id) \circ \rho(P) &= \sum_{(r, O) \in A} (G/r)_{|[n] \setminus O} \otimes (G/r)_O \otimes [G|_r] \\ &= \sum_{(O, s, s') \in B} (P_{[n] \setminus O})/s \otimes (P_O)/s' \otimes [(P_{[n] \setminus O})|_s] [(P_O)|_{s'}] \\ &= m_{2,4}^3 \circ (\rho \otimes \rho) \circ \Delta(P). \end{aligned}$$

Moreover, $(\varepsilon \otimes Id) \circ \rho(P) = \delta_{P,1} 1 \otimes 1 = \varepsilon(P) 1 \otimes 1$. $\square$

Remark. As noticed in [8], $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ are not in cointeraction through δ .

Taking the quotient through $\sim$:

Corollary 19 *The bialgebras $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ are in cointeraction via δ . Moreover, $\mathcal{H}_{\mathbf{qp}}$ is given a graduation by:*

$$\forall n \geq 0, (\mathcal{H}_{\mathbf{qp}})_n = Vect(P \in \mathbf{qp} \mid cl(P) = n).$$

With this graduation, hypotheses 1–5 of section 1.3 are satisfied.

Proof. Here, V is the space generated by the set of connected quasi-posets; the basis $(g_i)_{i \in I}$ of group-like elements of $(\mathcal{H}_{\mathbf{qp}})_1$ is $(\bullet_n)_{n \geq 1}$. $\square$

We denote by $M_{\mathbf{qp}}$ the monoid of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$. Its product is denoted by $*$. Using proposition 4 on $\mathcal{H}_{\mathbf{QP}}$:

Corollary 20 *Let $\lambda \in M_{\mathbf{qp}}$. The following map is a Hopf algebra endomorphism:*

$$\phi_\lambda : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P \in \mathbf{QP} & \longrightarrow \sum_{\sim \triangleleft P} \lambda_{[P| \sim]} P / \sim . \end{cases}$$

It is bijective if, and only if, for all $n \geq 1$, $\lambda_{\bullet_n} \neq 0$. If this holds, $\phi_\lambda^{-1} = \phi_{\lambda^}^{-1}$.*

Proof. $\phi_\lambda = Id \leftarrow \lambda$, so is an element of $E_{\mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{QP}}}$.

$\implies$. For all $n \geq 0$, $\phi_\lambda(\cdot_n) = \lambda \cdot_n \cdot_n$. As ϕ_λ is injective, $\lambda \cdot_n \neq 0$.

$\impliedby$. By proposition 5, λ is invertible in $M_{\mathbf{QP}}$: let us denote its inverse by μ . Then, by proposition 4:

$$\phi_\lambda \circ \phi_\mu = Id \leftarrow (\lambda * \mu) = Id \leftarrow \varepsilon' = Id.$$

Similarly, $\phi_\mu \circ \phi_\lambda = Id$. □

3 Ehrhart polynomials

Notations. For all $k \geq 0$, we denote by H_k the k -th Hilbert polynomial:

$$H_k(X) = \frac{X(X-1)\dots(X-k+1)}{k!}.$$

3.1 Definition

Definition 21 Let $P \in \mathbf{QP}(n)$ and $k \geq 1$. We put:

$$\begin{aligned} L_P(k) &= \{f : [n] \longrightarrow [k] \mid \forall i, j \in [n], i \leq_P j \implies f(i) \leq f(j)\}, \\ L_P^{str}(k) &= \{f \in L_P(k) \mid \forall i, j \in [n], (i \leq_P j \text{ and } f(i) = f(j)) \implies i \sim_P j\}, \\ W_P(k) &= \{w \in L_P(k) \mid w([n]) = [k]\}, \\ W_P^{str}(k) &= \{w \in L_P^{str}(k) \mid w([n]) = [k]\}. \end{aligned}$$

By convention:

$$L_P(0) = L_P^{str}(0) = W_P(0) = W_P^{str}(0) = \begin{cases} \emptyset & \text{if } P \neq 1, \\ \{1\} & \text{if } P = 1. \end{cases}$$

We also put:

$$L_P = \bigcup_{k \geq 0} L_P(k), \quad L_P^{str} = \bigcup_{k \geq 0} L_P^{str}(k), \quad W_P = \bigsqcup_{k \geq 0} W_P(k), \quad W_P^{str} = \bigsqcup_{k \geq 0} W_P^{str}(k).$$

Note that the elements of W_P and W_P^{str} are packed words (see definition 40).

Proposition 22 Let $P \in \mathbf{QP}$. There exist unique polynomials ehr_P and $ehr_P^{str} \in \mathbb{Q}[X]$, such that for $k \geq 0$:

$$ehr_P(k) = \#L_P(k), \quad ehr_P^{str}(k) = \#L_P^{str}(k).$$

Proof. This is obvious if $P = 1$, with $ehr_1(X) = ehr_1^{str}(X) = 1$. Let us assume that $P \in \mathbf{QP}(n)$, $n \geq 1$. Note that if $i > n$, $W_P(i) = 0$. For all $k \geq 1$:

$$\#L_P(k) = \sum_{i=1}^k \#W_P(i) \binom{k}{i} = \sum_{i=1}^k \#W_P(i) H_i(k) = \sum_{i=1}^n \#W_P(i) H_i(k).$$

So:

$$ehr_P(X) = \sum_{i=1}^n \#W_P(i) H_i(X).$$

Moreover, if $k = 0$:

$$ehr_P(0) = \sum_{i=1}^n \#W_P(i)H_i(0) = \#L_P(0).$$

In the same way:

$$ehr_P^{str}(X) = \sum_{i=1}^n \#W_P^{str}(i)H_i(X).$$

These are indeed elements of $\mathbb{Q}[X]$. □

Remarks.

1. Let $P, Q \in \mathbf{QP}(n)$.

- If they are isomorphic, then $ehr_P(k) = ehr_Q(k)$ for all $k \geq 1$, so $ehr_P = ehr_Q$.
- If $w \in L_P$, for all $x, y \in P$ such that $x \sim_P y$, then $w(x) \leq w(y)$ and $w(y) \leq w(x)$, so $w(x) = w(y)$: w goes through the quotient by $\sim_P$. We obtain in this way a bijection from $L_P(k)$ to $L_{\overline{P}}(k)$ for all k , so $ehr_P = ehr_{\overline{P}}$. Similarly, $ehr_P^{str} = ehr_{\overline{P}}^{str}$.

Hence, we obtain maps, all denoted by ehr and ehr^{str} , such the following diagrams commute:

$$\begin{array}{ccc} \mathcal{H}_{\mathbf{QP}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{qp}} \xrightarrow{\xi} \mathcal{H}_{\mathbf{p}} \\ & \searrow \text{ehr} & \downarrow \text{ehr} \nearrow \text{ehr} \\ & & \mathbb{K}[X] \end{array} \quad \begin{array}{ccc} \mathcal{H}_{\mathbf{QP}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{qp}} \xrightarrow{\xi} \mathcal{H}_{\mathbf{p}} \\ & \searrow \text{ehr}^{str} & \downarrow \text{ehr}^{str} \nearrow \text{ehr}^{str} \\ & & \mathbb{K}[X] \end{array}$$

2. Let $P \in \mathbf{P}(n)$. The classical definition of the Ehrhart polynomial $ehr^{cl}(t)$ is the number of integral points of $tPol(P)$, where $Pol(P)$ is the polytope associated to P . Hence, $ehr^{cl}(X) = ehr(X + 1)$.

Theorem 23 *The morphisms $ehr, ehr^{str} : \mathcal{H}_{\mathbf{QP}}, \mathcal{H}_{\mathbf{qp}}, \mathcal{H}_{\mathbf{p}} \longrightarrow \mathbb{K}[X]$ are Hopf algebra morphisms.*

Proof. It is enough to prove it for $ehr, ehr^{str} : \mathcal{H}_{\mathbf{p}} \longrightarrow \mathbb{K}[X]$.

First step. Let $P \in \mathbf{P}(n)$. Let us prove that for all $k, l \geq 0$:

$$ehr_P(k + l) = \sum_{O \in Top(P)} ehr_{P_{[n] \setminus O}}(k) ehr_{P|_O}(l), \quad ehr_P^{str}(k + l) = \sum_{O \in Top(P)} ehr_{P_{[n] \setminus O}}^{str}(k) ehr_{P|_O}^{str}(l).$$

Let $f \in L_P(k + l)$. We put $O = f^{-1}(\{k + 1, \dots, k + l\})$. If $x \in O$ and $x \leq_P y$, then $f(x) \leq f(y)$, so $y \in O$: O is an open set of P . By restriction, the following maps are elements of respectively $L_{P_{[n] \setminus O}}(k)$ and $L_{P|_O}(l)$:

$$f_1 : \begin{cases} [n] \setminus O & \longrightarrow [k] \\ x & \longrightarrow f(x), \end{cases} \quad f_2 : \begin{cases} O & \longrightarrow [l] \\ x & \longrightarrow f(x) - k. \end{cases}$$

This defines a map:

$$v : \begin{cases} L_P(k + l) & \longrightarrow \bigsqcup_{O \in Top(P)} L_{P_{[n] \setminus O}}(k) \times L_{P|_O}(l) \\ f & \longrightarrow (f_1, f_2). \end{cases}$$

This map is clearly injective; moreover:

$$\nu(L_P^{str}(k+l)) \subseteq \bigsqcup_{O \in Top(P)} L_{P_{[n] \setminus O}}^{str}(k) \times L_{P|_O}^{str}(l).$$

Let us prove that f is surjective. Let $(f_1, f_2) \in L_{P_{[n] \setminus O}}(k) \otimes L_{P|_O}(l)$, with $O \in Top(P)$. We define a map $f : P \longrightarrow [k+l]$ by:

$$f(x) = \begin{cases} f_1(x) & \text{if } x \notin O, \\ f_2(x) + k & \text{if } x \in O. \end{cases}$$

Let $x \leq_P i$. As O is an open set of P , three cases are possible:

- $x, y \notin O$: then $f_1(x) \leq f_1(y)$, so $f(x) \leq f(y)$.
- $x, y \in O$: then $f_2(x) \leq f_2(y)$, so $f(x) \leq f(y)$.
- $x \notin O, y \in O$: then $f(x) \leq k < fyj$.

So $f \in L_P(k+l)$, and $\nu(f) = (f_1, f_2)$: ν is surjective, and finally bijective. Moreover, if $f_1 \in L_{[n] \setminus O}^{str}(k)$ and $f_2 \in L_{P|_O}^{str}(l)$, then $f = \nu^{-1}(f_1, f_2) \in L_P^{str}(k+l)$. Finally:

$$\begin{aligned} f(L_P(k+l)) &= \bigsqcup_{O \in Top(P)} L_{P_{[n] \setminus O}}(k) \times L_{P|_O}(l), \\ f(L_P^{str}(k+l)) &= \bigsqcup_{O \in Top(P)} L_{P_{[n] \setminus O}}^{str}(k) \times L_{P|_O}^{str}(l). \end{aligned}$$

Taking the cardinals, we obtain the announced result.

Second step. Let $P \in \mathbf{P}(m)$, $Q \in \mathbf{P}(n)$, and $f : [m+n] \longrightarrow [k]$. Then $f \in L_{PQ}(k)$ if, and only if, $f|_{[m]} \in L_P(k)$ and $Std(f|_{[m+n] \setminus [m]}) \in L_Q(k)$. So $ehr_{PQ}(k) = ehr_P(k)ehr_Q(k)$, and then $ehr_{PQ}(X) = ehr_P(X)ehr_Q(X)$: ehr is an algebra morphism.

Let P be a finite poset, and $k, l \geq 0$. By the first step:

$$\begin{aligned} (ehr \otimes ehr) \circ \Delta(P)(k, l) &= \sum_{O \in Top(P)} ehr_{P_{[n] \setminus O}}(k) ehr_{P|_O}(l) \\ &= ehr_P(k+l) \\ &= \Delta \circ ehr(P)(k, l). \end{aligned}$$

As this is true for all $k, l \geq 1$, $(ehr \otimes ehr) \circ \Delta(P) = \Delta \circ ehr(P)$. Moreover:

$$\varepsilon \circ ehr(P) = ehr_P(0) = \begin{cases} 1 & \text{if } P = 1, \\ 0 & \text{otherwise,} \end{cases}$$

so $\varepsilon \circ ehr = \varepsilon$.

The proof is similar for ehr^{str} . □

3.2 Recursive computation of ehr and ehr^{str}

Let us recall this classical result:

Lemma 24 *We consider the following maps:*

$$L : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K}[X] \\ H_k(X) & \longrightarrow H_{k+1}(X). \end{cases}$$

The map L is injective, and $L(\mathbb{K}[X]) = \mathbb{K}[X]_+$. Moreover, for all $P \in \mathbb{K}[X]$, for all $n \geq 0$:

$$L(P)(n+1) = P(0) + \dots + P(n).$$

Proof. Let us consider $P = H_k(X)$. For all $n \geq 0$:

$$\begin{aligned}
H_k(0) + \dots + H_k(n) &= \binom{0}{k} + \dots + \binom{n}{k} \\
&= \binom{k}{k} + \dots + \binom{n}{k} \\
&= \binom{n+1}{k+1} \\
&= H_{k+1}(n+1) \\
&= L(H_k)(n+1).
\end{aligned}$$

By linearity, for any $P \in \mathbb{K}[X]$, $L(P)(n+1) = P(0) + \dots + P(n)$ for all $n \geq 1$. □

Proposition 25 *Let $P \in \mathbf{P}(n)$.*

$$\begin{aligned}
ehr_P(X) &= L \left(\sum_{\emptyset \neq O \in Top(P)} ehr_{P|_{[n] \setminus O}}(X) \right), \\
ehr_P^{str}(X) &= L \left(\sum_{\emptyset \neq O \in Top(P), \text{ discrete}} ehr_{P|_{[n] \setminus O}}^{str}(X) \right).
\end{aligned}$$

Proof. Let $n \geq 1$. As $L_Q(1)$ is reduced to a singleton for all finite poset Q :

$$\begin{aligned}
ehr_P(n+1) &= \sum_{O \in Top(P)} ehr_{P|_{[n] \setminus O}}(n) ehr_{P|_O}(1) \\
&= \sum_{\emptyset \neq O \in Top(P)} ehr_{P|_{[n] \setminus O}}(n) + ehr_P(n).
\end{aligned}$$

We put:

$$Q(X) = \sum_{\emptyset \neq O \in Top(P)} ehr_{P|_{[n] \setminus O}}(X).$$

In particular:

$$Q(0) = \sum_{\emptyset \neq O \in Top(P)} ehr_{P|_{[n] \setminus O}}(0) = ehr_{\emptyset}(0) + 0 = 1 = ehr_P(1).$$

Then:

$$\begin{aligned}
ehr_P(n+1) &= Q(n) + ehr_P(n) \\
&= Q(n) + Q(n-1) + ehr_P(n-1) \\
&\vdots \\
&= Q(n) + Q(n-1) + \dots + Q(1) + ehr_P(1) \\
&= Q(n) + \dots + Q(1) + Q(0) \\
&= L(Q)(n+1).
\end{aligned}$$

So $ehr_P = L(Q)$.

For ehr_P^{str} , observe that $ehr_Q^{str}(1) = 1$ if Q is discrete, and 0 otherwise, which implies:

$$ehr_P^{str}(n+1) = \sum_{\emptyset \neq O \in Top(P), \text{ discrete}} ehr_{P_{[n] \setminus O}}^{str}(n) + ehr_P^{str}(n).$$

The end of the proof is similar. $\square$

Examples.

$$\begin{aligned} ehr_{\bullet}(X) &= H_1(X) = X, \\ ehr_{\mathbf{1}}(X) &= H_1(X) + H_2(X) = \frac{X(X+1)}{2}, \\ ehr_{\mathbf{V}}(X) &= ehr_{\mathbf{\wedge}}(X) = H_1(X) + 3H_2(X) + 2H_3(X) = \frac{X(X+1)(2X+1)}{6}, \\ ehr_{\mathbf{\dagger}}(X) &= H_1(X) + 2H_2(X) + H_3(X) = \frac{X(X+1)(X+2)}{6}; \\ \\ ehr_{\bullet}^{str}(X) &= H_1(X) = X, \\ ehr_{\mathbf{1}}^{str}(X) &= H_2(X) = \frac{X(X-1)}{2}, \\ ehr_{\mathbf{V}}^{str}(X) &= ehr_{\mathbf{\wedge}}^{str}(X) = H_2(X) + 2H_3(X) = \frac{X(X-1)(2X-1)}{6}, \\ ehr_{\mathbf{\dagger}}^{str}(X) &= H_3(X) = \frac{X(X-1)(X-2)}{6}. \end{aligned}$$

3.3 Characterization of quasi-posets by packed words

Lemma 26 *Let $P \in \mathbf{QP}(n)$ and let $I_1, \dots, I_k$ be distinct minimal classes of the poset $\overline{P}$; let $w' \in W_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}}^{str}$. The following map belongs to W_P^{str} :*

$$w : \begin{cases} [n] & \longrightarrow \mathbb{N}^* \\ x \in I_p, 1 \leq p \leq k & \longrightarrow p \\ x \notin I_1 \sqcup \dots \sqcup I_k & \longrightarrow w'(x). \end{cases}$$

Proof. Let us assume that $i \leq_P j$.

- If $i \in I_p$, as I_p is a minimal class of $\overline{P}$, $j \in I_p$ or $j \notin I_1 \sqcup \dots \sqcup I_k$. In the first case, $w(i) = w(j)$; in the second case, $w(i) \leq k < w(j)$. If moreover $w(i) = w(j)$, then necessarily $j \in I_p$, so $i \sim_P j$.
- If $i \notin I_1 \sqcup \dots \sqcup I_k$, as $i \leq_P j$, $j \notin I_1 \sqcup \dots \sqcup I_k$, so $i \leq_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}} j$ and $w'(i) \leq w'(j)$, so $w'(i) \leq w'(j)$. If moreover $w(i) = w(j)$, then $w'(i) = w'(j)$, so $i \sim_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}} j$ and finally $i \sim_P j$.

As a conclusion, $w \in W_P^{str}$. $\square$

Remark. This lemma implies that W_P^{str} is non-empty for any non-empty quasi-poset P .

Proposition 27 *Let $P = ([n], \leq_P)$ be a quasi-poset and let $i, j \in [n]$. The following assertions are equivalent:*

1. $i \leq_P j$.

2. For all $w \in L_P$, $w(i) \leq w(j)$.
3. For all $w \in L_P^{str}$, $w(i) \leq w(j)$.
4. For all $w \in W_P$, $w(i) \leq w(j)$.
5. For all $w \in W_P^{str}$, $w(i) \leq w(j)$.

Proof. Obviously:

$$\begin{array}{ccccc} 1. & \implies & 2. & \implies & 3. \\ & & \Downarrow & & \Downarrow \\ & & 4. & \implies & 5. \end{array}$$

It is enough to prove that $5. \implies 1.$ We proceed by induction on n . If $n = 1$, there is nothing to prove. Let us assume the result at all ranks $< n$. Let $i, j \in [n]$, such that we do not have $i \leq_P j$. Let us prove that there exists $w \in W_P^{str}$, such that $w(i) > w(j)$. There exists a minimal element $k \in [n]$, such that $k \leq_P j$; let I be the class of k in $\overline{P}$. By hypothesis on i , i and k are not equivalent for $\sim_P$, so $i \notin I$. If $j \in I$, let us choose an element $w' \in W_{P_{[n] \setminus I}}^{str}$; if $j \notin I$, then by the induction hypothesis, there exists $w' \in W_{P_{[n] \setminus I}}^{str}$, such that $w'(i) > w'(j)$. By lemma 26, the following map is an element of W_P^{str} :

$$w : \begin{cases} [n] & \longrightarrow \mathbb{N} \\ x \in I & \longrightarrow 1 \\ x \notin I & \longrightarrow w'(x) + 1 \end{cases}$$

If $j \in I$, then $w(j) = 1 < w(i)$; if $j \notin I$, $w(i) = w'(i) + 1 > w'(j) + 1 = w(j)$. In both cases, $w(i) > w(j)$. $\square$

3.4 Link with linear extensions

Let $P \in \mathbf{QP}(n)$. Linear extensions, as defined in [10], are elements of W_P^{str} : they are the elements $f \in W_P^{str}$ such that

$$\forall i, j \in [n], f(i) = f(j) \iff i \sim_P j.$$

It may happens that not all elements of W_P^{str} are linear extensions. For example, if $P = {}^2\mathbf{V}_1^3$, $W_P^{str}(3) = \{(123), (132), (122)\}$, and (122) is not a linear extension of P . The set of linear extensions of P will be denoted by E_P .

Definition 28 Let w and w' be two packed words of the same length n . We shall say that $w \leq w'$ if:

$$\forall i, j \in [n], w(i) < w(j) \implies w'(i) < w'(j).$$

Proposition 29 Let $P \in \mathbf{QP}(n)$. Then:

$$W_P = \bigcup_{w \in E_P} \{w' \mid w' \leq w\}.$$

This union may be not disjoint. Moreover, the maximal elements of W_P for the order of definition 28 are the elements of E_P .

Proof. $\subseteq$. Let $w \in W_P$. For all $1 \leq p \leq \max(w)$, we put $I_p = w^{-1}(p)$. Let f_p be a linear extension of $P_{|I_p}$. Let us consider:

$$f : \begin{cases} [n] & \longrightarrow \mathbb{N} \\ i & \longrightarrow \max(f_1) + \dots + \max(f_{p-1}) + f_p(i) \text{ if } i \in I_p. \end{cases}$$

By construction, if $w(i) < w(j)$, then $f(i) < f(j)$: $w \leq f$. Let us prove that $f \in E_P$.

If $i \leq_P j$, then as $w \in W_P$, $w(i) \leq w(j)$. If $w(i) = w(j) = p$, then $i \leq_{P|_{I_p}} j$, so $f_p(i) \leq f_p(j)$, and $f(i) \leq f(j)$. If $w(i) < w(j)$, then $f(i) < f(j)$.

If $f(i) = f(j)$, then $w(i) = w(j) = p$, and $f_p(i) = f_p(j)$. As $f_p \in E_{P|_{I_p}}$, $i \sim_{P|_{I_p}} j$, so $i \sim_P j$.

$\supseteq$. Let $w \in E_P$ and $w' \leq w$. If $i \leq_P j$, then $w(i) \leq w(j)$ as w is a linear extension of P . As $w' \leq w$, $w'(i) \leq w'(j)$, so $w' \in W_P$.

Let w be a maximal element of W_P . There exists a linear extension w' of P , such that $w \leq w'$. As w is maximal, $w = w'$ is a linear extension of P . Conversely, if w is a linear extension of P and $w \leq w'$ in W_P , then as w is a linear extension of P , $\max(w) = cl(P)$. Moreover, as $w \leq w'$, $\max(w) \leq \max(w')$. As $w' \in W_P$, $\max(w') \leq cl(P)$, which implies that $\max(w) = \max(w') = cl(P)$, and finally $w = w'$: w is a maximal element of W_P . $\square$

Example. For $P = {}^2\mathbf{V}_1^3$:

$$\begin{aligned} E_P &= \{(123), (132)\}; \\ W_P &= \{(123), (122), (112), (111)\} \cup \{(132), (122), (121), (111)\} \\ &= \{(123), (132), (122), (112), (121), (111)\}. \end{aligned}$$

Note that the two components of W_P are not disjoint.

Remark. A similar result is proved in [10] for T -partitions of a quasi-poset, generalizing Stanley's result [17] for P -partitions of posets; nevertheless, this is different here, as the union is not a disjoint one.

4 Characters associated to ehr and ehr^{str}

Recall that $(M_{\mathbf{qp}}, *)$ the monoid of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$.

By theorems 7 and 23:

Proposition 30 *1. There exists a unique homogeneous Hopf algebra morphism ϕ_0 from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$ such that:*

$$\forall n \geq 1, \phi_0(\bullet_n) = X.$$

There exists a unique character $\lambda \in M_{\mathbf{qp}}$ such that for all $P \in \mathbf{qp}$,

$$\phi_0(P) = \lambda_P X^{cl(P)}.$$

2. There exist unique characters $\alpha, \alpha^{str} \in M_{\mathbf{qp}}$, such that:

$$ehr = \phi_0 \leftarrow \alpha, \quad ehr^{str} = \phi_0 \leftarrow \alpha^{str}.$$

Remarks.

1. Let $P \in \mathbf{qp}$. Then $ehr_P = ehr_{\overline{P}}$ and $ehr_P^{str} = ehr_{\overline{P}}^{str}$, so $\alpha_P = \alpha_{\overline{P}}$ and $\alpha_P^{str} = \alpha_{\overline{P}}^{str}$.
2. Still by theorem 7, there exists a unique homogeneous Hopf algebra morphism ϕ'_0 from $\mathcal{H}_{\mathbf{p}}$ to $\mathbb{K}[X]$ such that $\phi'_0(\bullet) = X$. By unicity, $\phi_0 = \phi'_0 \circ \Xi$, so for any $P \in \mathbf{qp}$, $\lambda_P = \lambda_{\overline{P}}$.

4.1 The character λ

Lemma 31 For all $P \in \mathbf{P}(n)$, $n \geq 0$:

$$\lambda_{[P]} = \begin{cases} 1 & \text{if } P = 1, \\ \frac{1}{n} \sum_{M \in \max(P)} \lambda_{[P_{[n] \setminus \{M\}}]} & \text{otherwise.} \end{cases} = \frac{1}{n} \sum_{m \in \min(P)} \lambda_{[P_{[n] \setminus \{m\}}]}$$

Proof. Let $P \in \mathbf{P}(n)$, with $n \geq 0$. Recall that $\pi : \mathbb{K}[X] \rightarrow \text{Vect}(X)$ is the canonical projection.

$$\begin{aligned} (Id \otimes \pi) \circ \Delta \circ \phi_0([P]) &= \lambda_{[P]}(Id \otimes \pi) \circ \Delta(X^n) \\ &= \lambda_{[P]} n X^{n-1}; \\ &= (Id \otimes \pi) \circ (\phi_0 \otimes \phi_0) \circ \Delta([P]) = \sum_{O \in \text{Top}(P)} \lambda_{[P|O]} \lambda_{[P_{[n] \setminus O}]} X^{|[n] \setminus O|} \pi(X^{|O|}) \\ &= \sum_{O \in \text{Top}(P), |O|=1} \lambda_{[P|O]} \lambda_{[P_{[n] \setminus O}]} X^{n-1} \\ &= \sum_{M \in \max(P)} \lambda_{[P_{[n] \setminus \{M\}}]} X^{n-1}. \end{aligned}$$

This implies the first equality. The second is proved by considering $(\pi \otimes Id) \circ \Delta \circ \phi_0([P])$. $\square$

Remarks.

1. This lemma allows to inductively compute λ_P . This gives:

P	$\cdot$	$\downarrow$	$\vee$	$\wedge$	$\downarrow$	$\vee$	$\wedge$	$\downarrow$	$\vee$	$\wedge$	$\downarrow$	$\vee$	$\wedge$	$\downarrow$	$\vee$	$\wedge$
λ_P	1	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{24}$	$\frac{5}{24}$	$\frac{1}{6}$	$\frac{1}{12}$	

2. If $P = (P, \leq)$ is a finite poset, we denote by P^{op} the opposite poset $(P, \geq)$. It is not difficult to deduce from this lemma that $\lambda_P = \lambda_{P^{op}}$.

Proposition 32 Let $P \in \mathbf{P}(n)$. The number of elements of $W_P(n)$ of P is denoted by μ_P : in other words, μ_P is the number of bijections f from $[n]$ to $[n]$ such that for all $x, y \in [n]$,

$$x \leq_P y \implies f(x) \leq f(y).$$

These bijections are called *heap-orderings* of P . Then, for any finite poset P , $\lambda_P = \frac{\mu_P}{n!}$.

Proof. Let us fix a non-empty finite poset $P \in \mathbf{P}(n)$. The set of heap-orderings of P is $HO(P) = W_P(n)$. We consider the map:

$$\begin{cases} HO(P) & \longrightarrow \bigsqcup_{M \in \max(P)} HO(P \setminus \{M\}) \\ f & \longrightarrow f_{[n-1]} \in HO(P \setminus \{f^{-1}(n)\}). \end{cases}$$

It is not difficult to prove that this is a bijection. So:

$$\mu_P = \sum_{M \in \max(P)} \mu_{P \setminus \{M\}}; \quad \frac{\mu_P}{n!} = \frac{1}{n} \sum_{M \in \max(P)} \frac{\mu_{P \setminus \{M\}}}{|P \setminus \{M\}|!}.$$

An easy induction on $|P|$ then proves that $\lambda_P = \frac{\mu_P}{n!}$ for all P . $\square$

This formula can be simplified for rooted forests.

Definition 33 Let P be a non-empty finite poset.

1. We put:

$$P! = \prod_{i \in V(P)} \# \{j \in V(P) \mid i \leq_P j\}.$$

By convention, $1! = 1$.

2. We shall say that P is a rooted forest if P does not contain any subposet isomorphic to $\mathbb{A}$.

For example, here are isoclasses of rooted forests of cardinality ≤ 4 :

$$1; \quad \cdot; \quad \mathbf{1}, \dots; \quad \mathbb{V}, \mathbf{1}, \dots, \mathbf{1}, \dots; \quad \mathbb{V}, \mathbf{1}, \mathbb{Y}, \mathbf{1}, \mathbb{V}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \dots, \dots$$

Examples.

P															
$P!$	1	2	3	4	6	4	8	8	12	12	18	24	6	9	16

Proposition 34 For all finite poset P , $\lambda_P \geq \frac{1}{P!}$, with equality if, and only if, P is a rooted forest.

Proof. We proceed by induction on $n = |P|$. It is obvious if $n = 0$. Let us assume the result at all ranks $< n$.

$$\begin{aligned} \lambda_P &= \frac{1}{|P|} \sum_{m \in \min(P)} \lambda_{P \setminus \{m\}} \\ &\geq \frac{1}{|P|} \sum_{m \in \min(P)} \prod_{i \in V(P), i \neq m} \frac{1}{\# \{j \in V(P) \mid j \neq m, i \leq_P j\}} \\ &= \frac{1}{|P|} \sum_{m \in \min(P)} \prod_{i \in V(P), i \neq m} \frac{1}{\# \{j \in V(P) \mid i \leq_P j\}} \\ &= \frac{1}{P!} \frac{1}{|P|} \sum_{m \in \min(P)} \# \{j \in V(P) \mid m \leq_P j\}. \end{aligned}$$

For any $j \in A$, there exists $m \in \min(P)$ such that $m \leq_P j$, so:

$$\sum_{m \in \min(P)} \# \{j \in V(P) \mid m \leq_P j\} \geq |P|.$$

Consequently, $\lambda_P \geq \frac{1}{P!}$.

Let us assume that this is an equality. Then:

$$\sum_{m \in \min(P)} \# \{j \in V(P) \mid m \leq_P j\} = |P|.$$

Consequently, for all $j \in \min(P)$, there exists a unique $m \in \min(P)$ such that $m \leq_P j$. Moreover, for all $m \in \min(P)$, $\lambda_{P \setminus \{m\}} = \frac{1}{P \setminus \{m\}!}$. By the induction hypothesis, $P \setminus \{m\}$ is a rooted forest; this implies that P is also a rooted forest.

Let us assume that P is a rooted forest. For any $j \in V(P)$, there exists a unique $m \in \min(P)$ such that $m \leq_P j$, so:

$$\sum_{m \in \min(P)} \# \{j \in V(P) \mid m \leq_P j\} = |P|.$$

Moreover, for all $m \in \min(P)$, $P \setminus \{m\}$ is also a rooted forest. By the induction hypothesis, $\lambda_{P \setminus \{m\}} = \frac{1}{P \setminus \{m\}!}$. Hence, $\lambda_P = \frac{1}{P!}$. $\square$

4.2 The character α^{str}

Let us now apply theorem 8 to ehr and ehr^{str} :

Theorem 35 *For all finite connected quasi-poset P , we have:*

$$\alpha_P = \frac{d \, ehr_P}{dX}(0), \quad \alpha_P^{str} = \frac{d \, ehr_P^{str}}{dX}(0).$$

For any quasi-poset P :

$$ehr_P(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim} X^{cl(\sim)}, \quad ehr_P^{str}(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim}^{str} X^{cl(\sim)},$$

where $cl(\sim)$ is the number of equivalence classes of $\sim$.

Let us give a few values of α :

P	$\cdot$	$\mathfrak{I}$	$\mathfrak{V}$	$\mathfrak{A}$	$\mathfrak{I}$	$\mathfrak{V}$	$\mathfrak{A}$	$\mathfrak{I}$	$\mathfrak{V}$	$\mathfrak{A}$	$\mathfrak{I}$	$\mathfrak{V}$	$\mathfrak{A}$	$\mathfrak{I}$	$\mathfrak{V}$	$\mathfrak{A}$
α_P	1	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	0	0	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{12}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

Lemma 36 *Let $P \in \mathbf{QP}$, not discrete. Then:*

$$(-1)^{cl(P)} ehr_P(-1) = ehr_P^{str}(1) = \begin{cases} 1 & \text{if } P \text{ is discrete,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If P is discrete, then $ehr_P(X) = ehr_P^{str}(X) = X^{cl(P)}$ and the result is obvious. Let us assume that P is not discrete. There exists a unique map f from $[n]$ to $[1]$; as P is not discrete, $f \notin L_P^{str}(1)$, so $ehr_P^{str}(1) = 0$. We now proceed with $ehr_P(-1)$.

First step. Let us prove that $L(H_k(-X)) = -H_{k+1}(-X)$ for all $k \geq 0$. For all $l, n \geq 0$:

$$H_l(-n) = (-1)^l \binom{n+l-1}{l}.$$

For all $k, n \geq 0$:

$$\begin{aligned} L(H_k(-X))(n+1) &= H_k(0) + \dots + H_k(-n) \\ &= (-1)^k \sum_{i=0}^n \binom{i+k-1}{k} \\ &= (-1)^k \sum_{j=k}^{n+k-1} \binom{j}{k} \\ &= (-1)^k \binom{n+k}{k+1} \\ &= -H_{k+1}(-(n+1)). \end{aligned}$$

Second step. Let us prove that $L((X+1)\mathbb{K}[X]) \subseteq (X+1)\mathbb{K}[X]$. For all $k \geq 2$, let us put $H_k(-X) = X(X+1)L_k(X)$; $(L_k(X))_{k \geq 2}$ is a basis of $\mathbb{K}[X]$, which implies that $(H_k(-X))_{k \geq 2}$ is a basis of $X(X+1)\mathbb{K}[X]$, and that $(X+1) \sqcup (H_k(-X))_{k \geq 2}$ is a basis of $(X+1)\mathbb{K}[X]$. First:

$$L(X+1) = L(H_1(X) + H_0(X)) = H_2(X) + H_1(X) = \frac{X(X-1)}{2} + X = \frac{X(X+1)}{2} \in (X+1)\mathbb{K}[X];$$

if $k \geq 2$, by the first step, $L(H_k(-X)) = -H_{k+1}(-X) \in (X+1)\mathbb{K}[X]$.

Last step. We can replace P by $\overline{P}$, and we now assume that $P \in \mathbf{P}(n)$. There is nothing to prove if $n = 0, 1$. Let us assume the result at all rank $< n$. Then, by the second step and the induction hypothesis:

$$\begin{aligned}
ehr_{[P]}(-1) &= L \left(\sum_{\emptyset \neq O \in Top(P)} ehr_{[P_{[n] \setminus O}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{\substack{\emptyset \neq O \in Top(P) \\ P_{[n] \setminus O} \text{ discrete}}} ehr_{[P_{[n] \setminus O}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{[n] \neq J \subseteq min(P)} ehr_{[P_{|J}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{J \subseteq min(P)} ehr_{[P_{|J}]} J(X) \right)_{|X=-1} \\
&= L \left(\sum_{J \subseteq min(P)} X^{|J|} \right)_{|X=-1} \\
&= L \left(\underbrace{(1+X)^{|min(P)|}}_{\in (X+1)\mathbb{K}[X]} \right)_{|X=-1} \\
&= 0.
\end{aligned}$$

For the fourth equality, note that P is not discrete, so $min(P) \neq P$. □

In other words, for any $P \in \mathbf{qp}$, $(-1)^{cl(P)} ehr_P(-1) = ehr_P^{str}(1) = \varepsilon'(P)$. By corollary 10:

Theorem 37 *ehr^{str} is the unique morphism from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$ such that:*

1. *ehr^{str} is a Hopf algebra morphism from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$.*
2. *ehr^{str} is a bialgebra morphism from $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ to $(\mathbb{K}[X], m, \delta)$.*

Moreover, the character α^{str} is the inverse of λ in M_b .

4.3 The character α and the duality principle

Theorem 38 1. *(Duality principle). For any quasi-poset P :*

$$ehr_P^{str}(X) = (-1)^{cl(P)} ehr_P(-X).$$

2. *For any quasi-poset P , $\alpha_P = (-1)^{cl(P)+cc(P)} \alpha_P^{str}$.*

3. *α is invertible in $M_{\mathbf{qp}}$. We denote by β its inverse. For any quasi-poset P :*

$$\beta_P = (-1)^{cl(P)+cc(P)} \lambda_P = (-1)^{cl(P)+cc(P)} \frac{\mu_P}{cl(P)!}.$$

Proof. 1. We consider the morphism:

$$\phi : \begin{cases} \mathcal{H}_{\mathbf{qp}} & \longrightarrow \mathbb{K}[X] \\ P \in \mathbf{qp} & \longrightarrow (-1)^{cl(P)} ehr_P(-X). \end{cases}$$

We put:

$$\theta_1 : \begin{cases} \mathcal{H}_{\mathbf{qp}} & \longrightarrow \mathcal{H}_{\mathbf{qp}} \\ P \in \mathbf{qp} & \longrightarrow (-1)^{cl(P)} P, \end{cases} \quad \theta_2 : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K}[X] \\ P(X) & \longrightarrow P(-X). \end{cases}$$

Both are Hopf algebra morphisms, and $\phi = \theta_2 \circ ehr \circ \phi_1$, so ϕ is a Hopf algebra morphism. If P is a non discrete quasi-poset, then $\phi(P)(1) = (-1)^{cl(P)} ehr_P(-1) = 0 = \varepsilon'(x)$. If P is a discrete quasi-poset, then $ehr_P(X) = X^{cl(P)}$, so $\phi(P)(1) = 1 = \varepsilon'(x)$. By corollary 10, $\phi = \phi_1 = ehr^{str}$.

2. and 3. Immediate consequences of the first point. □

Proposition 39 *The following map is a Hopf algebra automorphism:*

$$\theta : \begin{cases} (\mathcal{H}_{\mathbf{qp}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} P / \sim. \end{cases}$$

Its inverse is:

$$\theta^{-1} : \begin{cases} (\mathcal{H}_{\mathbf{qp}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} (-1)^{cl(\sim) + cl(P)} P / \sim. \end{cases}$$

Moreover:

$$ehr^{str} \circ \theta = ehr, \quad ehr \circ \theta^{-1} = ehr^{str}.$$

Proof. Let ι be the character of $\mathcal{H}_{\mathbf{qp}}$ which sends any quasi-poset to 1. By corollary 20, θ is an automorphism and $\theta^{-1} = \phi_{\iota^* - 1}$. For any quasi-poset P :

$$\iota(1) = 1 = ehr_P(1) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(P/\sim)!} \alpha_{P/\sim} = \sum_{\sim \triangleleft P} \lambda_{P/\sim} \alpha_{P/\sim} = \lambda * \alpha(P),$$

so $\iota = \lambda * \alpha$; hence, its inverse is $\beta * \alpha^{str}$, and for any quasi-poset P , as $cl(P/\sim) = cl(\sim)$ and $cc(P/\sim) = cc(P)$ for any $\sim \triangleleft P$:

$$\begin{aligned} \beta * \alpha^{str}(P) &= \sum_{\sim \triangleleft P} (-1)^{cl(\sim) + cc(P)} \frac{\mu_{P/\sim}}{cl(P/\sim)!} \alpha_{P/\sim}^{str} \\ &= (-1)^{cc(P)} ehr_P^{str}(-1) \\ &= (-1)^{cc(P) + cl(P)} ehr^{str}(1) \\ &= (-1)^{cc(P) + cl(P)}. \end{aligned}$$

Hence:

$$\theta^{-1}(P) = Id \leftarrow (\beta * \alpha^{str})(P) = \sum_{\sim \triangleleft P} (-1)^{cc(P/\sim) + cl(P/\sim)} P / \sim = \sum_{\sim \triangleleft P} (-1)^{cl(\sim) + cl(P)} P / \sim.$$

Moreover:

$$\begin{aligned} ehr^{str} \circ \theta &= (\phi_0 \leftarrow \alpha^{str}) \circ (Id \leftarrow \iota) \\ &= ((\phi_0 \leftarrow \alpha^{str}) \circ Id) \leftarrow \iota \\ &= (\phi_0 \leftarrow \alpha^{str}) \leftarrow \iota \\ &= \phi_0 \leftarrow (\alpha^{str} * \iota) \\ &= \phi_0 \leftarrow (\alpha^{str} * \lambda * \alpha) \\ &= \phi_0 \leftarrow \alpha \\ &= ehr. \end{aligned}$$

□

4.4 A link with Bernoulli numbers

For any $k \in \mathbb{N}$, let c_k be the corolla quasi-poset with k leaves: $c_k = ([k+1], \leq_{c_k})$, with $1 \leq_{c_k} 2, \dots, k+1$:

$$c_0 = \bullet_1, \quad c_1 = \mathbf{1}_1^2, \quad c_2 = {}^2\mathbf{V}_1^3, \quad c_3 = {}^2\mathbf{V}_1^4 \dots$$

By proposition 34, $\lambda_{c_k} = \frac{1}{k+1}$. Moreover:

$$\begin{aligned} L_{c_k} &= \{f : [k+1] \longrightarrow \mathbb{N}^* \mid f(1) \leq f(2), \dots, f(k+1)\}, \\ L_{c_k}^{str} &= \{f : [k+1] \longrightarrow \mathbb{N}^* \mid f(1) < f(2), \dots, f(k+1)\}, \end{aligned}$$

so, for all $n \geq 1$:

$$Ehr_{c_k}^{str}(n) = (n-1)^k + \dots + 1^k = S_k(n),$$

where $S_k(X)$ is the unique polynomial such that for all $n \geq 1$, $S_k(n) = 1^k + \dots + (n-1)^k$. As a consequence, $\alpha_{c_k}^{str}$ is equal to the k -th Bernoulli number b_k .

Let $\sim \triangleleft c_k$. As the equivalence classes of $\sim$ are connected:

- The equivalence class of the minimal element 1 of c_k contains i leaves, $0 \leq i \leq k$.
- The other equivalence classes are formed by a unique leaf.

Hence:

$$\delta([c_k]) = \sum_{i=0}^k \binom{k}{i} [c'_{i,k-i}] \otimes [c_i] \bullet^{k-i},$$

where $c'_{i,k-i}$ is the quasi poset on $[k+1]$ such that:

$$1 \sim_{c'_{i,k-i}} \dots \sim_{c'_{i,k-i}} i+1 \leq_{c'_{i,k-i}} i+2, \dots, k+1.$$

Hence, by theorem 35:

$$\begin{aligned} S_k(X) &= \sum_{i=0}^k \binom{k}{i} \lambda_{c'_{i,k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \lambda_{c'_{i,k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \lambda_{c_{k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \frac{b_i}{k-i+1} X^{k-i+1}. \end{aligned}$$

We recover in this way Faulhaber's formula. For all $n \geq 1$, $ehr_{c_k}(n) = n^k + \dots + 1^k$, and the duality principle gives, for all $n \geq 1$:

$$(-1)^{k+1} S_k(-n) = 1^k + \dots + n^k = S_k(n) + n^k.$$

5 Noncommutative version

5.1 Reminders on packed words

Let us recall the construction of the Hopf algebra of packed words **WQSym** [15, 16].

Definition 40 Let $w = x_1 \dots x_n$ be a word which letters are positive integers.

1. We shall say that w is a packed word if there exists an integer k such that $\{x_1, \dots, x_n\} = [k]$. The set of packed words of length n is denoted by $\mathbf{PW}(n)$; the set of all packed words is denoted by $\mathbf{PW}$.
2. There exists a unique increasing bijection $f : \{x_1, \dots, x_n\} \rightarrow [k]$ for a well-chosen k . We denote by $\text{Pack}(w)$ the packed word $f(x_1) \dots f(x_k)$. Note that w is packed if, and only if, $w = \text{Pack}(w)$.
3. Let $I \subseteq \mathbb{N}$. Let $i_1 < \dots < i_p$ be the indices i such that $x_i \in I$. We denote by w_I the word $x_{i_1} \dots x_{i_p}$.

As a vector space, **WQSym** is generated by the set $\mathbf{PW}$. The product is given by:

$$\forall u \in \mathbf{PW}(k), \forall v \in \mathbf{PW}(l), u.v = \sum_{\substack{w=x_1 \dots x_{k+l} \in \mathbf{PW}(k+l), \\ \text{Pack}(x_1 \dots x_k)=u, \\ \text{Pack}(x_{k+1} \dots x_{k+l})=v}} w.$$

The unit is the empty word 1. The coproduct is given by:

$$\forall w \in \mathbf{PW}, \Delta(w) = \sum_{k=0}^{\max(w)} w_{\{1, \dots, k\}} \otimes \text{Pack}(w_{\{k+1, \dots, \max(w)\}}).$$

For example:

$$\begin{aligned} (11).(11) &= (1111) + (1122) + (2211), \\ (11).(12) &= (1112) + (1123) + (2212) + (2213) + (3312), \\ (11).(21) &= (1121) + (1132) + (2231) + (3321), \\ (12).(11) &= (1211) + (1222) + (1233) + (1322) + (2311), \\ (12).(12) &= (1212) + (1213) + (1223) + (1234) + (1323) + (1324) \\ &\quad + (1423) + (2312) + (2313) + (2314) + (2413) + (3412), \\ (12).(21) &= (1221) + (1231) + (1232) + (1243) + (1332) + (1342) \\ &\quad + (1432) + (2321) + (2331) + (2341) + (2431) + (3421), \end{aligned}$$

$$\begin{aligned} \Delta(111) &= (111) \otimes 1 + (111) \otimes 1, \\ \Delta(212) &= (212) \otimes 1 + (1) \otimes (11) + 1 \otimes (212), \\ \Delta(312) &= (312) \otimes 1 + (1) \otimes (21) + (12) \otimes (1) + 1 \otimes (312). \end{aligned}$$

5.2 Hopf algebra morphisms in WQSym

Proposition 41 The two following maps are surjective Hopf algebra morphisms:

$$\begin{aligned} EHR : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow \mathbf{WQSym} \\ P & \longrightarrow \sum_{w \in W_P} w, \end{cases} \\ EHR^{str} : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow \mathbf{WQSym} \\ P & \longrightarrow \sum_{w \in W_P^{str}} w. \end{cases} \end{aligned}$$

Proof. Let $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, and w be a packed word of length $k + l$. Then:

- $w \in W_{PQ}$ if, and only if, $Pack(w_1 \dots w_k) \in W_P$ and $Pack(w_{k+1} \dots w_{k+l}) \in W_Q$.
- $w \in W_{PQ}^{str}$ if, and only if, $Pack(w_1 \dots w_k) \in W_P^{str}$ and $Pack(w_{k+1} \dots w_{k+l}) \in W_Q^{str}$.

This implies that :

$$EHR(PQ) = EHR(P)EHR(Q), \quad EHR^{str}(PQ) = EHR^{str}(P)EHR^{str}(Q).$$

Let $P \in \mathbf{QP}(n)$. We consider the two sets:

$$\begin{aligned} A &= \{(w, k) \mid w \in W_P, 0 \leq k \leq \max(w)\}, \\ B &= \{(O, w_1, w_2) \mid O \in Top(P), w_1 \in W_{Pack(P_{[n] \setminus O})}, w_2 \in W_{Pack(P|_O)}\}. \end{aligned}$$

We define a bijection between A and B by $F(w, k) = (O, w_1, w_2)$, where:

- $O = w^{-1}(\{k + 1, \dots, \max(w)\})$.
- $w_1 = Pack(w_{\{1, \dots, k\}})$.
- $w_2 = Pack(w_{\{k+1, \dots, \max(w)\}})$.

Then:

$$\begin{aligned} \Delta \circ EHR(P) &= \sum_{(w, k) \in A} Pack(w_{\{1, \dots, k\}}) \otimes Pack(w_{\{k+1, \dots, \max(w)\}}) \\ &= \sum_{(O, w_1, w_2) \in B} w_1 \otimes w_2 \\ &= (EHR \otimes EHR) \circ \Delta(P). \end{aligned}$$

So EHR is a Hopf algebra morphism. In the same way, EHR^{str} is a Hopf algebra morphism.

Let w be a packed word of length n . We define a quasi-poset structure on $[n]$ by $i \leq_P j$ if, and only if, $w_i \leq w_j$. Then $W_P^{str} = \{w\}$, so $EHR^{str}(P) = w$: EHR^{str} is surjective. If $w' \in W_P$, then $\max(w') \leq \max(w)$ with equality if, and only if, $w = w'$. Hence:

$$EHR(P) = w + \text{words } w' \text{ with } \max(w') < \max(w).$$

By a triangular argument, EHR is surjective. □

Examples.

$$\begin{aligned} EHR(\bullet_1) &= (1), & EHR^{str}(\bullet_1) &= (1), \\ EHR(\mathbf{1}_1^2) &= (12) + (11), & EHR^{str}(\mathbf{1}_1^2) &= (12), \\ EHR(\mathbf{1}_2^1) &= (21) + (11), & EHR^{str}(\mathbf{1}_2^1) &= (21), \\ EHR(\bullet_1 \bullet_2) &= (12) + (21) + (11), & EHR^{str}(\bullet_1 \bullet_2) &= (12) + (21) + (11), \\ EHR(\bullet_{1,2}) &= (11), & EHR^{str}(\bullet_{1,2}) &= (11). \end{aligned}$$

Remark. The Hopf algebra $\mathbf{WQSym}$ has a polynomial representation [16]. Let $X = \{x_1, x_2, \dots\}$ be an infinite, totally ordered alphabet; then, for any packed word w :

$$P_w(X) = \sum_{w' \in X^*, Pack(w')=w} w'.$$

With this polynomial representation, for any $P \in \mathbf{QP}(n)$:

$$EHR(P) = \sum_{f \in L_P} x_{f(1)} \dots x_{f(n)}, \quad EHR^{str}(P) = \sum_{f \in L_P^{str}} x_{f(1)} \dots x_{f(n)}.$$

Proposition 42 *The following map is a Hopf algebra automorphism:*

$$\Theta : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} P / \sim. \end{cases}$$

Its inverse is:

$$\Theta^{-1} : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} (-1)^{cl(P)+cl(\sim)} P / \sim. \end{cases}$$

Moreover, $EHR^{str} \circ \Theta = EHR$ and $EHR \circ \Theta^{-1} = EHR^{str}$.

Proof. By corollary 20, $\Theta = \phi_\iota$ is a Hopf algebra automorphism, where ι is defined in the proof of proposition 39. Its inverse is $\phi_{\iota^{-1}}$.

Let us prove that:

$$W_G = \sum_{\sim \triangleleft P} W_{G/\sim}^{str}.$$

Let $w \in W_G$; we define an equivalence $\sim_w$ by $x \sim_w y$ if $w(x) = w(y)$ and x and y are in the same connected component of $w^{-1}(w(x))$. By definition, the equivalence classes of $\sim_w$ are connected. If $x \sim_{P/\sim_w} y$, there exists $x_1, x'_1, \dots, x_k, x'_k, y_1, y'_1, \dots, y_l, y'_l$ such that:

$$\begin{aligned} x &\leq_P x_1 \sim_w x'_1 \leq_P \dots \leq_P x_k \sim_w x'_k \leq_P y, \\ y &\leq_P y_1 \sim_w y'_1 \leq_P \dots \leq_P y_l \sim_w y'_l \leq_P x. \end{aligned}$$

As $w \in W_P$, $w(x) \leq w(x_1) = w(x'_1) \leq \dots \leq w(x_k) = w(x'_k) \leq w(y)$; by symmetry, $w(x) = w(x_1) = \dots = w(x'_k) = w(y) = i$. Moreover, as the equivalence classes of $\sim_w$ are connected, x and y are in the same connected component of $w^{-1}(i)$, so $x \sim_w y$: $\sim_w \triangleleft P$.

If $x \leq_P y$ or $x \sim_w y$, then $w(x) \leq w(y)$. By transitive closure, if $x \leq_{P/\sim_w} y$, then $w(x) \leq w(y)$, so $w \in W_{P/\sim_w}$. Moreover, if $w(i) \neq w(j)$, we do not have $x \sim_w y$, so $w \in W_{P/\sim_w}^{str}$.

Let us assume that $\sim \triangleleft P$ and let $w \in W_{P/\sim}^{str}$. If $x \leq_P y$, then $x \leq_{P/\sim} y$, so $w(x) \leq w(y)$: $W_{P/\sim}^{str} \subseteq W_P$.

Let us assume that $w \in W_{P/\sim}^{str}$, with $\sim \triangleleft P$. If $x \sim y$, then $w(x) = w(y) = i$ and x and y are in the same connected component of $P| \sim$, so are in the the same connected component of $w^{-1}(i)$: $x \sim_w y$. If $x \sim_w y$, then $w(x) = w(y) = i$ and there exists $x_1, x'_1, \dots, x_k, x'_k$ with $w(x_1) = w(x'_1) = \dots = w(x_k) = w(x'_k) = i$ such that:

$$x \leq_P x_1 \geq_P x'_1 \leq_P \dots \geq_P x'_k \leq_P y.$$

As $w \in W_{P/\sim}^{str}$, $x \sim_{P/\sim} x_1$, $x_1 \sim_{P/\sim} x'_1, \dots, x'_k \sim_{P/\sim} y$. So $x \sim_{P/\sim} y$; as $\sim \triangleleft P$, $x \sim y$. Finally, $\sim = \sim_w$.

We obtain that:

$$EHR(P) = \sum_{w \in W_P} w = \sum_{\sim \triangleleft P} \sum_{w \in W_{P/\sim}^{str}} w = \sum_{\sim \triangleleft P} EHR^{str}(P / \sim) = EHR^{str}(\Theta(P)).$$

So $Ehr^{str} \circ \Theta = EHR$. □

Proposition 43 *Let us consider the following map:*

$$H : \begin{cases} \mathbf{WQSym} & \longrightarrow \mathbb{K}[X] \\ w \in \mathbf{PW} & \longrightarrow H_{\max(w)}(X). \end{cases}$$

This is a surjective Hopf algebra morphism, making the following diagram commuting:

$$\begin{array}{ccccc}
& & \mathcal{H}_{\mathbf{QP}} & & \\
& \swarrow \sqcup & \downarrow \Theta & \searrow EHR & \\
\mathcal{H}_{\mathbf{QP}} & & \mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR^{str}} & \mathbf{WQSym} \\
\downarrow \theta \downarrow ehr & \swarrow \sqcup & & \searrow H & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{ehr^{str}} & \mathbb{K}[X] & &
\end{array}$$

Proof. Let $P \in \mathbf{QP}$. Then:

$$ehr(\lfloor P \rfloor) = \sharp W_P(k) H_k(X) = \sum_{w \in W_P} H_{\max(w)}(X) = \sum_{w \in W_P} H(w) = H \circ EHR(P).$$

So $ehr \circ \lfloor \rfloor = H \circ EHR$. Similarly, $ehr^{str} \circ \lfloor \rfloor = H \circ EHR^{str}$.

Let us prove that H is a Hopf algebra morphism. Let $w_1, w_2 \in \mathbf{WQSym}$. There exist $x_1, x_2 \in \mathcal{H}_{\mathbf{QP}}$, such that $w_1 = EHR(x_1)$ and $w_2 = EHR(x_2)$. Then:

$$\begin{aligned}
H(w_1 w_2) &= H(EHR(x_1) EHR(x_2)) \\
&= H \circ EHR(x_1 x_2) \\
&= ehr(\lfloor x_1 x_2 \rfloor) \\
&= ehr(\lfloor x_1 \rfloor) ehr(\lfloor x_2 \rfloor) \\
&= H \circ EHR(x_1) H \circ EHR(x_2) \\
&= H(w_1) H(w_2).
\end{aligned}$$

Let $w \in \mathbf{WQSym}$. There exists $x \in \mathcal{H}_{\mathbf{QP}}$ such that $w = EHR(x)$.

$$\begin{aligned}
\Delta \circ H(w) &= \Delta \circ H \circ EHR(x) \\
&= (H \otimes H) \circ (EHR \otimes EHR) \circ \Delta(x) \\
&= (H \otimes H) \circ \Delta \circ EHR(x) \\
&= (H \otimes H) \circ \Delta(w).
\end{aligned}$$

So H is a Hopf algebra morphism. □

5.3 The non-commutative duality principle

Lemma 44 *The following map is an involution and a Hopf algebra automorphism:*

$$\Phi_{-1} : \begin{cases} \mathbf{WQSym} & \longrightarrow \mathbf{WQSym} \\ w & \longrightarrow (-1)^{\max(w)} \sum_{\sigma : [\max(w)] \rightarrow [l], \text{ non-decreasing}} \sigma \circ w. \end{cases}$$

Proof. Using the surjective morphisms EHR^{str} and ehr^{str} , taking the quotients of the cointeracting bialgebras $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$, we obtain that $(\mathbf{WQSym}, m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ are cointeracting bialgebras, with the coaction defined by:

$$\rho = (Id \otimes H) \circ \delta : \mathbf{WQSym} \longrightarrow \mathbf{WQSym} \otimes \mathbb{K}[X]$$

For any packed word w :

$$\rho(w) = \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} \sigma \circ w \otimes H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(1)}))}(X) \dots H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(l)}))}(X).$$

Using proposition 4, for any $\lambda \in \mathbb{K}$, considering the character:

$$ev_\lambda : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K} \\ P & \longrightarrow P(\lambda), \end{cases}$$

we obtain an endomorphism Φ_λ of $(\mathbf{WQSym}, m, \Delta)$ defined by $\Phi_\lambda = Id \leftarrow ev_\lambda$. if $\lambda \neq 0$, Φ_λ is invertible, of inverse $\Phi_{\lambda^{-1}}$. For any packed word w , denoting by k its maximum:

$$\Phi_\lambda(w) = \sum_{\sigma : [k] \twoheadrightarrow [l], \text{ non-decreasing}} H_{\max(Pack(w|_{(\sigma \circ w)^{-1}(1)}))}(\lambda) \dots H_{\max(Pack(w|_{(\sigma \circ w)^{-1}(l)}))}(\lambda) \sigma \circ w.$$

In particular, for $\lambda = -1$, for any $p \in \mathbb{N}$:

$$H_p(-1) = \frac{(-1)(-2) \dots (-k)}{k!} = (-1)^k.$$

Hence:

$$\begin{aligned} \Phi_{-1}(w) &= \sum_{\sigma : [k] \twoheadrightarrow [l], \text{ non-decreasing}} (-1)^{\max(Pack(w|_{(\sigma \circ w)^{-1}(1)})) + \dots + \max(Pack(w|_{(\sigma \circ w)^{-1}(l)}))} \sigma \circ w \\ &= (-1)^k \sum_{\sigma : [k] \twoheadrightarrow [l], \text{ non-decreasing}} \sigma \circ w. \end{aligned}$$

Indeed, if $x \in (\sigma \circ w)^{-1}(p)$ and $y \in (\sigma \circ w)^{-1}(q)$, with $p < q$, then $\sigma \circ w(x) < \sigma \circ w(y)$; as σ is non-decreasing, $x < y$. So there exists $n_1 < n_2 < \dots < n_l = k$ such that for all p , the values taken by w on $(\sigma \circ w)^{-1}(p)$ are $n_{p-1} + 1, \dots, n_p$. Hence, the values taken by $Pack(w|_{(\sigma \circ w)^{-1}(p)})$ are $1, \dots, n_p - n_{p-1}$, so:

$$\max(Pack(w|_{(\sigma \circ w)^{-1}(1)})) + \dots + \max(Pack(w|_{(\sigma \circ w)^{-1}(l)})) = n_1 + n_2 - n_1 + \dots + n_l - n_{l-1} = n_l = k.$$

In particular, Φ_{-1} is an involution and a Hopf algebra automorphism of $(\mathbf{WQSym}, m, \Delta)$. $\square$

Theorem 45 (Non commutative duality principle) *For any quasi-poset $P \in \mathbf{QP}$:*

$$EHR(P) = (-1)^{cl(P)} \Phi_{-1} \circ ERH^{str}(P), \quad EHR^{str}(P) = (-1)^{cl(P)} \Phi_{-1} \circ ERH(P).$$

Proof. We shall use the following involution and Hopf algebra automorphism:

$$\Psi : \begin{cases} \mathcal{H}_{\mathbf{QP}} & \longrightarrow \mathcal{H}_{\mathbf{QP}} \\ p \in \mathbf{QP} & \longrightarrow (-1)^{cl(P)} P. \end{cases}$$

Recall that the character ι of $\mathcal{H}_{\mathbf{QP}}$ sends any $P \in \mathbf{QP}$ to 1. By the duality principle:

$$\iota \circ \Psi(P) = (-1)^{cl(P)} = (-1)^{cl(P)} ehr(P)(1) = ehr^{str}(-1) = ev_{-1} \circ ehr^{str}(P).$$

So $\iota \circ \Psi = ev_{-1} \circ ehr^{str}$.

Let $P \in \mathbf{QP}$. Recalling that if $\sim \triangleleft P$, $cl(P|_{\sim}) = cl(P)$:

$$\delta \circ \Psi(P) = (-1)^{cl(P)} \sum_{\sim \triangleleft P} P / \sim \otimes P | \sim = \sum_{\sim \triangleleft P} P / \sim \otimes (-1)^{cl(P|_{\sim})} P | \sim = (Id \otimes \Psi) \circ \delta(P).$$

So $\delta \circ \Psi = (Id \otimes \Psi) \circ \delta$. Hence, for any $x \in \mathcal{H}_{\mathbf{QP}}$:

$$\begin{aligned} EHR \circ \Psi(x) &= EHR^{str} \circ (Id \leftarrow \iota) \circ \Psi(x) \\ &= EHR^{str}(\Psi(x)_0) \iota \circ \Psi(x)_1 \\ &= EHR^{str}(x_0) \iota \circ \Psi(x_1) \\ &= EHR^{str}(x_0) ev_{-1} \circ ehr^{str}(x_1) \\ &= EHR^{str}(x^{(1)}) ev_{-1} \circ EHR^{str}(x^{(2)}) \\ &= EHR \leftarrow ev_{-1}(x) \\ &= (Id \leftarrow ev_{-1}) \circ EHR^{str}(x) \\ &= \Phi_{-1} \circ EHR^{str}(x), \end{aligned}$$

where we denote $\delta(x) = x^{(1)} \otimes x^{(2)}$ and $\rho(x) = x_0 \otimes x_1$. As Φ_{-1} and Ψ are involutions, $EHR^{str} \circ \Psi = \Phi_{-1} \circ EHR$. $\square$

In $E_{\mathbb{K}[X] \rightarrow \mathbb{K}[X]}$, putting $\phi_\lambda = Id \leftarrow ev_\lambda$, for any $P \in \mathbb{K}[X]$, $\phi_\lambda(P) = P(\lambda X)$. Moreover, as H is compatible with the coactions:

$$H \circ \Phi_\lambda = H \circ (Id \leftarrow ev_\lambda) = H \leftarrow ev_\lambda = (Id \leftarrow ev_\lambda) \circ H = \phi_\lambda \circ H,$$

so:

$$ehr \circ \Psi = H \circ EHR \circ \Psi = H \circ \Phi_{-1} \circ EHR^{str} = \phi_{-1} \circ H \circ EHR^{str} = \phi_{-1} \circ ehr^{str}.$$

In other words, for any $P \in \mathbf{QP}$, $(-1)^{cl(P)} ehr_P(X) = ehr_P^{str}(-X)$: we recover the duality principle.

We obtain the commutative diagram of Hopf algebra morphisms:

$$\begin{array}{ccccc}
\mathcal{H}_{\mathbf{QP}} & & & & \\
\downarrow \theta & \searrow EHR & & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR^{str}} & \mathbf{WQSym} & \sqcup & \\
\downarrow \Psi & & \downarrow \Phi_{-1} & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR} & \mathbf{WQSym} & \sqcup & \\
& & & \sqcup & \\
& & & & \mathcal{H}_{\mathbf{qp}} \\
& & & \searrow H & \downarrow \theta \\
& & & & \mathcal{H}_{\mathbf{qp}} \xrightarrow{ehr} \mathbb{K}[X] \\
& & & \searrow H & \downarrow \psi \\
& & & & \mathcal{H}_{\mathbf{qp}} \xrightarrow{ehr^{str}} \mathbb{K}[X] \\
& & & & \downarrow \phi_{-1} \\
& & & & \mathbb{K}[X]
\end{array}$$

5.4 Compatibility with the other product and coproduct

Theorem 46 We define a second coproduct δ on $\mathbf{WQSym}$:

$$\forall w \in \mathbf{PW}, \delta(w) = \sum_{(\sigma, \tau) \in A_w} (\sigma \circ w) \otimes (\tau \circ w),$$

where A_w is the set of pairs of packed words (σ, τ) of length $\max(w)$ such that:

- σ is non-decreasing.
- If $1 \leq i < j \leq \max(w)$ and $\sigma(i) = \sigma(j)$, then $\tau(i) < \tau(j)$.

Then $(\mathbf{WQSym}, m, \delta)$ is a bialgebra and EHR^{str} is a bialgebra morphism from $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ to $(\mathbf{WQSym}, m, \delta)$.

Proof. Let us prove that $\delta \circ EHR^{str} = (EHR^{str} \otimes EHR^{str}) \circ \delta$. Let $P \in \mathbf{QP}$. We consider the two following sets:

$$\begin{aligned}
A &= \{(\sim, w_1, w_2) \mid \sim \triangleleft P, w_1 \in W_{P/\sim}^{str}, w_2 \in W_{P|\sim}^{str}\}, \\
B &= \{(w, \sigma, \tau) \mid w \in W_P^{str}, (\sigma, \tau) \in A_w\}.
\end{aligned}$$

Let $(\sim, w_1, w_2) \in A$. We put $I_p = w_1^{-1}(p)$ for all $1 \leq p \leq \max(w_1)$, and $w_2^{(p)}$ the standardization of the restriction of w_2 to I_p . We define w by:

$$w(i) = w_2^{(p)}(i) + \max w_1^{(2)} + \dots + \max w_{p-1}^{(2)} \text{ if } i \in I_p.$$

Let us prove that $w \in W_P^{str}$. If $x \leq_P y$, then $x \leq_{P/\sim} y$, so $p = w_1(x) \leq w_2(y) = q$.

- If $p < q$, then $w(x) < w(y)$.
- If $p = q$, then $w_1(x) = w_2(y)$ and, as $x \leq_P y$, x and y are in the same connected component of $w^{-1}(p)$. So $x \sim_{w_1} y$, that is to say $x \sim y$ as $w_1 \in W_{P/\sim}^{str}$, and $x \leq_{P|\sim} y$, which implies that $w_2(x) \leq w_2(y)$ and finally $w(x) \leq w(y)$.

Let us assume that moreover $w(x) = w(y)$. Then $p = q$ and necessarily, $w_2(x) = w_2(y)$. As $w_2 \in W_{P/\sim}^{str}$, $x \sim_{P|\sim} y$, so $x \sim_P y$.

If $w(x) = w(y)$, then by definition of w , $w_1(x) = w_1(y)$. So there exists a unique $\sigma : [\max(w)] \rightarrow [\max(w_1)]$, such that $w_1 = \sigma \circ w$. If $w(x) < w(y)$, then, by construction of w , $w_1(x) \leq w_1(y)$: σ is non-decreasing.

There exists a unique $\tau : [\max(w)] \rightarrow [\max(w_2)]$, such that $w_2 = \tau \circ \sigma$. As $Pack(w|_{I_p}) = Pack((w_2)|_{I_p})$ for all p , τ is increasing on I_p .

To any $(\sim, w_1, w_2) \in A$, we associate $(w, \sigma, \tau) = F(\sim, w_1, w_2) \in B$, such that $w_1 = \sigma \circ \tau$, $w_2 = \tau \circ \sigma$, and $\sim = \sim_{\sigma \circ \tau}$. This defines a map $F : A \rightarrow B$.

Let $(w, \sigma, \tau) \in B$. We put $G(w, \sigma, \tau) = (\sim, \sigma, \tau) = (\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w)$. If $x \leq_P y$, then $w(x) \leq w(y)$, so $w_1(x) = \sigma \circ w(x) \leq \sigma \circ w(y) = w_1(y)$. If moreover $w_1(x) = w_1(y)$, then as $x \leq_P y$, x and y are in the same connected component of $w_1^{-1}(w_1(x))$, so $x \sim_{w_1} y$: $w_1 \in W_{P/\sim}^{str}$.

If $x \leq_{P|\sim} y$, then $x \sim_{w_1} y$ and $x \leq_P y$, so $w_1(x) = w_1(y)$ and $w(x) \leq w(y)$. By hypothesis on τ , $\tau \circ w(x) \leq \tau \circ w(y)$, so $w_2(x) \leq w_2(y)$. If moreover $w_2(x) = w_2(y)$, by hypothesis on τ , $w(x) = w(y)$. As $w \in W_P^{str}$, $x \sim_P y$, so $x \sim_{P|\sim} y$: $w_2 \in W_{P|\sim}^{str}$.

We defined in this way a map $G : B \rightarrow A$. If $(\sim, w_1, w_2) \in A$:

$$G \circ F(\sim, w_1, w_2) = G(w, \sigma, \tau) = (\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w) = (\sim_{w_1}, w_1, w_2) = (\sim, w_1, w_2).$$

So $G \circ F = Id_A$. If $(w, \sigma, \tau) \in B$:

$$F \circ G(w, \sigma, \tau) = F(\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w) = (w, \sigma, \tau).$$

So $F \circ G = Id_B$: F and G are inverse bijections.

We obtain:

$$\begin{aligned} (EHR^{str} \otimes EHR^{str}) \circ \delta(P) &= \sum_{(\sim, w_1, w_2) \in A} w_1 \otimes w_2 \\ &= \sum_{(w, \sigma, \tau) \in B} \sigma \circ w \otimes \tau \circ w \\ &= \sum_{w \in W_P^{str}} \delta(w) \\ &= \delta \circ EHR^{str}(P). \end{aligned}$$

So EHR^{str} is compatible with δ .

As EHR^{str} is compatible with the product m and the coproduct δ , $Ker(EHR^{str})$ is a biideal of $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$, and $(\mathbf{WQSym}, m, \delta)$ is identified with the quotient $\mathcal{H}_{\mathbf{QP}}/Ker(EHR^{str})$, so is a bialgebra. $\square$

Examples.

$$\begin{aligned} \delta(11) &= (11) \otimes (11), \\ \delta(12) &= (12) \otimes ((11) + (12) + (21)) + (11) \otimes (12), \\ \delta(21) &= (21) \otimes ((11) + (12) + (21)) + (11) \otimes (21). \end{aligned}$$

This coproduct δ on **WQSym** is the internal coproduct of [16], dual to the product of the Solomon-Tits algebra.

Remarks.

1. The count of $(\mathbf{WQSym}, m, \delta)$ is given by:

$$\varepsilon_B(w) = \begin{cases} 1 & \text{if } w = (1 \dots 1), \\ 0 & \text{otherwise.} \end{cases}$$

2. There is no coproduct δ' on **WQSym** such that $(EHR \otimes EHR) \circ \delta = \delta' \circ EHR$. Indeed, if δ' is any coproduct on **WQSym**, for $x = \mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_1 \bullet_2 - \bullet_{1,2}$:

$$\delta' \circ EHR(x) = \delta'(0) = 0,$$

but:

$$\begin{aligned} & (EHR \otimes EHR) \circ \delta(x) \\ &= (EHR \otimes EHR)((\mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_1 \bullet_2) \otimes \bullet_{1,2} + \bullet_{1,2} \otimes (\mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_1 \bullet_2 - \bullet_{1,2})) \\ &= (11) \otimes (11). \end{aligned}$$

Proposition 47 $H : (\mathbf{WQSym}, m, \delta) \longrightarrow (\mathbb{K}[X], m, \delta)$ is a bialgebra morphism.

Proof. Let w be a packed word. We denote $k = \max(w)$. Let $a, b \in \mathbb{N}$.

$$\begin{aligned} (H \otimes H) \circ \delta(w)(a, b) &= \sum_{(\sigma, \tau) \in A_w} H_{\max(\sigma \circ w)}(a) H_{\max(\tau \circ w)}(b) \\ &= \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} \binom{a}{l} \binom{b}{|\sigma^{-1}(1)|} \cdots \binom{b}{|\sigma^{-1}(l)|} \\ &= \sum_{\substack{1 \leq l \leq k, \\ i_1 + \dots + i_l = k}} \binom{a}{l} \binom{b}{i_1} \cdots \binom{b}{i_l} \\ &= \binom{ab}{k} \\ &= H_k(ab) \\ &= \delta(H(w))(a, b). \end{aligned}$$

As this is true for any $a, b \in \mathbb{N}$, $(H \otimes H) \circ \delta(w) = \delta \circ H$. □

Definition 48 Let $w = w_1 \dots w_k$ and $w' = w'_1 \dots w'_l$ be two packed words. We put:

$$\begin{aligned} w \downarrow w' &= w_1 \dots w_k (w'_1 + \max(w)) \dots (w'_l + \max(w)), \\ w \circledast w' &= w_1 \dots w_k (w'_1 + \max(w) - 1) \dots (w'_l + \max(w) - 1), \\ w \downarrow\!\!\downarrow w' &= w \downarrow w' + w \circledast w'. \end{aligned}$$

These three products are extended to **WQSym** by bilinearity.

Proposition 49 For all $x, y \in \mathcal{H}_{\mathbf{QP}}$:

$$EHR^{str}(x \downarrow y) = EHR^{str}(x) \downarrow EHR^{str}(y), \quad EHR(x \downarrow\!\!\downarrow y) = EHR(x) \downarrow\!\!\downarrow EHR(y).$$

Proof. Let $P \in \mathbf{QP}(k)$ and $Q \in \mathbf{QP}(l)$. If $w = w_1 \dots w_{k+l}$ is a packed word of length $k+l$:

$$\begin{aligned} w \in W_{P \downarrow Q}^{str} &\iff w_1 \dots w_k \in L_P^{str}, w_{k+1} \dots w_{k+l} \in L_Q^{str}, w_1, \dots, w_k < w_{k+1}, \dots, w_{k+l} \\ &\iff w = w_P \downarrow w_Q, \text{ with } w_P \in W_P^{str}, w_Q \in W_Q^{str}. \end{aligned}$$

So $W_{P \downarrow Q}^{str} = W_P^{str} \downarrow W_Q^{str}$, and:

$$EHR^{str}(P \downarrow Q) = \sum_{w_P \in W_P^{str}, w_Q \in W_Q^{str}} w_P \downarrow w_Q = EHR^{str}(P) \downarrow EHR^{str}(Q).$$

If $w = w_1 \dots w_{k+l}$ is a packed word of length $k+l$:

$$\begin{aligned} w \in W_{P \downarrow Q} &\iff w_1 \dots w_k \in L_P, w_{k+1} \dots w_{k+l} \in L_Q, w_1, \dots, w_k \leq w_{k+1}, \dots, w_{k+l} \\ &\iff w = (w_P \downarrow w_Q, \text{ with } w_P \in W_P, w_Q \in W_Q) \\ &\text{or } w = (w_P \otimes w_Q, \text{ with } w_P \in W_P, w_Q \in W_Q). \end{aligned}$$

Note that these two conditions are incompatible:

- in the first case, $\max(w_1 \dots w_k) = \min(w_{k+1} \dots w_{k+l}) - 1$;
- in the second case, $\max(w_1 \dots w_k) = \min(w_{k+1} \dots w_{k+l})$.

So $W_{P \downarrow Q} = (W_P \downarrow W_Q) \sqcup (W_P \otimes W_Q)$, and:

$$\begin{aligned} EHR(P \downarrow Q) &= \sum_{w_P \in W_P, w_Q \in W_Q} w_P \downarrow w_Q + w_P \otimes w_Q \\ &= EHR(P) \downarrow EHR(Q) + EHR(P) \otimes EHR(Q), \end{aligned}$$

so $EHR(P \downarrow Q) = EHR(P) \sharp EHR(Q)$. □

Remark. As a consequence, $(\mathbf{WQSym}, \downarrow, \Delta)$ and $(\mathbf{WQSym}, \sharp, \Delta)$ are infinitesimal bialgebras [12], as $(\mathcal{H}_{\mathbf{QP}}, \downarrow, \Delta)$ is [11, 10].

Corollary 50 *For all $x, y \in \mathbf{WQSym}$:*

$$\Phi_{-1}(x \downarrow y) = \Phi_{-1}(x) \sharp \Phi_{-1}(y) \quad \Phi_{-1}(x \sharp y) = \Phi_{-1}(x) \downarrow \Phi_{-1}(y).$$

Proof. If $P, Q \in \mathbf{QP}$, then $cl(P \downarrow Q) = cl(P) + cl(Q)$, so:

$$\Psi(P \downarrow Q) = (-1)^{cl(P)+cl(Q)} P \downarrow Q = \Psi(P) \downarrow \Psi(Q).$$

Let $x, y \in \mathbf{WQSym}$. There exist $x', y' \in \mathcal{H}_{\mathbf{QP}}$, such that $EHR^{str}(x') = x$ and $EHR^{str}(y') = y$. Hence, using the non-commutative duality principle:

$$\begin{aligned} \Phi_{-1}(x \downarrow y) &= \Phi_{-1}(EHR^{str}(x') \downarrow EHR^{str}(y')) \\ &= \Phi_{-1} \circ EHR^{str}(x' \downarrow y') \\ &= \Phi_{-1} \circ EHR^{str} \circ \Psi(\Psi(x') \downarrow \Psi(y')) \\ &= EHR(\Psi(x') \downarrow \Psi(y')) \\ &= EHR \circ \Psi(x') \sharp EHR \circ \Psi(y') \\ &= \Phi_{-1}(\Phi_{-1} \circ EHR \circ \Psi(x')) \sharp \Phi_{-1}(\Phi_{-1} \circ EHR \circ \Psi(y')) \\ &= \Phi_{-1}(EHR^{str}(x')) \sharp \Phi_{-1}(EHR^{str}(y')) \\ &= \Phi_{-1}(x) \sharp \Phi_{-1}(y). \end{aligned}$$

As Φ_{-1} is an involution, we obtain the second point. □

5.5 Restriction to posets

In [10], the image of the restriction to $\mathcal{H}_{\mathbf{P}}$ of the map from $\mathcal{H}_{\mathbf{QP}}$ to $\mathbf{WQSym}$ defined by T -partitions is a Hopf subalgebra, isomorphic to the Hopf algebra of permutations $\mathbf{FQSym}$ [13, 7]. This is not the case here:

Proposition 51 $EHR(\mathcal{H}_{\mathbf{P}}) = EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \mathbf{WQSym}$.

Proof. Let w be a packed word of length n . We define a poset P on $[n]$ by:

$$\forall i, j \in [n], i \leq_P j \text{ if } (i = j) \text{ or } (w(i) < w(j)).$$

Note that if $i \leq_P j$, then $w(i) \leq w(j)$. If $i \leq_P j$ and $j \leq_P k$, then:

- if $i = j$ or $j = k$, then obviously $i \leq_P k$.
- Otherwise, $w(i) < w(j)$ and $w(j) < w(k)$, so $w(i) < w(k)$ and $i \leq_P k$.

Let us assume that $i \leq_P j$ and $j \leq_P i$. Then $w(i) \leq w(j)$ and $w(j) \leq w(i)$, so $w(i) = w(j)$. As $i \leq_P j$, $i = j$. So P is indeed a poset, and we observed that $w \in W_P$.

Let w' be a packed word of length n . Let us prove that $w' \in W_P^{str}$ if, and only if, $w \leq w'$, where $\leq$ is the order on packed words defined in definition 28.

$\implies$. Let us assume that $w' \in W_P^{str}$. If $w(i) < w(j)$, then $i \leq_P j$, so $w'(i) \leq w'(j)$. Moreover, if $w'(i) = w'(j)$, then $i \leq_P j$, so $i = j$ as P is a poset, and finally $w(i) = w(j)$: contradiction. So $w'(i) < w'(j)$, we shows that $w \leq w'$.

$\impliedby$. Let us assume that $w' \leq w$. If $i \leq_P j$, then $i = j$ or $w(i) < w(j)$, so $w'(i) = w'(j)$ or $w'(i) < w'(j)$. If, moreover, $w'(i) = w'(j)$, then $i = j$; so $w' \in W_P^{str}$.

We obtain an element $P \in \mathcal{H}_{\mathbf{P}}$ such that:

$$EHR^{str}(P) = \sum_{w \leq w'} w'.$$

As this holds for any w , by a triangularity argument, $EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \mathbf{WQSym}$. By the non-commutative duality principle:

$$EHR(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1} \circ EHR^{str} \circ \Psi(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1} \circ EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1}(\mathbf{WQSym}) = \mathbf{WQSym},$$

as Φ_{-1} is an automorphism of $\mathbf{WQSym}$. $\square$

References

- [1] P. Alexandroff, *Diskrete Räume.*, Rec. Math. Moscou, n. Ser. **2** (1937), 501–519 (German).
- [2] Matthias Beck and Sinai Robins, *Computing the continuous discretely*, second ed., Undergraduate Texts in Mathematics, Springer, New York, 2015, Integer-point enumeration in polyhedra, With illustrations by David Austin.
- [3] Ch. Brouder, *Trees, renormalization and differential equations*, BIT **44** (2004), no. 3, 425–438.
- [4] J. C. Butcher, *An algebraic theory of integration methods*, Math. Comp. **26** (1972), 79–106.
- [5] Damien Calaque, Kurusch Ebrahimi-Fard, and Dominique Manchon, *Two interacting Hopf algebras of trees: a Hopf-algebraic approach to composition and substitution of B-series*, Adv. in Appl. Math. **47** (2011), no. 2, 282–308, arXiv:0806.2238.

- [6] F. Chapoton, *Sur une série en arbres à deux paramètres*, Sémin. Lothar. Combin. **70** (2013), Art. B70a, 20.
- [7] Gérard Duchamp, Florent Hivert, and Jean-Yves Thibon, *Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras*, Internat. J. Algebra Comput. **12** (2002), no. 5, 671–717.
- [8] Frédéric Fauvet, Loïc Foissy, and Dominique Manchon, *The Hopf algebra of finite topologies and mould composition*, arXiv:1503.03820, 2015.
- [9] Loïc Foissy, *Chromatic polynomials and bialgebras of graphs*, in preparation.
- [10] Loïc Foissy and Claudia Malvenuto, *The Hopf algebra of finite topologies and T -partitions*, J. Algebra **438** (2015), 130–169, arXiv:1407.0476.
- [11] Loïc Foissy, Claudia Malvenuto, and Frédéric Patras, *Infinitesimal and B_∞ -algebras, finite spaces, and quasi-symmetric functions*, J. Pure Appl. Algebra **220** (2016), no. 6, 2434–2458, arXiv:1403.7488.
- [12] Jean-Louis Loday and María Ronco, *On the structure of cofree Hopf algebras*, J. Reine Angew. Math. **592** (2006), 123–155, arXiv:math/0405330.
- [13] Claudia Malvenuto and Christophe Reutenauer, *A self paired Hopf algebra on double posets and a Littlewood-Richardson rule*, J. Combin. Theory Ser. A **118** (2011), no. 4, 1322–1333, arXiv:0905.3508.
- [14] Dominique Manchon, *On bialgebras and Hopf algebras or oriented graphs*, Confluentes Math. **4** (2012), no. 1, 1240003, 10, arXiv:1011.3032.
- [15] J.-C. Novelli and J.-Y. Thibon, *Construction of dendriform trialgebras*, C. R. Acad. Sci. Paris **342** (2006), no. 6, 365–446, arXiv:math/0605061.
- [16] ———, *Polynomial realization of some trialgebras*, FPSAC’06 (San Diego) (2006), arXiv:math/0605061v1.
- [17] Richard P. Stanley, *Ordered structures and partitions*, American Mathematical Society, Providence, R.I., 1972, Memoirs of the American Mathematical Society, No. 119.
- [18] R. E. Stong, *Finite topological spaces*, Trans. Amer. Math. Soc. **123** (1966), 325–340.
- [19] David Wright and Wenhua Zhao, *D -log and formal flow for analytic isomorphisms of n -space*, Trans. Amer. Math. Soc. **355** (2003), no. 8, 3117–3141 (electronic), arXiv:math/0209274.