

Grothendieck constants: bounds and comparison

Literature before and after arXiv:2609.07000 · 13 September 2026

The classical constants compare the vector optimum of a finite bilinear form with its scalar optimum, over all matrix sizes and vector dimensions. The scalar variables are signs over $\mathbb{R}$ and unit complex phases over $\mathbb{C}$. Our work gives the following bounds; neither exact constant is determined.

$$1375\pi/2454 \leq K_G^{\mathbb{R}} < 1.779754412112$$
$$1.373 < K_G^{\mathbb{C}} < 1.404698554831$$

Comparison with the literature

The literature columns contain publicly stated results and their direct consequences. “Before” includes the August 2026 real-case results [1]; “after” adds Guo, Fang and Lu [4]. The “Our work” column summarizes unpublished computer-assisted proofs.

Bound	Before the new paper	After the new paper	Our work
Real lower	$6\pi/11$ [1] ≈ 1.713595992867	$6\pi/11$ [1] ≈ 1.713595992867	$1375\pi/2454$ ≈ 1.760264832390
Real upper	$1.781841324234^*[1]$	$1.781841324234^*[1]$	1.779754412112
Complex lower	≈ 1.33807 [3] (Davie)	1.35584631827168 [4]	1.373
Complex upper	1.404909132736 [2] (Haagerup)	1.404909132736 [2] (Haagerup)	1.404698554831

Entries are endpoint values. Displayed decimal upper endpoints are rounded upward; values marked $\approx$ are approximations to the expressions shown. The exact inequalities for our work appear above. Bold entries indicate the strongest bounds in each row.

*Theorem 6.4 of [1] states the real upper bound 1.7818666069360661 . Its Proposition 6.3 and Lemma 6.2, combined with Proposition 6.1, give the stronger comparison endpoint used here:

$$\pi/\{2[0.881573822049 - 0.000011328860 - 14.443/\sqrt{(10 \cdot 251^5)}]\}$$
$$= 1.78184132423347648...$$

Size of the improvement. Relative to these real literature bounds, the lower endpoint increases by 0.04666884 and the upper endpoint decreases by 0.00208691, reducing the interval width by **71.44%**. The complex lower endpoint exceeds [4] by **0.01715368**. Our complex interval is **52.57%** narrower than the Davie-Haagerup interval and **35.39%** narrower than the interval after [4]. The new real upper endpoint removes **7.33%** of our preceding real interval. These percentages compare interval widths, not relative errors in the unknown constants.

Other reported claims. Li et al. [5, §3, Table 2] report internally tested real bounds $51\pi/92 \approx 1.741535057968$ and 1.7813319810625639 , expressly not stated as theorems. Our endpoints are also stronger numerically, by approximately 0.01872977 and 0.00157757, respectively.

Methods and prior work

Real lower bound. The proof bounds the scalar norm of the Gaussian operator $(33/20)P_1 - P_3 - (3/20)P_5$, where P_j projects onto Hermite degree j . For signs f and g , the midpoint functions $h = (f + g)/2$ and $k = (f - g)/2$ satisfy $hk = 0$ and $h^2 + k^2 = 1$. Full transverse majorants, a circle inequality and certified one-dimensional ternary estimates control every Gaussian dimension. Explicit finite approximation then gives matrix witnesses. This shares the agreement/disagreement and Gaussian-coefficient analysis of [1], with a sharper universal inequality and a direct finite-matrix conclusion.

Real upper bound. The construction extends the Gaussian rounding and tensor preprocessing framework of [1]. A five-variable sign function combines six inner and four outer sine harmonics. A signed odd degree-81 polynomial preprocesses the primary correlation; four auxiliary coordinates are reflected. The guarantee is the linear correlation coefficient minus the absolute sum of all nonlinear coefficients. The complete signed Fourier calculation, including both zero-frequency corrections, retains degrees 0–403. Localized fourth- and sixth-derivative estimates bound every remaining odd coefficient from degree 405. The supplied endpoint checker treats the primitive integrations and derivative enclosures as numerical premises.

Complex lower bound. The operator is $P - \lambda I - \beta C_3$, with P the holomorphic linear projection and C_3 the full real cubic chaos, complexified. The Gaussian projection framework and proposed cubic perturbation precede both constructions [4, 6]. Our separate radial weights for real and imaginary components use the exact relation $\text{Re}(h \text{ conj}(k)) = 0$. Pointwise maximization and a finite family of auxiliary weights bound the whole remaining scalar parameter range for one fixed operator. Exact rational certificates and finite approximation yield the bound 1.373.

Relation to Guo-Fang-Lu. Their common radial-weight Gram/Schur criterion estimates the scalar norm within the Gaussian multiplier family shared by the two constructions. Theorem 5.2 of [4] confines that criterion's optimum to (1.35584631827168, 1.35584697425050); this is not an upper bound for the constant or for the entire operator family. Their §5.4 identifies discarded joint phase information. Our anisotropic estimate uses this information and lies outside that restricted criterion. Rotation-character compression places our construction in their multiplier family, so the sharper scalar estimate is the substantive distinction.

Complex upper bound. Three Gaussian coordinates combine signed tensor preprocessing, a mixed linear phase and bounded nonlinear phase perturbations. Complete contour enclosures, zero winding numbers and a homotopy argument justify all scalar Cauchy tails. The calculation retains 9,948 modes, charges every omission and phase remainder, and independently aggregates 2,415,418 coefficient contributions. Exact reweighting of six phase classes further refines the endpoint, which improves Haagerup's bound by approximately 0.000210578. The original complete computation supports the result; a full rerun through the public portable entypoints remains to be completed.

References

- [1] R. Saha et al. *New Lower and Upper Bounds for the Grothendieck Constant*. [arXiv:2608.11158v2](#), §§2, 6-7 (2026).
- [2] U. Haagerup. *A new upper bound for the complex Grothendieck constant*. *Israel J. Math.* 60, 199-224 (1987). [doi:10.1007/BF02790792](#).
- [3] S. Friedland, L.-H. Lim and J. Zhang. *An elementary and unified proof of Grothendieck's inequality*. *L'Enseignement Math.* 64, 327-351 (2018), §1(v). [arXiv:1711.10595](#).
- [4] S. Guo, E. X. Fang and J. Lu. *An Improved Lower Bound for the Complex Grothendieck Constant*. [arXiv:2609.07000v1](#), Theorems 1.1, 3.1, 5.2 and §5.4 (2026).
- [5] A. Li et al. *Long-Horizon AI Research for Grothendieck Constant: A Case Study in Human-AI Mathematical Collaboration*. [arXiv:2608.11195v3](#), §3, Table 2 (2026).
- [6] S. Heilman. *A lower bound for Grothendieck's constant*. [arXiv:2603.22616v1](#), §1.8 (2026).