

Grothendieck constants: current bounds and comparison

Literature before and after arXiv:2609.07000 · 11 September 2026

The classical constants compare the vector optimum of a finite bilinear form with its scalar optimum, over all matrix sizes and vector dimensions. The scalar variables are signs over $\mathbb{R}$ and unit complex phases over $\mathbb{C}$. Our work gives the following bounds; neither exact constant is determined.

$$1375\pi/2454 \leq K_G^{\mathbb{R}} \leq \pi/[2(0.8818276493988)]$$
$$1.373 < K_G^{\mathbb{C}} \leq 100000000/71188883$$

Comparison with the literature

The literature columns contain publicly stated results and their direct consequences. “Before” includes the August 2026 real-case results [1]; “after” adds Guo, Fang and Lu [4]. The “Our work” column summarizes unpublished computer-assisted proofs.

Bound	Before the new paper	After the new paper	Our work
Real lower	$6\pi/11$ [1] ≈ 1.713595992867	$6\pi/11$ [1] ≈ 1.713595992867	$1375\pi/2454$ ≈ 1.760264832390
Real upper	1.781841324234^* [1]	1.781841324234^* [1]	1.781296297373
Complex lower	≈ 1.33807 [3] (Davie)	1.35584631827168 [4]	1.373
Complex upper	1.404909132736 [2] (Haagerup)	1.404909132736 [2] (Haagerup)	1.404713710707

Entries are endpoint values. Displayed decimal upper endpoints are rounded upward; values marked $\approx$ are approximations to the expressions shown. The exact inequalities for our work appear above. Bold entries indicate the strongest bounds in each row.

*Theorem 6.4 of [1] states the real upper bound 1.7818666069360661 . Its Proposition 6.3 and Lemma 6.2, combined with Proposition 6.1, give the stronger comparison endpoint used here:

$$\pi/\{2[0.881573822049 - 0.000011328860 - 14.443/\sqrt{(10 \cdot 251^5)}]\}$$
$$= 1.78184132423347648...$$

Size of the improvement. Relative to these real literature bounds, the lower endpoint increases by 0.04666884 and the upper endpoint decreases by 0.00054503 , reducing the interval width by **69.18%**. The complex lower endpoint exceeds [4] by **0.01715368**. Our complex interval is **52.55%** narrower than the Davie-Haagerup interval and **35.36%** narrower than the interval after [4]. These percentages compare interval widths, not relative errors in the unknown constants.

Other reported claims. Li et al. [5, §3, Table 2] report internally tested real bounds $51\pi/92 \approx 1.741535057968$ and 1.7813319810625639 , not independently verified by the authors and not stated as theorems. Our endpoints improve on these reported values by approximately 0.01872977 and 0.00003568 , respectively; the interval is **47.15%** narrower.

Methods and prior work

Real lower bound. The proof bounds the scalar norm of $(33/20)P_1 - P_3 - (3/20)P_5$, where P_j projects onto Hermite degree j . Full three-by-three transverse majorants, agreement estimates, and a circle inequality reduce the bound to universal one-dimensional ternary inequalities. Outward Gaussian integration and complete rectangle covers certify them in every dimension. Finite Gaussian partitions give explicit matrix witnesses, including one with ratio greater than 1.76; their certified ratio bounds tend to $1375\pi/2454$. The proof extends the agreement/disagreement analysis of [1]; the certified quintic scalar estimate is the substantive contribution.

Real upper bound. The construction extends Gaussian rounding and tensor preprocessing, including the coordinate-dependent framework of [1]. It uses a degree-nine polynomial threshold and a degree-eighty-one preprocessing polynomial. A lower bound for the linear correlation coefficient, minus the absolute sum of all nonlinear coefficients, gives the universal rounding guarantee. Interval enclosures cover coefficients through degree 501; a fourth-derivative bound controls the entire remaining tail. The contribution is a stronger explicit scheme and its complete coefficient estimate.

Complex lower bound. The operator is $P - \lambda I - \beta C_3$, with P the holomorphic linear projection and C_3 the full real cubic chaos, complexified. The Gaussian projection framework and proposed cubic perturbation precede both constructions [4, 6]. Our separate radial weights for real and imaginary components use the exact relation $\text{Re}(h \text{ conj}(k)) = 0$. Pointwise maximization and a finite family of auxiliary weights bound the whole remaining scalar parameter range for one fixed operator. Exact rational certificates and finite approximation yield the bound 1.373.

Relation to Guo-Fang-Lu. Their common radial-weight Gram/Schur criterion estimates the scalar norm within the Gaussian multiplier family shared by the two constructions. Theorem 5.2 of [4] confines that criterion's optimum to (1.35584631827168, 1.35584697425050); this is not an upper bound for the constant or for the entire operator family. Their §5.4 identifies discarded joint phase information. Our anisotropic estimate uses this information and lies outside that restricted criterion. Rotation-character compression places our construction in their multiplier family, so the sharper scalar estimate is the substantive distinction.

Complex upper bound. Complex Gaussian phase rounding combines signed tensor preprocessing with nonlinear phase perturbations, including a radial damping factor. For its radial correlation expansion, the linear coefficient minus the full absolute nonlinear coefficient mass bounds the retained objective. Complete contour enclosures establish zero winding numbers; a homotopy excludes the hypergeometric branch cut. Cauchy estimates then control every scalar tail. Exact bounds also cover omitted phase orders and primitive modes. This improves Haagerup's endpoint by approximately 0.000195422. The new lower-bound paper [4] does not change this upper-bound comparison.

References

- [1] R. Saha et al. *New Lower and Upper Bounds for the Grothendieck Constant*. [arXiv:2608.11158v2](#), §§2, 6-7 (2026).
- [2] U. Haagerup. *A new upper bound for the complex Grothendieck constant*. Israel J. Math. 60, 199-224 (1987). [doi:10.1007/BF02790792](#).
- [3] S. Friedland, L.-H. Lim and J. Zhang. *An elementary and unified proof of Grothendieck's inequality*. L'Enseignement Math. 64, 327-351 (2018), §1(v). [arXiv:1711.10595](#).
- [4] S. Guo, E. X. Fang and J. Lu. *An Improved Lower Bound for the Complex Grothendieck Constant*. [arXiv:2609.07000v1](#), Theorems 1.1, 3.1, 5.2 and §5.4 (2026).
- [5] A. Li et al. *Long-Horizon AI Research for Grothendieck Constant: A Case Study in Human-AI Mathematical Collaboration*. [arXiv:2608.11195v3](#), §3, Table 2 (2026).

[6] S. Heilman. *A lower bound for Grothendieck's constant*. [arXiv:2603.22616v1](#), §1.8 (2026).