WEEKLY METABOLIC RESEARCH LOG

Eloralintide Companion Tool | Structured tracking for clinician-guided metabolic therapy or formal research participation

WHY USE A LOG?	A single scale reading rarely explains what is happening. Tracking dose timing, symptoms, nutrition, hydration, bowel pattern and trend data helps separate three very different possibilities: **insufficient exposure, escalation that is poorly tolerated, or a nutrition/hydration problem.**

1 | WEEKLY SETUP

Week #		**Week of**		**Compound / therapy**	Eloralintide / _____
Exact weekly dose	_____ mg	**Administration date**	_____	**Time**	_____
Escalation this week?	Yes / No	**Previous dose**	_____ mg	**Escalation date**	_____
Dose held / delayed?	Yes / No	**If yes, why?**			

2 | 72-HOUR POST-ADMINISTRATION CHECK

Rate nausea from 0-10 and note other symptoms during the period when tolerability changes may be easiest to identify.

0-24 hours		Low / Typical / High		
24-48 hours		Low / Typical / High		
48-72 hours		Low / Typical / High		

3 | NUTRITION + HYDRATION

Daily Metric	Mon	Tue	Wed	Thu	Fri	Sat	Sun
Protein (g)							
Meals / day							
Fluids (oz)							
Appetite 0-10							

4 | GI + HYDRATION CHECK

Bowel pattern this week	Normal / Constipation / Diarrhea / Mixed	**Vomiting?**	No / Yes
Hydration concerns?	No / Mild / Moderate / Significant	**Abdominal symptoms?**	No / Yes
Notes			

WEEKLY METABOLIC RESEARCH LOG

Trend review + interpretation worksheet

5 | BODY-WEIGHT + WAIST TREND

Weight								

Waist this week	_____ in	**Prior week**	_____ in	**Change**	_____ in

6 | REASON FOR HOLDING OR REDUCING A DOSE

[ ] Persistent nausea	[ ] Poor oral intake	[ ] Illness
[ ] Dehydration	[ ] Procedure	[ ] Medication change
[ ] Clinician instruction	[ ] Other: _____	

7 | THREE-QUESTION TROUBLESHOOTING FRAMEWORK

1 | INSUFFICIENT EXPOSURE?	2 | ESCALATION / TOLERABILITY?	3 | NUTRITION / HYDRATION?
Is the current dose or duration simply too early to judge? Look at the trend rather than a single week.	Did symptoms increase after a dose change? Was intake disrupted for 48-72 hours after administration?	Are low protein intake, inadequate fluids, constipation, vomiting or very low food intake affecting the picture?
My notes:	My notes:	My notes:

8 | WEEKLY REVIEW

What changed this week?	
What appears to be working?	
What needs review with the clinician / study team?	
Plan for next week	

IMPORTANT	This worksheet is an educational tracking tool. It does not establish an eloralintide dosing schedule and should not be used to direct self-administration. Eloralintide is investigational and not FDA-approved. Clinical-trial dose arms are research designs, not prescribing instructions. Persistent or severe symptoms, inability to maintain hydration, or other concerning changes require appropriate medical evaluation.