

Part 1: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 2: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 3: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is

the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 4: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 5: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 6: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being

converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 7: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30

to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 8: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 9: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one

molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 10: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later

stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 11: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.

Part 12: Cellular Respiration Review

Cellular respiration is the set of metabolic reactions that cells use to convert the chemical energy stored in glucose into adenosine triphosphate, commonly called ATP. ATP is the main energy currency of the cell and powers processes such as muscle contraction, active transport and protein synthesis.

Overview of the Process

Aerobic respiration requires oxygen and occurs in three main stages: glycolysis, the Krebs cycle and the electron transport chain. The overall equation shows glucose and oxygen being converted into carbon dioxide, water and energy. A single molecule of glucose can yield up to 38 molecules of ATP under ideal conditions, although the realistic figure is closer to 30 to 32 ATP.

Stage 1: Glycolysis

Glycolysis takes place in the cytoplasm and does not require oxygen. During glycolysis one molecule of glucose is split into two molecules of pyruvate. This stage produces a net gain of two ATP molecules and two molecules of NADH, which carry high energy electrons to later stages.

Stage 2: The Krebs Cycle

The Krebs cycle, also known as the citric acid cycle, takes place in the mitochondrial

matrix. Before the cycle begins, pyruvate is converted into acetyl coenzyme A, releasing carbon dioxide. Each turn of the Krebs cycle produces one ATP, three NADH and one FADH₂, and releases two molecules of carbon dioxide as waste.

Stage 3: Electron Transport Chain

The electron transport chain is located in the inner mitochondrial membrane. NADH and FADH₂ donate electrons to a series of protein complexes, and the energy released pumps hydrogen ions across the membrane. The resulting gradient drives the enzyme ATP synthase, a process called oxidative phosphorylation, which produces the majority of ATP. Oxygen acts as the final electron acceptor and combines with hydrogen ions to form water.

Anaerobic Respiration

When oxygen is unavailable, cells rely on fermentation to regenerate NAD⁺ so that glycolysis can continue. In human muscle cells, pyruvate is converted into lactic acid, which contributes to muscle fatigue during intense exercise. In yeast, alcoholic fermentation converts pyruvate into ethanol and carbon dioxide. Fermentation produces only two ATP per glucose molecule, making it far less efficient than aerobic respiration.

Key Exam Points

Students should be able to name the location of each stage, state the inputs and outputs of glycolysis, explain why oxygen is essential as the final electron acceptor, and compare the ATP yield of aerobic respiration with fermentation.