

Simple Linear Model

Notes by Moutushi Chatterji L^AT_EXed by Sathvik H S

Model Assumptions and Model Objectives

Consider the general linear model of the form

$$Y = X\beta + \varepsilon \quad (1)$$

where

$$Y = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} \quad X = \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1p} \\ x_{21} & x_{22} & \cdots & x_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{np} \end{pmatrix} \quad \beta = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{pmatrix} \quad \varepsilon = \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{pmatrix}$$

Here Y is the vector of n observations, β is the vector of parameters, ε is the vector of random errors and X is the design matrix. Both X and β are fixed.

Assumptions

We assume ε to have the following properties:

$$\mathbb{E}(\varepsilon) = 0 \quad (2)$$

$$\text{Var}(\varepsilon) = \sigma^2 I_n \quad (3)$$

The second assumption implies that $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n$ have the same but unknown variances σ^2 and are uncorrelated.

Objectives

- i) To estimate the unknown parameters $\beta_1, \beta_2, \dots, \beta_p$ is possible OR at least estimate those linear combinations of β that can be estimated.
- ii) To test suitable statistical hypotheses about β OR at least functions of β .

Derivation

The rank of X is assumed to be r such that $r \leq \min(n, p)$. If $r = p \leq n$, the model (1) is said to be a *full rank model*, otherwise it is called a non full rank model.

Define $\hat{\beta}$ as a function of known quantities like Y, X , etc. such that $\hat{\beta}$ is “close” to β in some sense. Then,

1. $e = Y - X\hat{\beta}$ is the *vector of residuals*.
2. $\varepsilon = Y - X\beta$ is the *vector of errors*.

One way to obtain $\hat{\beta}$ is to use *method of least squares*.

Consider the sum of squares of residuals, i.e.,

$$\begin{aligned} e'e &= (Y - X\hat{\beta})'(Y - X\hat{\beta}) \\ &= Y'Y - Y'X\hat{\beta} - \hat{\beta}'X'Y - \hat{\beta}'X'X\hat{\beta} \\ &= Y'Y - 2\hat{\beta}'X'Y + \hat{\beta}'X'X\hat{\beta} \end{aligned}$$

The last equality follows since $Y'X\hat{\beta} = \hat{\beta}'X'Y$ as both sides are 1×1 matrices.

$$\implies \frac{\partial}{\partial \hat{\beta}}(e'e) = -2X'Y + 2(X'X)\hat{\beta}$$

Equating this to 0 we have,

$$X'Y = (X'X)\hat{\beta} \quad (4)$$

The above collection of equations are called *normal equations*.

When the inverse of $(X'X)$ exists, we simply have

$$\hat{\beta} = (X'X)^{-1}X'Y \quad (5)$$

Some Results

1. The matrix $S = X'X$ is a symmetric matrix having same rank as X .

Proof. If a vector α is orthogonal to the rows of the matrix X , then $X\alpha = 0 \implies X'X\alpha = 0$, i.e., α is orthogonal to the rows of $X'X$ as well.

Conversely, if $X'X\alpha = 0$, then $\alpha'X'X\alpha = 0$ or $Z'Z = 0$ where $Z = X\alpha$. But $Z'Z = z_1^2 + z_2^2 + \dots + z_n^2$ which implies $Z = X\alpha = 0$, i.e., α is orthogonal to the rows of X .

Thus X and $X'X$ has the same rank r . This also shows that the row space of X and row space of $S = X'X$ are the same. $\square$

2. If $\hat{\beta}$ be any particular solution to the (4), then it minimizes $e'e$, the sum of square of residuals.

Proof. Consider any other values of β , say β_0 . Then,

$$\begin{aligned} (Y - X\beta_0)'(Y - X\beta_0) &= (Y - X\hat{\beta} + X\hat{\beta} - X\beta_0)'(Y - X\hat{\beta} + X\hat{\beta} - X\beta_0) \\ &= (Y - X\hat{\beta})'(Y - X\hat{\beta}) + \left\{ X(\hat{\beta} - \beta_0) \right\}'(Y - X\hat{\beta}) \\ &\quad + (Y - X\hat{\beta})' \left\{ X(\hat{\beta} - \beta_0) \right\} + \left\{ X(\hat{\beta} - \beta_0) \right\}' \left\{ X(\hat{\beta} - \beta_0) \right\} \\ &= (Y - X\hat{\beta})'(Y - X\hat{\beta}) + (\hat{\beta} - \beta_0)'X'(Y - X\hat{\beta}) \\ &\quad + (Y - X\hat{\beta})'X(\hat{\beta} - \beta_0) + \left\{ X(\hat{\beta} - \beta_0) \right\}' \left\{ X(\hat{\beta} - \beta_0) \right\} \\ &= (Y - X\hat{\beta})'(Y - X\hat{\beta}) + (\hat{\beta} - \beta_0)'(X'Y - X'X\hat{\beta}) \\ &\quad + (X'Y - X'X\hat{\beta})'(\hat{\beta} - \beta_0) + m'm \quad \text{where } m = X(\hat{\beta} - \beta_0) \\ &= (Y - X\hat{\beta})'(Y - X\hat{\beta}) + m'm \quad \text{(from (4))} \\ &\geq (Y - X\hat{\beta})'(Y - X\hat{\beta}) \quad \square \end{aligned}$$

Thus the minimized value can be unambiguously denoted by $SSE = (Y - X\hat{\beta})'(Y - X\hat{\beta})$ and stands for *sum of squares due to error*.

Generalized Inverse and It's Properties

Definition

1. Any $n \times m$ matrix A^- satisfying the relation $AA^-A = A$ is defined as a *generalized inverse* (*g-inverse*) of the $m \times n$ matrix A .
2. An $n \times m$ matrix A^- is defined to be a *generalized inverse* of the $m \times n$ matrix A if for every u satisfying $\text{rank}(A) = \text{rank}(A|u)$, A^-u is a solution of the equations $Ax = u$ for unknown u . In other words, A^-u is a solution of the equations $Ax = u$ for unknown u whenever the system is *consistent*.

Properties

Define an $n \times n$ matrix $H = A^-A$. Then,

1. $AH = A$
2. $H^2 = H$
3. $\text{Tr}(H) = \text{rank}(H) = \text{rank}(A)$

Proof. The first two properties follow directly from definition.

The first equality in property 3 follows as H is an idempotent matrix (property 2) and is similar to a diagonal matrix containing eigenvalues of H (spectral theorem). But the only eigenvalues of an idempotent matrix are 0 and 1 as the polynomial $x^2 - x$ annihilates H .

For the second equality, observe that $A^-A\alpha = 0 \implies AA^-A\alpha = 0 \implies A\alpha = 0$ and $A\alpha = 0 \implies A^-A\alpha = 0$. i.e, if a vector α is orthogonal to the rows of A , then it is orthogonal to the rows of H and vice versa. Hence $\text{rank } A = \text{rank } H$. $\square$

General Solution for a System of Homogenous Equations

Observe that, $A(I - H) = A - AH = 0$, by the first property of g-inverse. i.e, each of the n columns $h_1, h_2, \dots, h_n$ of $I - H$ are orthogonal to the rows of A .

Also, $(I - H)^2 = I - H - H + H^2 = I - H$ by second property of g-inverse. i.e, $I - H$ is idempotent.¹ This implies $\text{rank}(I - H) = \text{Tr}(I - H) = \text{Tr}(I) - \text{Tr}(H) = n - r$, where $r = \text{rank}(A) = \text{rank}(H)$ from third property of g-inverse.

Thus, only $(n - r)$ of the columns vectors $h_1, h_2, \dots, h_n$ are linearly independent. Without loss of generality we assume that the first $(n - r)$ columns of $I - H$, $h_1, h_2, \dots, h_{n-r}$ are linearly independent.

Since A is an $m \times n$ matrix of rank r , its rows are n -vectors and therefore we can find at most $(n - r)$ linearly independent vectors orthogonal to them. $h_1, h_2, \dots, h_{n-r}$ is one such set. If there exists any other vector orthogonal to the rows of A , it must be a linear combination of $h_1, h_2, \dots, h_{n-r}$.

Let us now consider a system of homogenous equations

$$Ax = 0 \tag{6}$$

By definition, x is orthogonal to the rows of A and so any vector satisfying (6) must be a linear combination of $h_1, h_2, \dots, h_{n-r}$. This is equivalent to saying that x will be a linear combination of columns of $I - H$, $h_1, h_2, \dots, h_n$, because $h_{n-r+1}, h_{n-r+2}, \dots, h_n$ are linear combinations of $h_1, h_2, \dots, h_{n-r}$ themselves.

¹Let R be a ring with unity 1_R and e be an idempotent element of this ring. Then, $1_R - e$ is also idempotent.

Therefore, x must be of the form

$$x = z_1 h_1 + z_2 h_2 + \cdots + z_n h_n = (h_1 \ h_2 \ \dots \ h_n)z = (I - H)z \quad (7)$$

for some $z = (z_1 \ z_2 \ \dots \ z_n)'$.²

Conversely, if (7) holds, $Ax = A(I - H)z = (A - AH)z = 0$, by the first property of g-inverse. This shows that the general solution of (6) is given by $\tilde{x} = (I - H)z$.

General Solution for a System of Non-Homogenous Equations

Consider a consistent system of non-homogenous equations given by

$$Ax = u \quad (8)$$

If A^- is any g-inverse of A , by definition of A^- , A^-u is a particular solution to (8). Thus $A(x - A^-u) = u - u = 0$ is a system of homogenous equations in $x - A^-u$. The corresponding general solution of this homogenous system is $\tilde{x} - A^-u = (I - H)z$ which implies, the general solution to (8) is,

$$\tilde{x} = A^-u + (I - H)z \quad (9)$$

Solution of the Normals Equations (General Form)

The normal equations are given by (as seen from (4)),

$$X'Y = (X'X)\hat{\beta}$$

A particular solution of these equations will be

$$\hat{\beta} = S^-X'Y$$

where S^- is any g-inverse of $S = X'X$. Thus, the general solution of normal equations will be,

$$\tilde{\beta} = \hat{\beta} + (I - H)z \quad (10)$$

where $H = S^-S$ is a $p \times p$ matrix.

Some More Results

3. $X = XH$

Proof. From the first property of g-inverse, we have $SH = S$. Therefore,

$$\begin{aligned} 0 &= (I - H)'(S - SH) \\ &= (I - H)'(X'X - X'XH) \\ &= (I - H)'(X'(X - XH)) \\ &= (X(I - H))'(X - XH) \\ &= (X - XH)'(X - XH) \end{aligned}$$

Note that both the sides of the equality are $p \times p$ matrices. Thus equating the i th diagonal elements of both sides, we get that the sum of squares of the entries in i th row of $(X - XH)$ is zero for every $i = 1, 2, \dots, p$. Thus every element of $(X - XH)$ is zero. $\square$

²These z_i 's need not be unique even for a fixed x as columns of $I - H$ are not necessarily linearly independent.

4. A solution of the normal equations (4) is unique iff $\text{rank}(X) = \text{rank}(X'X) = p$.

Proof. A solution to the normal equations is unique iff the general solution (10) of normal equations doesn't contain the arbitrary vector.

This happens iff $I - H = 0$ i.e, $I = H = S^-S$ which happens iff S is non-singular and has regular inverse S^{-1} . Thus the result follows. $\square$