$success: 1 \rightarrow p$ $failure: 0 \rightarrow q$ p+q=1

$$\mu_1 = 1 \cdot p = p$$

$$\sigma_1^2 = (1 - \mu_1)^2 p + (0 - \mu_1)^2 q = (1 - p)^2 p + (0 - p)^2 (1 - p)$$

$$= (1 - p)(p - p^2 + p^2) = p(1 - p) = pq$$

 $\therefore \mu = np$

 $\therefore \sigma^2 = npq$

(2)

$$f(x) = \binom{n}{x} p^x q^{n-x}$$
 $@x = m: f(x) \to maxima$

 $@ x = m : ln f(x) \rightarrow maxima$

$$f = f(x)$$
 $f' = f'(x)$ $f'(m) = [f'(x)]_{x=m}$
 $f'(m) = 0$ $(\ln f)'(m) = 0$

$$\ln f = \ln f(m) + (\ln f)'(m) (x - m) + \frac{1}{2} (\ln f)''(m) (x - m)^2 + \sum_{k=3}^{\infty} \frac{1}{k!} (\ln f)^{(k)} (m) (x - m)^k$$

$$f = f(m) e^{\frac{1}{2} (\ln f)''(m)(x - m)^2} e^{\sum_{k=3}^{\infty} \frac{1}{k!} (\ln f)^{(k)} (m)(x - m)^k}$$

(3) $\ln x! = \sum_{k=2}^{x} \ln k \approx \int_{1}^{x} \ln t \, dt \approx x \ln x - x$

$$\ln f = \ln \left(\frac{n!}{(n-x)!x!} p^x q^{n-x} \right) = \ln n! - \ln(n-x)! - \ln x! + \ln p^x + \ln q^{n-x}$$

$$\approx n \ln n - n - (n-x) \ln(n-x) + (n-x) - x \ln x + x + x \ln p + (n-x) \ln q$$

$$= n \ln n - n - n \ln(n-x) + x \ln(n-x) + n - x - x \ln x + x + x \ln p + n \ln q - x \ln q$$

$$(\ln f)' = -n \frac{-1}{n-x} + \ln(n-x) + x \frac{-1}{n-x} - 1 - (\ln x + 1) + 1 + \ln p - \ln q$$

$$= \frac{n}{n-x} + \ln(n-x) - \frac{x}{n-x} - 1 - \ln x - 1 + 1 + \ln p - \ln q$$

$$= 1 + \ln(n-x) - 1 - \ln x - 1 + 1 + \ln p - \ln q = \ln(n-x) - \ln x + \ln p - \ln q = \ln \left(\frac{(n-x)p}{xq} \right)$$

$$(\ln f)'(m) = 0$$
 $\ln \left(\frac{(n-m)p}{mq}\right) = 0$ $\frac{(n-m)p}{mq} = 1$

$$(n-m)p = mq$$
 $np - mp = mq$ $np = m(p+q)$ $\therefore m = np = \mu$

$$(\ln f)'' = (\ln(n-x) - \ln x + \ln p - \ln q)' = -\frac{1}{n-x} - \frac{1}{x}$$

$$(\ln f)''(m) = -\frac{1}{n-m} - \frac{1}{m} = -\frac{1}{n-np} - \frac{1}{np} = -\frac{1}{nq} - \frac{1}{np} = \frac{-p-q}{npq} = -\frac{1}{npq} = -\frac{1}{\sigma^2}$$

$$(\ln f)^{(k)} = (k-2)! \left(-(n-x)^{-(k-1)} + (-1)^{k-1} x^{-(k-1)} \right)$$

$$(\ln f)^{(k)}(m) = (k-2)! \left(-(n-np)^{-(k-1)} + (-1)^{k-1} (np)^{-(k-1)} \right) = (k-2)! \cdot \frac{-p^{k-1} + (-q)^{k-1}}{(npq)^{k-1}}$$

$$\lim_{n \to \infty} \frac{1}{k!} (\ln f)^{(k)}(m) = \lim_{n \to \infty} \left(\frac{1}{k(k-1)} \cdot \frac{-p^{k-1} + (-q)^{k-1}}{n^{k-1} (pq)^{k-1}} \right) = 0$$

(4)

$$f \approx f(m) e^{\frac{1}{2}(\ln f)''(m)(x-m)^2} = f(\mu) e^{\frac{1}{2}(-\frac{1}{\sigma^2})(x-\mu)^2} = f(\mu) e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$

$$\therefore \binom{n}{x} p^x (1-p)^{n-x} \approx \binom{n}{\mu} p^\mu (1-p)^{n-\mu} e^{-\frac{1}{2}(\frac{x-\mu}{\sigma})^2}$$