

Solution no – 01

```
KD1-86660-pratik@>select onum,cname from orders,customers where customers.cnum=orders.cnum;
```

onum	cname
3001	Cisneros
3003	Hoffman
3002	Pereira
3005	Liu
3006	Cisneros
3009	Giovanni
3007	Grass
3008	Clemens
3010	Grass
3011	Clemens

10 rows in set (0.08 sec)

Solution no – 02

```
KD1-86660-pratik@>select onum,cname,sname from orders,customers,salespeople where orders.cnum=customers.cnum and orders.snum=salespeople.snum;
```

onum	cname	sname
3003	Hoffman	Peel
3009	Giovanni	Axelrod
3005	Liu	Serres
3010	Grass	Serres
3007	Grass	Serres
3011	Clemens	Peel
3008	Clemens	Peel
3006	Cisneros	Rifkin
3001	Cisneros	Rifkin
3002	Pereira	Motika

10 rows in set (0.00 sec)

Solution no – 03

```
KD1-86660-pratik@>select cname,sname,comm from customers,salespeople where salespeople.snum=customers.snum and comm>0.12
;
```

cname	sname	comm
Liu	Serres	0.13
Grass	Serres	0.13
Cisneros	Rifkin	0.15

```
3 rows in set (0.00 sec)
```

Solution no – 04

```
KD1-86660-pratik@>select distinct onum,rating,comm*amt as comm from orders,customers,salespeople where salespeople.snum
=orders.snum and (orders.cnum=customers.cnum and customers.rating > 100);
```

onum	rating	comm
3010	300	40.2935
3007	300	9.8475
3005	200	670.8585
3006	300	164.7240
3001	300	2.8035
3009	200	171.3230

```
6 rows in set (0.00 sec)
```