

Solution no -01

```
KD1-86660-pratik@>select onum,snum,amt*0.12 as amt from orders;
```

onum	snum	amt
3001	1007	2.2428
3003	1001	92.0628
3002	1004	228.0120
3005	1002	619.2540
3006	1007	131.7792
3009	1003	205.5876
3007	1002	9.0900
3008	1001	566.7600
3010	1002	37.1940
3011	1001	1187.0256

```
10 rows in set (0.00 sec)
```

Solution no – 02

```
KD1-86660-pratik@>select distinct city,max(rating) as rating from customers group by city;
```

city	rating
London	100
Rome	200
San Jose	300
Berlin	300

```
4 rows in set (0.00 sec)
```

Solution no – 03

```
KD1-86660-pratik@>select rating,cname,cnum from customers order by rating desc;
```

rating	cname	cnum
300	Grass	2004
300	Cisneros	2008
200	Giovanni	2002
200	Liu	2003
100	Hoffman	2001
100	Clemens	2006
100	Pereira	2007

```
7 rows in set (0.00 sec)
```

Solution no – 04

```
KD1-86660-pratik@>select distinct odate,count(distinct onum) as count from orders group by odate order by count desc;
```

odate	count
1990-10-03	5
1990-10-04	2
1990-10-06	2
1990-10-05	1

```
4 rows in set (0.00 sec)
```