

Contents

1 08-26-2025

- 1.1 Basics
- 1.2 Planes and Spheres
- 1.3 Vector Notation
- 1.4 Computing 3x3 Determinants
 - 1.4.1 Cofactor Method
 - 1.4.2 Diagonal Method

2 08-28-2025

- 2.1 Section 2.3, Checkpoint 2.23: Finding the Angle between Two Vectors
- 2.2 Section 2.3, Checkpoint 2.24: Identifying Orthogonal Vectors
- 2.3 Section 2.3, Checkpoint 2.27: Resolving Vectors into Components
- 2.4 Section 2.4, Checkpoint 2.33 (quick)
- 2.5 Section 2.4, Checkpoint 2.38
- 2.6 Section 2.4, Example 2.44: Evaluating Torque

3 09-02-2025

- 3.1 Section 2.5, Checkpoint 2.45
- 3.2 Section 2.5, Checkpoint 2.46
- 3.3 Section 2.5, Checkpoint 2.47
- 3.4 Section 2.5, Checkpoint 2.48

4 09-04-2025

- 4.1 Section 2.6, Exercises 309-312
- 4.2 Section 2.6, Example 2.59 Identifying Equations of Quadric Surfaces
- 4.3 Section 2.6, Checkpoint 2.54
- 4.4 Section 2.6, Exercise 350

5 09-09-2025

- 5.1 Section 3.1, Exercise 8
- 5.2 Section 3.1, Exercise 22

5.3	Section 3.2, Checkpoint 3.5
5.4	Section 3.2, Checkpoint 3.7
5.5	Section 3.2, Checkpoint 3.8
6	09-11-2025	
6.1	Section 3.3, Checkpoint 3.9
6.2	Section 3.3, Checkpoint 3.13
6.3	Section 3.3, Exercise 3.143
7	09-16-2025 - Midterm 1 tomorrow	
8	09-18-2025 - Post Midterm 1, No Content	
9	09-23-2025	
9.1	Section 4.1, Checkpoint 4.4
9.2	Section 4.1, Checkpoint 4.5
9.3	Section 4.2, Exercise 60
9.4	Section 4.2, Exercise 77
9.5	Section 4.2, Exercise 81
9.6	Section 4.2, Exercise 86
9.7	Section 4.2, Exercise 88
9.8	Section 4.2, Exercise 95
10	09-25-2025	
10.1	Section 4.3, Checkpoint 4.13
10.2	Section 4.3, Exercises 114-117
10.3	Section 4.4, Checkpoint 4.19
10.4	Section 4.4, Checkpoint 4.22
11	09-30-2025	
11.1	Section 4.5, Checkpoint 4.23
11.2	Section 4.5, Exercise 253
11.3	Section 4.5, Example 4.30
11.4	Section 4.6, Example 4.31
11.5	Section 4.6, Example 4.32
11.6	Section 4.6, Checkpoint 4.30

12 10-02-2025

- 12.1 Section 4.7, Checkpoint 4.35
- 12.2 Section 4.7, Checkpoint 4.36

13 10-07-2025 - Midterm 2 tomorrow

- 13.1 Section 4.8, Exercise 369
- 13.2 Section 4.8, Exercise 373
- 13.3 Section 4.8, Exercise 374
- 13.4 Section 4.8, Exercise 379
- 13.5 Section 4.8, Exercise 388

14 10-09-2025 - Post Midterm 2, No Content

15 10-21-2025

16 10-23-2025

17 10-28-2025

18 10-30-2025

19 11-06-2025

20 11-11-2025 - Midterm 3 tomorrow

21 11-13-2025 - Post Midterm 3, No Content

22 11-18-2025

23 11-20-2025

24 11-25-2025

25 12-02-2025 - Final Review

26 12-04-2025 - Final Review

1 08-26-2025

1.1 Basics

This class is **Calculus in Three Dimensions**, thus the class will require you to think in 3-D.

- Get used to drawing in 3-D.

My preference is to draw the usual vertical and horizontal axes.

Label the horizontal axis y and the vertical axis z .

Then draw a diagonal from the bottom left to the top right, passing through the origin. Label this axis x . Note that you can interchange any of the axis labels as necessary (which will come in handy when we start doing 3-D integration).

- In \mathbb{R}^3 , the (Euclidean) distance (also referred to as the L_2 norm) between any two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

1.2 Planes and Spheres

In this course, you will often deal with planes and spheres (e.g., spherical coordinates).

- You will learn the canonical formula for a plane later (finding it involves cross products). For now, we will find equations for planes parallel to another plane.

Ex. Write an equation of the plane passing through point $(21, 2, 59)$ that is parallel to the xz -plane.

When a plane is parallel to the xz -plane, it means only the x and z coordinates may vary.

Thus taking the y -value, we get the equation $y = 2$.

Ex. Write an equation of the plane passing through points $(2, 125, 9)$, $(21, 25, 9)$, $(5, 7, 9)$ that is parallel to the xy -plane.

When a plane is parallel to the xy -plane, it means only the x and y coordinates may vary.

Conveniently, the z -value in all 3 coordinates are equal; we get the equation $z = 9$.

- A sphere is the set of all points in space equidistant from a fixed point, the center of the sphere. For center (a, b, c) and radius r , we represent the sphere by the (“canonical”) equation:

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$$

Ex. Find the equation of the sphere with diameter \overline{PQ} where $P = (2, -1, -3)$ and $Q = (-2, 5, -1)$.

First, find the center, which is at the midpoint of the diameter \overline{PQ} :

$$C = \left(\frac{2 + (-2)}{2}, \frac{-1 + 5}{2}, \frac{-3 + (-1)}{2} \right) = (0, 2, -2)$$

Next, find the radius using the distance formula (half the length of the diameter)

$$r = \frac{1}{2} \|\overline{PQ}\| = \frac{1}{2} \sqrt{(-2 - 2)^2 + (5 - (-1))^2 + (-1 - (-3))^2} = \frac{1}{2} \sqrt{56} = \sqrt{\frac{56}{4}} = \sqrt{14}$$

Thus the sphere is given by $x^2 + (y - 2)^2 + (z + 2)^2 = 14$

1.3 Vector Notation

Vectors are quantities with magnitude and direction. In \mathbb{R}^3 , the standard unit vectors are

$$\hat{i} = \langle 1, 0, 0 \rangle \quad \hat{j} = \langle 0, 1, 0 \rangle \quad \hat{k} = \langle 0, 0, 1 \rangle$$

There are several notations. Fix points in \mathbb{R}^3 $P = (0, 2, 1)$ and $Q = (2, 5, 9)$

Let's represent the vector $\overrightarrow{PQ} = \langle x_Q - x_P, y_Q - y_P, z_Q - z_P \rangle$.

- Component Form: $\overrightarrow{PQ} = \langle 2 - 0, 5 - 2, 9 - 1 \rangle = \langle 2, 3, 8 \rangle$
- Using the unit vectors: $\overrightarrow{PQ} = 2\hat{i} + 3\hat{j} + 8\hat{k}$

Ex: Consider the vectors $v = \langle 0, 2, 1 \rangle$ and $u = \langle 2, 5, 9 \rangle$. Find a unit vector in the direction of $3v + u$.

You will need the following rules (taken from your textbook)

RULE: PROPERTIES OF VECTORS IN SPACE

Let $\mathbf{v} = \langle x_1, y_1, z_1 \rangle$ and $\mathbf{w} = \langle x_2, y_2, z_2 \rangle$ be vectors, and let k be a scalar.

Scalar multiplication: $k\mathbf{v} = \langle kx_1, ky_1, kz_1 \rangle$

Vector addition: $\mathbf{v} + \mathbf{w} = \langle x_1, y_1, z_1 \rangle + \langle x_2, y_2, z_2 \rangle = \langle x_1 + x_2, y_1 + y_2, z_1 + z_2 \rangle$

Vector subtraction: $\mathbf{v} - \mathbf{w} = \langle x_1, y_1, z_1 \rangle - \langle x_2, y_2, z_2 \rangle = \langle x_1 - x_2, y_1 - y_2, z_1 - z_2 \rangle$

Vector magnitude: $\|\mathbf{v}\| = \sqrt{x_1^2 + y_1^2 + z_1^2}$

Unit vector in the direction of \mathbf{v} : $\frac{1}{\|\mathbf{v}\|}\mathbf{v} = \frac{1}{\|\mathbf{v}\|} \langle x_1, y_1, z_1 \rangle = \left\langle \frac{x_1}{\|\mathbf{v}\|}, \frac{y_1}{\|\mathbf{v}\|}, \frac{z_1}{\|\mathbf{v}\|} \right\rangle$, if $\mathbf{v} \neq \mathbf{0}$

Solution: first find $3v + u$, then find unit vector in that direction.

$$3v + u = \langle 3(0), 3(2), 3(1) \rangle + \langle 2, 5, 9 \rangle \quad (\text{scalar multiplication})$$

$$= \langle 0, 6, 3 \rangle + \langle 2, 5, 9 \rangle = \langle 2, 11, 12 \rangle$$

$$\frac{1}{\|3v + u\|}(3v + u) = \frac{\langle 2, 11, 12 \rangle}{\sqrt{2^2 + 11^2 + 12^2}} = \left\langle \frac{2}{\sqrt{269}}, \frac{11}{\sqrt{269}}, \frac{12}{\sqrt{269}} \right\rangle \quad (\text{unit vector in direction})$$

Observe that 269 is prime, so you can't simplify the denominator. Thus we're done. Not all numbers will be pretty, but as a tip: make sure your answers are feasible. If you are taking the length of a vector with 1-digit coordinates and get a 5-digit number, that's probably wrong.

DON'T FORGET THE SQUARE ROOT $\sqrt{\quad}$ when taking norms!!!!

1.4 Computing 3x3 Determinants

Useful for computing cross products. You are free to use whatever trick you want so long as you show your work. (note vertical bars means determinant, while square brackets mean matrix).

1.4.1 Cofactor Method

Key notes: $+$, $-$, $+$ (**DON'T FORGET THE MIDDLE IS NEGATIVE**) and the 2x2 sub-determinants are the elements not in the same row or column.

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix} = a(ei - fh) - b(di - fg) + c(dh - eg) \\ = aei - afg - bdi + bfg + cdh - ceg$$

For cross products between two vectors $u = \langle u_x, u_y, u_z \rangle$ and $v = \langle v_x, v_y, v_z \rangle$, you just compute the 3x3

determinant $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ u_x & u_y & u_z \\ v_x & v_y & v_z \end{vmatrix}$

1.4.2 Diagonal Method

Recall that the 2x2 determinant is computed using the diagonals: $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$, where the diagonals going from right to left are positive and diagonals going from left to right are negative. We extend this idea to 3x3 determinants:

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} \xrightarrow{\text{expand}} \begin{matrix} & a & b & c \\ f & d & e & f & d \\ h & i & g & h & i & g & h \end{matrix} \xrightarrow{\text{compute}} aei + bfg + cdh - afh - bdi - ceg$$

which is the same result as derived from the cofactor method.

2 08-28-2025

2.1 Section 2.3, Checkpoint 2.23: Finding the Angle between Two Vectors

Find the measure of the angle, in radians, formed by vectors $a = \langle 1, 2, 0 \rangle$ and $b = \langle 2, 4, 1 \rangle$. Round to the nearest hundredth.

2.2 Section 2.3, Checkpoint 2.24: Identifying Orthogonal Vectors

For which value of x is $p = \langle 2, 8, -1 \rangle$ orthogonal to $q = \langle x, -1, 2 \rangle$?

2.3 Section 2.3, Checkpoint 2.27: Resolving Vectors into Components

Express $v = 5i - j$ as a sum of orthogonal vectors such that one of the vectors has the same direction as $u = 4i + 2j$.

Properties of the Cross Product

Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in space, and let c be a scalar.

- | | | |
|------|---|---------------------------------------|
| i. | $\mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u})$ | Anticommutative property |
| ii. | $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = \mathbf{u} \times \mathbf{v} + \mathbf{u} \times \mathbf{w}$ | Distributive property |
| iii. | $c(\mathbf{u} \times \mathbf{v}) = (c\mathbf{u}) \times \mathbf{v} = \mathbf{u} \times (c\mathbf{v})$ | Multiplication by a constant |
| iv. | $\mathbf{u} \times \mathbf{0} = \mathbf{0} \times \mathbf{u} = \mathbf{0}$ | Cross product of the zero vector |
| v. | $\mathbf{v} \times \mathbf{v} = \mathbf{0}$ | Cross product of a vector with itself |
| vi. | $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$ | Scalar triple product |

2.4 Section 2.4, Checkpoint 2.33 (quick)

Use the properties of the cross product to calculate $(i \times k) \times (k \times j)$.

2.5 Section 2.4, Checkpoint 2.38

Find the area of the parallelogram $PQRS$ with vertices $P(1, 1, 0)$, $Q(7, 1, 0)$, $R(9, 4, 2)$, and $S(3, 4, 2)$.

2.6 Section 2.4, Example 2.44: Evaluating Torque

A bolt is tightened by applying a force of 6 N to a 0.15-m wrench (Figure 2.62). The angle between the wrench and the force vector is 40° . Find the magnitude of the torque about the center of the bolt. Round the answer to two decimal places.




Figure 2.62 Torque describes the twisting action of the wrench.

3 09-02-2025

3.1 Section 2.5, Checkpoint 2.45

Find the distance between point $(0, 3, 6)$ and the line with parametric equations

$$x = 1 - t, y = 1 + 2t, z = 5 + 3t$$

3.2 Section 2.5, Checkpoint 2.46

Describe the relationship between the lines with the following parametric equations:

$$x = 1 - 4t, y = 3 + t, z = 8 - 6t$$

$$x = 2 + 3s, y = 2s, z = -1 - 3s$$

3.3 Section 2.5, Checkpoint 2.47

Find an equation of the plane containing the lines L_1 and L_2 :

$$L_1 : x = -y = z$$

$$L_2 : \frac{x-3}{2} = y = z - 2$$

On the original handout, I mistyped and had $L_2 = x - 32$. This has been corrected, as the error makes the question unsolvable since the lines would be skew.

Worked Solution

Let $L_1 : x = -y = z = t$. The parameterized form of L_1 is thus $\langle t, -t, t \rangle = (0, 0, 0) + \langle 1, -1, 1 \rangle t$

Let $L_2 : \frac{x-3}{2} = y = z - 2 = t$. The parameterized form of L_2 is thus $\langle 2t + 3, t, t + 2 \rangle = (3, 0, 2) + \langle 2, 1, 1 \rangle t$.

A normal vector to the plane that contains both lines is the cross product of the two lines' direction vectors, which was $\vec{n} = \langle -2, 1, 3 \rangle$ (*work done in recitation*).

Recall the equation of a plane is $\vec{n} \cdot (\vec{r} - \vec{r}_0) = 0$, where $\vec{r} = (x, y, z)$, \vec{n} is a normal vector of the plane, and \vec{r}_0 is some arbitrary point in the plane. *Note I said a normal vector, since we could scale \vec{n} arbitrarily.*

We have \vec{n} , so we just need to get \vec{r}_0 . *In recitation, I mentioned that we can pick any arbitrary point on either line to create the equation of the plane that contains L_1 and L_2 . This holds true, and I will arbitrarily pick some points below and prove that the resulting plane equation is the same.*

- In recitation, we picked the easy point $(0, 0, 0)$, which we know to lie in L_1 .

$$\begin{aligned}\vec{n} \cdot (\vec{r} - \vec{r}_0) &= \langle -2, 1, 3 \rangle \cdot (x - 0, y - 0, z - 0) \\ &= -2x + y + 3z = 0\end{aligned}$$

- But what if we had picked $(3, 0, 2)$, which was our anchor point for L_2 ?

$$\begin{aligned}\vec{n} \cdot (\vec{r} - \vec{r}_0) &= \langle -2, 1, 3 \rangle \cdot (x - 3, y - 0, z - 2) \\ &= -2(x - 3) + y + 3(z - 2) = 0 \\ &= -2x + 6 + y + 3z - 6 = 0 \\ &= -2x + y + 3z = 0\end{aligned}\quad \text{(the 6's cancel!)}$$

- As an extension, derive the plane equation you get if you choose the intersection point of L_1 and L_2 ! You'll see that you still reach the same plane equation.

3.4 Section 2.5, Checkpoint 2.48


Find the distance between point $P = (5, -1, 0)$ and the plane given by $4x + 2y - z = 3$.

Note: We do NOT cover the foci of paraboloids or ellipsoids.


4.1 Section 2.6, Exercises 309-312

For the following exercises, the graph of a quadric surface is given.


- Specify the name of the quadric surface.
- Determine the axis of the quadric surface.




309.



310.



311.



312.

4.2 Section 2.6, Example 2.59 Identifying Equations of Quadric Surfaces

Identify the surfaces represented by the given equations.

- (a) $16x^2 + 9y^2 + 16z^2 = 144$ **Solution** Observe that all the terms are squared, and all the coefficients are positive (but not equal) so this is an ellipsoid.
- (b) $9x^2 - 18x + 4y^2 + 16y - 36z + 25 = 0$ **Solution** We have two squared terms, and z term of degree one. We could rearrange this to get $36z = 9x^2 - 18x + 4y^2 + 16y + 25$, which appears to be a elliptic paraboloid.

4.3 Section 2.6, Checkpoint 2.54

Identify the surface represented by the equation $9x^2 + y^2 - z^2 + 2z - 10 = 0$.

Solution We have three squared terms, two positive coefficients and one negative coefficient. So this is a double cone.

4.4 Section 2.6, Exercise 350

Find the equation of the quadric surface with points $P(x, y, z)$ that are equidistant from point $Q(0, 2, 0)$ and plane of equation $y = -2$. Identify the surface.

Solution First, the distance from a point (x, y, z) to the plane $y = -2$ is just $|y|$. *This is because the plane $y = -2$ has normal vector \hat{j} ; projecting \overrightarrow{PQ} onto \hat{j} just yields the y component.*

To do this, we want $\langle x, y, z \rangle$ that satisfies

$$\begin{aligned}\sqrt{(x-0)^2 + (y-2)^2 + (z-0)^2} &= y - (-2) \\ \sqrt{x^2 + (y-2)^2 + z^2} &= y + 2 \\ x^2 + (y-2)^2 + z^2 &= (y+2)^2 \\ x^2 + y^2 - 4y + 4 + z^2 &= y^2 + 4y + 4 \\ x^2 + z^2 &= 8y\end{aligned}$$

This is the form of an elliptic paraboloid.

5 09-09-2025

5.1 Section 3.1, Exercise 8

Find the limit of the following vector valued function at the indicated value of t .

$$\lim_{t \rightarrow 4} \left\langle \sqrt{t-3}, \frac{\sqrt{t}-2}{t-4}, \tan\left(\frac{\pi}{t}\right) \right\rangle$$

Solution We can just substitute for the x and z components. The middle one requires more thought; direct substitution gives $\frac{0}{0}$, so we apply L'Hopital's rule.

$$\lim_{t \rightarrow 4} \frac{\sqrt{t}-2}{t-4} = \lim_{t \rightarrow 4} \frac{\frac{1}{2\sqrt{t}}}{1} = \frac{1}{4}$$

So the limit is

$$\left\langle 1, \frac{1}{4}, 1 \right\rangle$$

5.2 Section 3.1, Exercise 22

Eliminate the parameter t , write the equation in Cartesian coordinates, then sketch the graphs of the vector-valued functions. *Hint: solve first equation for x in terms of t and substitute this result into the second equation.*

$$\mathbf{r}(t) = 2t\hat{i} + t^2\hat{j} \quad \text{let } x = 2t, y = t^2$$

THEOREM 3.3

Properties of the Derivative of Vector-Valued Functions

Let \mathbf{r} and \mathbf{u} be differentiable vector-valued functions of t , let f be a differentiable real-valued function of t , and let c be a scalar.

- | | | |
|------|--|--------------------|
| i. | $\frac{d}{dt}[c\mathbf{r}(t)] = c\mathbf{r}'(t)$ | Scalar multiple |
| ii. | $\frac{d}{dt}[\mathbf{r}(t) \pm \mathbf{u}(t)] = \mathbf{r}'(t) \pm \mathbf{u}'(t)$ | Sum and difference |
| iii. | $\frac{d}{dt}[f(t)\mathbf{u}(t)] = f'(t)\mathbf{u}(t) + f(t)\mathbf{u}'(t)$ | Scalar product |
| iv. | $\frac{d}{dt}[\mathbf{r}(t) \cdot \mathbf{u}(t)] = \mathbf{r}'(t) \cdot \mathbf{u}(t) + \mathbf{r}(t) \cdot \mathbf{u}'(t)$ | Dot product |
| v. | $\frac{d}{dt}[\mathbf{r}(t) \times \mathbf{u}(t)] = \mathbf{r}'(t) \times \mathbf{u}(t) + \mathbf{r}(t) \times \mathbf{u}'(t)$ | Cross product |
| vi. | $\frac{d}{dt}[\mathbf{r}(f(t))] = \mathbf{r}'(f(t)) \cdot f'(t)$ | Chain rule |
| vii. | If $\mathbf{r}(t) \cdot \mathbf{r}(t) = c$, then $\mathbf{r}(t) \cdot \mathbf{r}'(t) = 0$. | |

5.3 Section 3.2, Checkpoint 3.5

Calculate the derivative of the function

$$\mathbf{r}(t) = (t \ln t)\hat{i} + (5e^t)\hat{j} + (\cos t - \sin t)\hat{k}$$

5.4 Section 3.2, Checkpoint 3.7

Find the unit tangent vector for the vector-valued functions

$$\mathbf{r}(t) = (t^2 - 3)\hat{i} + (2t + 1)\hat{j} + (t - 2)\hat{k}$$

5.5 Section 3.2, Checkpoint 3.8

Calculate the following integral:

$$\int_1^3 \left[(2t + 4)\hat{i} + (3t^2 - 4t)\hat{j} \right] dt$$

6 09-11-2025

6.1 Section 3.3, Checkpoint 3.9

Calculate the arc length of the parameterized curve

$$\mathbf{r}(t) = \langle 2t^2 + 1, 2t^2 - 1, t^3 \rangle, \quad 0 \leq t \leq 3$$

Extension: write the arc length parametrization by solving for t in terms of s

6.2 Section 3.3, Checkpoint 3.13

Find the equation of the osculating circle of the curve defined by the vector-valued function

$$y = 2x^2 - 4x + 5 \text{ at } x = 1$$

6.3 Section 3.3, Exercise 3.143

Find the equation for the osculating plane at point $t = \frac{\pi}{4}$ on the curve

$$\mathbf{r}(t) = \cos(2t)\hat{i} + \sin(2t)\hat{j} + t\hat{k}$$

7 09-16-2025 - Midterm 1 tomorrow

Logistics: see @35 on Piazza.

- 50 mins
- no electronics at all
- single-page, typical 8.5x11 or A4 size paper, front and back, can typeset or handwrite
- write your answer in the provided answer box so we can grade faster

Points, vectors, lines, planes, quadrics. See the handout for quadrics that I gave out, and consult the wikipedia if you need help making your formula sheet.

Dot and Cross products and their meanings

- how does the dot product relate to projections/components?
- how does the cross product relate to areas/volumes?
- how do dot and cross products help when calculating work and torque?

Calculus on vector-valued functions (limits, derivatives, integrals)

- Remember examples from previous recitations - sometimes we had you do L'Hopital's or do questions that required knowledge of limits from 2-D calculus.

Arc length + arc length parameterization

- What is the formula?

Curvature + unit tangent, normal, binormal

- Also includes osculating planes/circles
- know the formulas (or write them down on your formula sheet)

motion in space

- apply what you know about projections/components to find out the "component" of the tangential/normal components of acceleration
- plus other questions regarding position, velocity, speed, acceleration.

What can I do to study? **DO A PRACTICE TEST** and check your answers afterwards. Read up on the textbook (see @35 on Piazza for which sections to ignore) and get a good night's sleep. I will be proctoring your exam, so I'll see you all tomorrow :-)

8 09-18-2025 - Post Midterm 1, No Content

9 09-23-2025

9.1 Section 4.1, Checkpoint 4.4

Find the domain and range of the function $f(x, y) = \sqrt{36 - 9x^2 - 9y^2}$

9.2 Section 4.1, Checkpoint 4.5

Find the equation of the level surface of the function

$$g(x, y, z) = x^2 + y^2 + z^2 - 2x + 4y - 6z$$

corresponding to $c = 2$, and describe the surface, if possible.

9.3 Section 4.2, Exercise 60

Evaluate $\lim_{(x,y) \rightarrow (1,2)} x$ or explain why the limit does not exist.

9.4 Section 4.2, Exercise 77

Evaluate $\lim_{(x,y) \rightarrow (0,0)} \ln(x^2 + y^2)$ or explain why the limit does not exist.

*Does the limit exist on **any** path?*

9.5 Section 4.2, Exercise 81

Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{x^4 - 4y^4}{x + 2y^2}$ or explain why the limit does not exist.

Do you see any algebraic manipulations?

9.6 Section 4.2, Exercise 86

Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{xy + y^3}{x^2 + y^2}$ or explain why the limit does not exist.

(a) Along the x -axis ($y = 0$)

(b) Along the y -axis $x = 0$

(c) Along the path $y = 2x$

9.7 Section 4.2, Exercise 88

Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^4 + y^2}$ or explain why the limit does not exist.

(a) Along the x -axis ($y = 0$)

(b) Along the y -axis $x = 0$

(c) Along the path $y = x^2$

9.8 Section 4.2, Exercise 95

Determine the region in which the function is continuous. Explain your answer

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

Hint: maybe this coordinate system isn't helpful. Also remember the Squeeze Theorem?

We do NOT expect you to use Squeeze Theorem with completely general functions


10 09-25-2025

10.1 Section 4.3, Checkpoint 4.13

Calculate $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ for the function $f(x, y) = \tan(x^3 - 3x^2y^2 + 2y^4)$ by holding the opposite variable constant, then differentiating.

10.2 Section 4.3, Exercises 114-117

Calculate the sign of the partial derivative using the graph of the surface f below.



114. $f_x(1, 1)$

115. $f_x(-1, 1)$

116. $f_y(1, 1)$

117. $f_x(0, 0)$

10.3 Section 4.4, Checkpoint 4.19

Find the equation of the tangent plane to the surface defined by the function

$$f(x, y) = x^3 - x^2y + y^2 - 2x + 3y - 2 \text{ at the point } (-1, 3)$$

10.4 Section 4.4, Checkpoint 4.22

Find the differential dz of the function

$$f(x, y) = 4y^2 + x^2y - 2xy$$

and use it to approximate Δz at point $(1, -1)$. Use $\Delta x = 0.03$ and $\Delta y = -0.02$. What is the exact value of Δz ?

11 09-30-2025

11.1 Section 4.5, Checkpoint 4.23

Calculate $\frac{dz}{dt}$ given the following functions. Express your final answer in terms of t .

$$z = f(x, y) = x^2 - 3xy + 2y^2$$

$$x = x(t) = 3 \sin 2t$$

$$y = y(t) = 4 \cos 2t$$

11.2 Section 4.5, Exercise 253

The radius of a right circular cone is increasing at 3 cm/min whereas the height of the cone is decreasing at 2 cm/min. Find the rate of change of the volume of the cone when the radius is 13 cm and the height is 18 cm.

11.3 Section 4.5, Example 4.30

- (a) Calculate $\frac{dy}{dx}$ if y is defined implicitly as a function of x via the equation

$$3x^2 - 2xy + y^2 + 4x - 6y - 11 = 0$$

What is the equation of the tangent line to the graph of this curve at point $(2, 1)$?

- (b) Calculate $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$, given $x^2e^y - yze^x = 0$.

11.4 Section 4.6, Example 4.31

Let $\theta = \arccos(\frac{3}{5})$. Find the directional derivative $D_u f(x, y)$ of $f(x, y) = x^2 - xy + 3y^2$ in the direction of $u = \cos \theta \hat{i} + \sin \theta \hat{j}$. What is $D_u f(-1, 2)$?

Use the definition of directional derivative (x-h defn)

11.5 Section 4.6, Example 4.32

Let $\theta = \arccos(\frac{3}{5})$. Find the directional derivative $D_u f(x, y)$ of $f(x, y) = x^2 - xy + 3y^2$ in the direction of $u = \cos \theta \hat{i} + \sin \theta \hat{j}$. What is $D_u f(-1, 2)$?

Use the partial derivatives

11.6 Section 4.6, Checkpoint 4.30

Find the direction for which the directional derivative of $g(x, y) = 4x - xy + 2y^2$ at $(-2, 3)$ is a maximum. What is the maximum value?

12 10-02-2025

12.1 Section 4.7, Checkpoint 4.35

Use the second derivative to find the local extrema of the function

$$f(x, y) = x^3 + 2xy - 6x - 4y^2$$

12.2 Section 4.7, Checkpoint 4.36

Given $z = f(x, y)$ is continuous and differentiable on a closed, bounded set D , the strategy to find absolute extrema of f on D is to

1. Determine the critical points of f in D
2. Calculate f at each of these critical points
3. Determine the maximum and minimum values of f on the boundary of its domain
4. Find the absolute maximum and minimum values of f by comparing the values from steps 2 and 3.

Use the above strategy to find the absolute extrema of

$$f(x, y) = 4x^2 - 2xy + 6y^2 - 8x + 2y + 3 \text{ on } \{(x, y) : 0 \leq x \leq 2 \text{ and } -1 \leq y \leq 3\}$$

13 10-07-2025 - Midterm 2 tomorrow

Use the method of Lagrange Multipliers to find the maximum and minimum values of the function subject to the given constraints.

13.1 Section 4.8, Exercise 369

Minimize $f(x, y) = x^2 + y^2$ on the hyperbola $xy = 1$.

13.2 Section 4.8, Exercise 373

The curve $x^3 - y^3 = 1$ is asymptotic to the line $y = x$. Find the point(s) on the curve $x^3 - y^3 = 1$ farthest from the line $y = x$.

What is the function we seek to minimize?

13.3 Section 4.8, Exercise 374

Maximize $U(x, y) = 8x^{\frac{4}{5}}y^{\frac{1}{5}}$ under constraint $4x + 2y = 12$.

13.4 Section 4.8, Exercise 379

Maximize $f(x, y, z) = x^2 + y^2 + z^2$ under constraints $x + y + z = 9$ and $x + 2y + 3z = 20$.

13.5 Section 4.8, Exercise 388

A large container in the shape of a rectangular solid must have a volume of $480m^3$. The bottom of the container costs $\$5/m^2$ to construct, whereas the top and sides cost $\$3/m^2$ to construct. Use Lagrange multipliers to find the dimensions of the container of this size that has the minimum cost.

14 10-09-2025 - Post Midterm 2, No Content

20 11-11-2025 - Midterm 3 tomorrow

21 11-13-2025 - Post Midterm 3, No Content

