Please check the examination det	tails below	before ente	ring your can	didate information
Candidate surname			Other name	s
Pearson Edexcel Level 3 GCE	Centre	Number		Candidate Number
Specimen Paper				
(Time: 1 hour 30 minutes)		Paper R	eference <b>9</b>	FM0/3C
Further Mathe Advanced Paper 3C: Further Med				
You must have: Mathematical Formulae and Sta	atistical T	ables, cal	culator	Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

#### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



S61298A
©2018 Pearson Education Ltd.



DO NOT WRITE IN THIS AREA

# Answer ALL questions. Write your answers in the spaces provided.

Unless otherwise indicated, whenever a numerical value of g is required, take  $g = 9.8 \,\mathrm{m\,s^{-2}}$  and give your answer to either 2 significant figures or 3 significant figures.

1. A van of mass 750 kg is moving along a straight horizontal road.

At the instant when the speed of the van is  $v \, \text{m s}^{-1}$ , the resistance to the motion of the van is modelled as a force of magnitude  $(200 + v^2) \, \text{N}$ .

When the engine of the van is working at a constant rate of 12 kW, and the van is moving at a constant speed,

(a) show that the van must be moving at 20 m s<sup>-1</sup>, justifying your answer.

(4)

Later on, the van is moving up a straight road inclined at an angle  $\theta$  to the horizontal,

where 
$$\sin \theta = \frac{1}{15}$$

At the instant when the speed of the van is  $v \, \text{m s}^{-1}$ , the resistance to the motion of the van from non-gravitational forces is modelled as a force of magnitude  $(200 + v^2) \, \text{N}$ .

The engine of the van is now working at a constant rate of 15 kW.

(b) Find the acceleration of the van at the instant when v = 10

(5)

DO NOT WRITE IN THIS AREA

Question 1 continued
(Total for Question 1 is 9 marks)



DO NOT WRITE IN THIS AREA

2.




Figure 1

One end of a light elastic string, of natural length l and modulus of elasticity  $\frac{3}{4}mg$ , is

attached to a particle of mass m. The other end of the string is attached to a fixed point A on a rough inclined plane. The plane is inclined at angle  $\alpha$  to the horizontal, where

$$\tan\alpha = \frac{5}{12}$$

Initially the particle is held at the point B on the plane, where AB = 2l and B lies below A on the line of greatest slope through A, as shown in Figure 1.

The particle is released from rest at B and first comes to instantaneous rest at the point C, where C is between A and B and  $AC = \frac{8}{5}l$ .

Find the coefficient of friction between the particle and the plane.

**(6)** 

DO NOT WRITE IN THIS AREA

Question 2 continued
(Table - O
(Total for Question 2 is 6 marks)



3.	A particle, $P$ , of mass $3  \text{kg}$ is moving with velocity $(2\mathbf{i} + \mathbf{j})  \text{m s}^{-1}$ when it receives an impulse $\mathbf{I}$ of magnitude $\sqrt{130}  \text{N s}$ . Immediately after receiving the impulse, $P$ is moving with velocity $(-\mathbf{i} + \lambda \mathbf{j})  \text{m s}^{-1}$ , where $\lambda$ is a positive constant.	
	(a) Find I, giving your answer in terms of i and j.	(6)
	The angle between the direction of motion of $P$ immediately before receiving the impulse and the direction of motion of $P$ immediately after receiving the impulse is $\theta^{\circ}$	
	(b) Find the value of $\theta$	(6)

DO NOT WRITE IN THIS AREA

Question 3 continued	
(Total for Question 3 is 9 marks)	



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

4.




Figure 2

A light elastic string, of natural length 8l and modulus of elasticity kmg, has its ends attached to two points A and B, where AB = 8l and AB is horizontal.

A pebble, P, of mass m is attached to the midpoint of the string. The pebble rests in equilibrium at a distance 3l vertically below AB, as shown in Figure 2. The pebble is modelled as a particle, and air resistance is modelled as negligible.

(a) Show that 
$$k = \frac{10}{3}$$
 (4)

The pebble is pulled vertically downwards from its equilibrium position until the total length of the string is  $\frac{40}{3}l$ . The pebble is released from rest.

(b) Find the acceleration of P at the instant it is released from rest.

(3)

At the instant the pebble crosses the line AB, the pebble has speed v.

(c) Find v.

(3)

In an experiment, when the natural length of the string was 2 m, it was found that the speed of P at the instant when it crossed the line AB was  $1.5 \,\mathrm{m\,s^{-1}}$ .

(d) Considering the model, suggest a reason, other than air resistance, why the model and the experiment give different values.

(1)



DO NOT WRITE IN THIS AREA

Question 4 continued



Question 4 continued	

DO NOT WRITE IN THIS AREA

Question 4 continued	
	-
	-
	_
	_
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
(Total for Question 4 is 11 marks)	_



DO NOT WRITE IN THIS AREA

# 5. [In this question $\mathbf{i}$ and $\mathbf{j}$ are perpendicular unit vectors in a horizontal plane]




Figure 3

Figure 3 represents the plan view of part of a horizontal floor, where AB and BC represent fixed vertical walls. The direction of  $\overrightarrow{AB}$  is in the direction of the vector  $\mathbf{i}$  and the direction of  $\overrightarrow{BC}$  is in the direction of the vector  $(-\mathbf{i} + 3\mathbf{j})$ .

A small ball is projected along the floor towards wall AB so that, immediately before hitting wall AB, the velocity of the ball is  $(2\mathbf{i} - 3\mathbf{j}) \,\mathrm{m} \,\mathrm{s}^{-1}$ .

The ball hits wall AB and then hits wall BC.

The coefficient of restitution between the ball and wall AB is  $\frac{1}{2}$ 

The coefficient of restitution between the ball and wall BC is  $\frac{1}{3}$ 

The velocity of the ball immediately after hitting wall BC is  $\mathbf{v} \, \mathbf{m} \, \mathbf{s}^{-1}$ .

The floor and the walls are modelled as being smooth. The ball is modelled as a particle.

Show that 
$$\mathbf{v} = \left(-\mathbf{i} + \frac{1}{2}\mathbf{j}\right)$$
.

(12)

DO NOT WRITE IN THIS AREA

Question 5 continued



estion 5 continued	



DO NOT WRITE IN THIS AREA

Question 5 continued	
	(Total for Question 5 is 12 marks)



6. A particle, P, of mass 4m is moving along a straight line on a smooth horizontal plane.
A particle, Q, of mass 3m is at rest on the plane on the same straight line.
Particle P collides directly with particle Q.
Immediately before the collision the speed of P is ku, where k is a constant.
Immediately after the collision the speed of P is u and the speed of Q is 3u/2
The coefficient of restitution between P and Q is e.
(a) (i) Show that there is only one possible value of k.
(ii) State the value of k and the value of e.
(11)
(b) Find the total kinetic energy lost in the collision between P and Q.
(3)

DO NOT WRITE IN THIS AREA

Question 6 continued



uestion 6 continued	



DO NOT WRITE IN THIS AREA

Question 6 continued	
	(Total for Question 6 is 14 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

7.




Figure 4

Two smooth uniform spheres, A and B, are moving with speeds u and 2u respectively on a smooth horizontal surface.

Sphere A has mass 3m and radius 2a. Sphere B has mass 2m and radius 2a.

The centres of the spheres are moving towards each other on parallel paths. The paths are at a distance 3a apart, as shown in Figure 4.

The spheres collide. The coefficient of restitution between A and B is  $\frac{1}{3}$ 

- (a) Show that the magnitude of the impulse received by A in the collision is  $\frac{6\sqrt{7}}{5}mu$ .
- (b) Find the speed of A immediately after the collision.

(3)

(c) State how you have used the fact that the spheres are smooth when considering their collision.

(1)

DO NOT WRITE IN THIS AREA

Question 7 continued	



uestion 7 continued	



DO NOT WRITE IN THIS AREA

Question 7 continued

