Please check the examination details bel	ow before ente	ering your candidate information
Candidate surname		Other names
Centre Number Candidate No	umber	
Pearson Edexcel Level	3 GCE	
	Paper reference	8FM0/25
Further Mathema Advanced Subsidiary Further Mathematics option 25: Further Mechanics 1 (Part of options C, E, H and J	ns	•
You must have: Mathematical Formulae and Statistica	al Tables (Gr	een), calculator

Candidates may use any calculator allowed by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \,\mathrm{m \, s^{-2}}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 40. There are 4 questions.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1. A car of mass 1200 kg moves up a straight road that is inclined to the horizontal at an angle α , where $\sin \alpha = \frac{1}{15}$

The total resistance to the motion of the car from non-gravitational forces is modelled as a constant force of magnitude R newtons.

At the instant when the engine of the car is working at a rate of $32\,kW$ and the speed of the car is $20\,m\,s^{-1}$, the acceleration of the car is $0.5\,m\,s^{-2}$

	Find	the	val	lue	of	R
I ma me varae or re						
I ma me varae of it						
1 1110 0110 1 0110 01 11						

- (5
٠.	J,

	ſ.
XXXXX	ı
	ı
	ı
	ı
	ı
◀	ı
M I	ı
\times 66 \times	ı
4	ı
XXXXX	ı
	ı
×±×	ı
$\otimes \mathbb{Z} \otimes \mathbb{Z}$	ı
	ı
ш	ı
$\times \mapsto \times$	ı
~	ı
	ı
	ı
Ö	ı
	ı
	ı
00	ı
	ı
	ı
******	ı
$\times \times \times \times \times \times$	ı
	ı
$\times\!\!\times\!\!\times\!\!\times\!\!\times$	ı
	ı
	ı
*****	ı
	1
	ı
	ı
	ı
\times	ı
	ı
× co×	ı
	ı
	ı
	ı
	ı
<u> </u>	ı
	ı
× <u>••</u> ×	ı
	1
XX	l

Ž	
Ž	
DON	
DO N	
DO.N	
DO N	
DON	
DOM	
DO N	
N OQ	
NOQ.	
N OQ	
N OQ	
DON	
A DO N	
EA DO N	
(REA DO N	
AREA DO N	
IS AREA DO N	
HIS AREA DO N	
THIS AREA DO N	
N THIS AREA DO N	
IN THIS AREA DO N	
TE IN THIS AREA DO N	
ITE IN THIS AREA DO N	
VRITE IN THIS AREA DO N	
RITE IN THIS AREA DO N	
OT WRITE IN THIS AREA DO N	
WRITE IN THIS AREA DO N	
NOT WRITE IN THIS AREA DO N	
NOT WRITE IN THIS AREA DO N	

Question 1 continued	
	(Total for Question 1 is 5 marks)



2.	Two particles, A and B, have masses m and 3m respectively. The particles are moving in
	opposite directions along the same straight line on a smooth horizontal plane when they
	collide directly.

Immediately before they collide, A is moving with speed 2u and B is moving with speed u.

The direction of motion of each particle is reversed by the collision.

In the collision, the magnitude of the impulse exerted on A by B is $\frac{9mu}{2}$

(a) Find the value of the coefficient of restitution between A and B.

(7)

(b) Hence, write down the total loss in kinetic energy due to the collision, giving a reason for your answer.

(1)

XXXX

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
XXXX
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
XXXX
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
\times
\times
$\times\!\times\!\times\!\times$
\times
\times
\times
××-×
DO NOT WRITE IN THIS AREA
$\otimes \triangle$

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
××××
$\times\!\!\times\!\!\times\!\!\times$
XXXX
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\times\!\times\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\!\times\!\!\!\!\times\!\!\!\!\times\!\!\!\!\times$
AREA
$\times\!\!\times\!\!\!\times\!\!\!\times$
$\times \times \times \times \times$
××××

Ŧ
Ę
E TE
N TH
EINTH
FE IN TH
ITE IN TH
SITE IN TH
/RITE IN TH
WRITE IN TH
WRITEHNTH
T WRITE IN TH
OT WRITE IN TH
OT WRITE IN TH
VOT WRITE IN TH
NOT WRITE IN TH
NOT WRITE IN THIS
$\times \times $
$\times \times $
DO NOT WRITE IN TH
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
$\times \times $
DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
AREA DO
E IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
WRITE IN THIS AREA DO
WRITE IN THIS AREA DO
WRITE IN THIS AREA DO
WRITE IN THIS AREA DO
WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO

Question 2 continued	



Question 2 continued

Question 2 continued
(Total for Question 2 is 8 marks)
(Total for Question 2 is 6 marks)



3. A plane is inclined to the horizontal at an angle α , where $\tan \alpha = \frac{3}{4}$

A particle *P* is held at rest at a point *A* on the plane.

The particle P is then projected with speed $25 \,\mathrm{m\,s^{-1}}$ from A, up a line of greatest slope of the plane.

In an initial model, the plane is modelled as being smooth and air resistance is modelled as being negligible.

Using this model and the principle of conservation of mechanical energy,

(a) find the speed of P at the instant when it has travelled a distance $\frac{25}{6}$ m up the plane from A.

(4)

In a refined model, the plane is now modelled as being rough, with the coefficient of friction between P and the plane being $\frac{3}{5}$

Air resistance is still modelled as being negligible.

Using this refined model and the work-energy principle,

(b) find the speed of P at the instant when it has travelled a distance $\frac{25}{6}$ m up the plane from A.

(8)

XXXX

\longleftrightarrow
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
\longleftrightarrow
$\times\!\!\times\!\!\times\!\!\times$
≪nin/i×
\times
\times
AREA
INTHIS
$\times \times \times \times$
$\times \times \times$
\times
() -
$\times\!\!\times\!\!\times\!\!\times$
\times
$\otimes \boxtimes \times$
\times
\times_{\sim}
$\times \times \times$
$\otimes \otimes \times$
X Z
\times
$\otimes \otimes \otimes$
DO NOT WRITE
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\langle \rangle \rangle \rangle \rangle \rangle$
$\Leftrightarrow \Leftrightarrow \Rightarrow$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
XXXX
XXXX
$\times\!\!\times\!\!\times$
$\!$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\!>$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\!\cdot\!\!\!\cdot\!\!\!\cdot\!\!\!\cdot\!\!\!\cdot$
$\times\!\!\times\!\!\times$
\times
\times
XXXX
AREA
$\sim\sim$
\times
$\times \times \times \times$
\times
Ė
IN THIS
EN TH
TE IN TH
RITERNAL
RITEINTE
WRITE IN TH
WRITEINTH
WRITEHNTH
T WRITE IN TH
OT WRITE IN TH
IOT WRITE IN TH
NOT WRITE
NOT WRITE
NOT WRITE
NOT WRITE
DO NOT WRITE IN TH
NOT WRITE
DO NOT WRITE
A DO NOT WRITE
A DO NOT WRITE
A DO NOT WRITE
A DO NOT WRITE
AREA DO NOT WRITE
AREA DO NOT WRITE
AREA DO NOT WRITE
AREA DO NOT WRITE
AREA DO NOT WRITE
THIS AREA DO NOT WRITE
THIS AREA DO NOT WRITE
THIS AREA DO NOT WRITE
IN THIS AREA DO NOT WRITE
IN THIS AREA DO NOT WRITE
E IN THIS AREA DO NOT WRITE
FE IN THIS AREA DO NOT WRITE
FE IN THIS AREA DO NOT WRITE
FE IN THIS AREA DO NOT WRITE
FE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
NOT WRITE IN THIS AREA DO NOT WRITE
NOT WRITE IN THIS AREA DO NOT WRITE
NOT WRITE IN THIS AREA DO NOT WRITE
OO NOT WRITE IN THIS AREA DO NOT WRITE
WRITE IN THIS AREA DO NOT WRITE
OO NOT WRITE IN THIS AREA DO NOT WRITE

Question 3 continued



Question 3 continued

	ς,	X.	2		
5			4	d	
_	×	е		15	
		79		Ŀ	
	×	. 2	S	Z	
	V	N	ν	ч	
	а				
	×	÷	G	à	
	a	N		₹	
	4				
			4	à	
	(i	ф		г	
	×	94		Ŀ	
	V	$^{\wedge}$		7	
/		V	7	2	
	8	n		9	
	U		r	Я	
1	Δ	Z	2	Z	
	×		7	7	
	Œ				
	×	И	I.	2	
	V.				
	a	ø.			
	м		ú	è	
>	a	К	7	7	
		7	1		
	м		ú	è	
		X	ð	۳	
	М		L	٠	
	4				
١,	7	₹	9	-	
		\rightarrow		>	
	6		ð	G	
	4	и	L	3	
	7	-	•	7	
	4	K.	2	C	
	м	P	ņ	ņ	
2	U	ĸ			
	/1	,	=	,	
	×	~	G	à	
	a	N		₹	
	7				
	Na.	ũ.	Z	7	
		Z			
	K		Ξ	2	
	i,	ä			
١,	ß	₹	7		
_	V				
	a	ĸ	2	_	
	4	7	₹	7	
	v	2	4	S.	
	a	r	7		
	Я	L2		а	
7		79	,	ч	
١.	ă	=	ь	ď	
		2		5	
	а		è	6	
	×	7		7	
	V	△	2	S	
5	S	ø	۹	ú	
0	И	D		3	
	U	×	ò	ø	
١,	Λ	-ã	ú	Z	
	M	r5	7	٩	
	ď	ĸ	۵	4	
	Я	7	۴	7	
1	V	$^{\sim}$		\	
				\rangle	
2				1	
	Λ	\checkmark		2	

Question 3 continued
(Total for Question 3 is 12 marks)



- **4.** A particle P of mass $2m \log 1$ is moving with speed $2u \operatorname{m} \operatorname{s}^{-1}$ on a smooth horizontal plane. Particle P collides with a particle Q of mass $3m \log 1$ which is at rest on the plane. The coefficient of restitution between P and Q is e. Immediately after the collision the speed of Q is $v \operatorname{m} \operatorname{s}^{-1}$
 - (a) Show that $v = \frac{4u(1+e)}{5}$

(6)

(b) Show that $\frac{4u}{5} \leqslant v \leqslant \frac{8u}{5}$

(2)

Given that the direction of motion of P is reversed by the collision,

(c) find, in terms of u and e, the speed of P immediately after the collision.

(2)

After the collision, Q hits a wall, that is fixed at right angles to the direction of motion of Q, and rebounds.

The coefficient of restitution between Q and the wall is $\frac{1}{6}$

Given that P and Q collide again,

(d) find the full range of possible values of e.

(5)



	- 1
	н
****	н
>>>>>	-1
	-1
<<	-1
	-1
\times	-1
<u> </u>	-1
	-1
× vo×	-1
	-1
	-1
$\times \times \times$	-1
$\times\!\!\times\!\!\times\!\!\times$	-1
	-1
\times	-1
×W×	-1
\bowtie	-1
	-1
	-1
	-1
$\times \times \times$	-1
	-1
$\sim\sim\sim$	-1
× - × ×	-1
\times	-1
$\times \odot \times$	-1
DO	-1
	-1

$\times\!\!\times\!\!\times\!\!\times$	
	- 1

*****	-1
XXXX	-1
$\times\!\!\times\!\!\times\!\!\times$	-1
****	-1
$\times\!\!\times\!\!\times\!\!\times$	-1
	-1
	-1
	-1
	-1
$\times \times $	-1
\times ω \times	-1
\times $$ \times	-1
	-1
$\otimes \not\models \otimes$	-1
×2×	-1
\cdots	-1
	-1
	-1
$\times \times $	-1
$\times \underline{\underline{e}} \times$	-1
XX = XXX	-1
XXXXX	-1
$\otimes \boxplus \otimes$	-1
$\times \circ \times$	-1
	-1
	-1
∞	-1
\times	
\bowtie	- 1
$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times$	
\cdots	
$\sim\sim\sim$	- 10
	- 1
A	
A	
A	
REA	
REA	
S AREA	
S AREA	
THIS AREA	
THIS AREA	
N THIS AREA	
IN THIS AREA	
IN THIS AREA	
E IN THIS AREA	
TE IN THIS AREA	
RITE IN THIS AREA	
VRITE IN THIS AREA	
VRITE IN THIS AREA	
T WRITE IN THIS AREA	
T WRITE IN THIS AREA	
VOT WRITE IN THIS AREA	
VOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	

Question 4 continued	



Question 4 continued

<					7
2					S
Þ			X		≺
			X		≺
S			Z		2
Κ					2
Κ		×			\rangle
Ż		×	Š		5
2			×,		S
>			×		K
					≺
۲,			~		2
Κ			>		>
K		×	٦		7
2		4	ë	9	Б
2		Ĵ	₹	Ħ	ij.
>		ì	кі	è	4
Ŋ			ы	ķ	d
Κ			Ŕ	ú	è
K					ì
Z			7	Z	S
2	<				К
			X	7	ę
S				S	2
Κ			la		я
Κ			2	Š	2
K		ł	7	e	5
2			7		5
)		j	¥		6
		ì	K		
۱			P	۹	۴
Κ		8	:>		>
×		d		ĕ	è
1		si	ä	Р	5
2		į		2	3
>		3		9	₹
5		ç		١	4
K			ŭ		J
K			7	ď	2
K			Þ	ø	è
2			ĸ		S
		8	Δ.	7	₹
Ď			Ħ	ø	۴
S		1	ij	á	ė
K		ą	9	6	è
K		d	伛		5
2		á	S		Ą
)			X		Κ
>			K	1	_
Κ			7	5	7
×				ij	è
K			D		3
2			۹	p	5
2		J	7		
>		ì		S	4
S			X		7
Κ			à	ń	×
K			D		1
<			9	ø	5
2		á	и	9	4
\rangle			K	ì	4
S		٦	X	2	7
S					2
K		×			>
K					S
2			Χ		ς
>			Χ		Κ
					<
۲					2
K		×	>		\geq
K		K	\supset		>
2			Χ		٩
>					<
۸,			V		2
×			>		>
K		×			>
2					S
2			×		≺
			×		
b					<
Κ					2
K		K	\Diamond		\rangle
ď		×	\supset		>
2			K		5
2			×		Κ
>					<
`			~		1
Κ		K	\geq		2
K		×	>		>
K					
				2	2
		J	É	j	Ì
Þ		4	É	9	į
3		j	E V		Ì
3		j	N. A.		
?		j	N M		
3		1			
		1			
?					
		1			
		4	Š		
		4	Š		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
		e e e	X		
			X		

	5				5					
						>				
5			(<	×
										C
1	/						١,			0
				2			ς			
1				١,	1					/
							>			
l		/							١	e.
>	⋖								≺	×
1	7		7	١		5				\
7	5		5	l						Ö
>		\geq	0							×
5								S		١,
	>							⟨		
	7				7	4	¥	ľ		
5					7	5	Ħ	Ŋ	<	×
ς	>				а	Ĺ	4	Ŕ		ς,
	>		У	ij	ij	ø	ø	ľ		
	5			à	á	à	ja	r		7
>					⊿		_	۵		×
ς	2	5	?	7	7	7	Ź	2	2	ς,
	>			ú	ġ	۹	P	۲		<
2	\			ø	٩	ú	Ĺ	2		×
\rangle		>		>		S	₹	>		×
5	2			š	2	'n	ŝ			S
	>		Ż		d	ĸ	1	K		
2	S	?		2	Z	2	S	2		7
>		\geq			Ź	Ź	ď	5		×
	2	5	2	ú	Ħ		è	١,		V
	>			4	۲	Ĺ	2	K		
2			S	Ź	3	7	3			8
					4	è	6	Ď		×
5	2							5		×
	>			S	2	S	2			
	\			Ę	5		ø	γ		0
	◁		S	á		2	٤	Þ		×
	Ĉ		(ř	Z	Z	Z	S	Č	×
⟨	>			ď	9	ď	9	۲		C
			١,	Z	>	2	1	ď		
>	₹	>			ч	ν	4	ν		×
ζ	S	₹	S	ij	Ц	Ķ	1	Ķ		ζ
3	\$?	S	ļ	2	Ľ	1	Ķ		8
2	3	?	5	l	7	K	1	Ķ		Š
3	?	?	8		7	5	2	5	5	>
3	>	>			7		2	Ş	3	?
3		>	>	i	2		1	Ş	3	2
3		?	>	i	2		2	Ş	3	
3		>>>>	>	i	2		2	Ş		
3		>>>>	>	i	2		2	Ş	\ \ \	
3		>>>>		1			2	Ş	\ \ \	
3		>>>>>		1			2	Ş		
3		>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>		1			2	Ş		
3		>>>>>		1			2	Ş		
		> > > >		1			2	Ş		
\ \ \ \		> > > >		1	2		2	Ş		
\ \ \ \		> > > >		1			2	Ş		
				1			2	Ş		
		> > > >		1			2	Ş		
		> > > >		1			2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		
							2	Ş		

Question 4 continued



Question 4 continued
(Total for Question 4 is 15 marks)
TOTAL FOR FURTHER MECHANICS 1 IS 40 MARKS

