

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
Level 3 GCE**

Centre Number

Candidate Number

--	--	--	--	--	--

--	--	--	--	--	--

Specimen Paper

(Time: 1 hour 40 minutes)

Paper Reference **8FM0/01**

Further Mathematics

**Advanced Subsidiary
Paper 1: Core Pure Mathematics**

You must have:

Mathematical Formulae and Statistical Tables, calculator

Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

Information

- A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.
- There are 9 questions in this question paper. The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S60739A

©2018 Pearson Education Ltd.

1/1/1



Answer ALL questions. Write your answers in the spaces provided.

1.

$$\mathbf{P} = \frac{1}{2} \begin{pmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix} \quad \mathbf{Q} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

The matrices \mathbf{P} and \mathbf{Q} represent linear transformations, P and Q respectively, of the plane.

The linear transformation M is formed by first applying P and then applying Q .

- (a) Find the matrix M that represents the linear transformation M .

(2)

- (b) Show that the invariant points of the linear transformation M form a line in the plane, stating the equation of this line.

(3)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 1 continued

(Total for Question 1 is 5 marks)



2. (a) Sketch, on an Argand diagram, the set of points

$$X = \{z \in \mathbb{C} : |z - 4 - 2i| < 3\} \cap \left\{z \in \mathbb{C} : 0 \leqslant \arg(z) \leqslant \frac{\pi}{4}\right\}$$

On your diagram

- shade the part of the diagram that is included in the set
- use solid lines to show the parts of the boundary that are included in the set, and use dashed lines to show the parts of the boundary that are not included in the set

(3)

(b) Show that the complex number $z = 5 + 4i$ is in the set X .

(3)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 2 continued

(Total for Question 2 is 6 marks)



3. (a) Find, in terms of the real constant k , the determinant of the matrix

$$\mathbf{M} = \begin{pmatrix} 3 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & k & 2 \end{pmatrix} \quad (2)$$

Three distinct planes, Π_1 , Π_2 and Π_3 , are defined by the equations

$$\begin{aligned}\Pi_1 : \mathbf{r} \cdot \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} &= 4 \\ \Pi_2 : \mathbf{r} &= \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \\ \Pi_3 : x + ky + 2z &= -1\end{aligned}$$

where λ and μ are scalar parameters.

- (b) Find an equation in Cartesian form for

- (i) Π_1
(ii) Π_2 (4)

Given that the three planes Π_1 , Π_2 and Π_3 form a sheaf,

- (c) use the answer to part (a) to explain why $k = -1$ (2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 8 marks)



4. A company manufacturing radios agreed a 20 year contract with a retailer to supply its radios. In the first year of the contract, 500 radios were supplied to the retailer. In each subsequent year, the number of radios supplied to the retailer was 50 more than in the previous year.

The amount received by the company for each radio during year n of the contract was £ $\left(20 + \frac{n^2}{45}\right)$

The total cost of producing the radios during year n was modelled as £ $(1000 + 10n^2)$

- (a) Show that, according to the model, the profit made by the company in year n , £ P_n , is given by

$$P_n = \frac{10}{9}(n^3 + 900n + 7200) \quad (2)$$

- (b) Use the standard results for summations to show that the total profit made by the company in the first N years of the contract, £ T_N , is given by

$$T_N = aN(N^3 + bN^2 + cN + d)$$

where a, b, c and d are constants to be found.

(5)

At the end of the 20 years, the company found that its total profit made from this contract just exceeded £500 000.

- (c) Assess the model in light of this information.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 4 continued

[A large rectangular area for writing, divided into four quadrants by vertical and horizontal lines. The top-left quadrant contains the question header 'Question 4 continued'. The other three quadrants are labeled 'DO NOT WRITE IN THIS AREA' vertically along their left edges.]

(Total for Question 4 is 9 marks)



5.

$$f(z) = 8z^3 + 12z^2 + 6z + 65$$

Given that $\frac{1}{2} - i\sqrt{3}$ is a root of the equation $f(z) = 0$

- (a) write down the other complex root of the equation, (1)
 - (b) use algebra to solve the equation $f(z) = 0$ completely. (3)
 - (c) Show the roots of $f(z)$ on a single Argand diagram. (2)
 - (d) Show that the roots of $f(z)$ form the vertices of an equilateral triangle in the complex plane. (2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 5 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 5 continued

[A large area for writing, consisting of 20 horizontal lines.]

(Total for Question 5 is 8 marks)



S 6 0 7 3 9 A 0 1 7 3 2

6. (a) Prove by induction that, for all $n \in \mathbb{Z}^+$

$$f(n) = n^5 + 4n$$

is divisible by 5

(6)

(b) Show that $f(-x) = -f(x)$ for all $x \in \mathbb{R}$

(1)

(c) Hence prove that $f(n)$ is divisible by 5 for all $n \in \mathbb{Z}$

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

(Total for Question 6 is 9 marks)



7. The population of Zebu cattle in a particular country is modelled by two sub-populations, adults and juveniles. In this model, the only factors affecting the population of the Zebu are the birth and survival rates of the population.

Data recorded in the years preceding 2018 was used to suggest the annual birth and survival rates of the population.

The results are shown in the table below, with values to 2 significant figures. It is assumed that these rates will remain the same in future years.

	Birth rate	Survival rate
Adult population	0.23	0.97
Juvenile population	0	0.87

It is also assumed that $\frac{1}{3}$ of the surviving juvenile population become adults each year.

Let A_n and J_n be the respective sub-populations, in millions, of adults and juveniles, n years after 1st January 2018. Then the adult population in year $n + 1$ satisfies the equation

$$A_{n+1} = 0.97A_n + \frac{1}{3}(0.87)J_n = 0.97A_n + 0.29J_n$$

- (a) Form the corresponding equation for the juvenile population in year $n + 1$ under this model, justifying your values.

(2)

The total population on 1st January 2018 was estimated, to 2 significant figures, as 1.5 million Zebu, with 1.2 million of these being adults.

- (b) Find the value of p and the matrix \mathbf{M} such that the population of Zebu can be modelled by the system

$$\begin{pmatrix} A_0 \\ J_0 \end{pmatrix} = \begin{pmatrix} 1.2 \\ p \end{pmatrix} \quad \begin{pmatrix} A_{n+1} \\ J_{n+1} \end{pmatrix} = \mathbf{M} \begin{pmatrix} A_n \\ J_n \end{pmatrix}$$

Give p to 2 significant figures and each entry of \mathbf{M} to 2 decimal places.

(3)

Using the model formed in (b), find, to 3 significant figures,

- (c) (i) the **total** Zebu population that was present on 1st January 2017
(ii) the predicted **juvenile** Zebu population on 1st January 2025

(5)

As a result of the predictions of this model, it is decided that the country will export 15 000 juveniles to a neighbouring country at the end of each year.

- (d) Adapt the model from 2018 onwards to include this export.

(2)

- (e) State one limitation of this model.

(1)



DO NOT WRITE IN THIS AREA

Question 7 continued

[Large blank area for writing, consisting of 20 horizontal lines.]



Question 7 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 7 continued

[A large rectangular area for writing, divided into four quadrants by vertical and horizontal lines. The top-left quadrant contains the question header 'Question 7 continued'. The other three quadrants are empty for writing.]

(Total for Question 7 is 13 marks)



8.


**Figure 1**

Figure 1 shows a sketch of the curve with equation $y = x^{\frac{2}{3}}, x \geq 0$

The curve passes through the point A with x coordinate 8

The region R_1 is bounded by the curve, the vertical line passing through A and the x -axis.

The region R_2 is bounded by the curve, the horizontal line passing through A and the y -axis.

The solid V_1 is formed by rotating the region R_1 through 360° about the x -axis.

The solid V_2 is formed by rotating the region R_2 through 360° about the y -axis.

- (a) Show that the exact volume of the solid V_1 is $\frac{384\pi}{7}$ (4)

The solids V_1 and V_2 are placed in an empty container. A solid is selected at random and then replaced in the container. This is carried out 10 times.

Given that the probability of selecting each type of solid is proportional to its volume,

- (b) find, to 4 decimal places, the probability that the solid V_2 is selected exactly 8 times. (7)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 8 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 8 continued

[A large rectangular area for writing, divided into four quadrants by vertical and horizontal lines. The top-left quadrant contains the question header 'Question 8 continued'. The other three quadrants are empty for writing.]

(Total for Question 8 is 11 marks)



S 6 0 7 3 9 A 0 2 7 3 2

9. A small comet C is passing near to a planet. The planet can be modelled as a sphere with centre O taken as a fixed point in space, so that the motion of the comet is relative to the origin O .

The diameter of the planet is 13 000 km.

The comet is monitored by satellites orbiting the planet.

When the monitoring begins the comet is at position $146\mathbf{i} + 234\mathbf{j} - 85\mathbf{k}$ and is moving with vector $-21\mathbf{i} - 33\mathbf{j} + 13\mathbf{k}$ every hour, where the units are in thousands of kilometres.

Assuming the comet maintains a straight line course throughout its motion,

- (a) determine whether or not the comet will collide with the planet.

(6)

Two of the satellites, A and B , have position vectors $\overrightarrow{OA} = 5\mathbf{i} + 12\mathbf{k}$ and $\overrightarrow{OB} = 4\mathbf{i} + 12\mathbf{j} - 3\mathbf{k}$ at the beginning of monitoring. They return to these positions every 4 hours.

- (b) Find the expected angle ACB between the comet and the satellites A and B when they first return to their initial positions. Give your answer to the nearest 0.1°

(4)

- (c) Give a reason why the answer to (b) may differ from the true value.

(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 9 continued

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 9 continued

[Large blank area for writing, consisting of 20 horizontal lines.]



Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 9 is 11 marks)

TOTAL FOR PAPER IS 80 MARKS

