

## Functional and logic programming

- written exam -

### **Important:**

1. Subjects are graded as follows: of - 1p; A – 1.5p; B - 2.5p; C - 2.5p; D - 2.5p.
2. Prolog problems will be resolved using SWI Prolog. The following are required: (1) explanation of the code and of the reasoning behind it; (2) recursive model that solves the problem, for all the predicates used; (3) specification of every predicate (parameters and their meaning, flow model, type of the predicate - deterministic/non-deterministic).
3. Lisp problems will be resolved using Common Lisp. The following are required: (1) explanation of the code and of the reasoning behind it; (2) recursive model that solves the problem, for each function used; (3) specification of every function (parameters and their meaning).

**A.** The following function definition in LISP is given

```
(DEFUN F(L)
  (COND
    ((NULL L) 0)
    (> (F (CAR L)) 2) (+ (F (CDR L)) (F (CAR L))))
    (T (+ (F (CAR L)) 1))
  )
)
```

Rewrite the definition in order to avoid the repeated recursive call **(F (CAR L))**. Do NOT redefine the function. Do NOT use SET, SETQ, SETF. Justify your answer.

**B.** Given a linear list containing positive numbers, write a SWI-Prolog program that returns (as a list of pairs) all possible partitions of the initial list in two sublists, such that the sum of the digits of the elements from the two sublists is the same (we assume there is at least one such partition of the initial list). **For example**, for the list [28, 21, 52, 34, 7], the result should be (not necessarily in this order): [[52, 28], [7, 34, 21]], [[34, 28], [7, 52, 21]], [[7, 28], [34, 52, 21]], [[34, 52, 21], [7, 28]], [[7, 52, 21], [34, 28]], [[7, 34, 21], [52, 28]]].

**C.** Write a PROLOG program that generates the list of permutations of the set  $1..N$ , having the property that the absolute value of the difference between 2 consecutive values from the permutation is  $\geq 2$ . Write the mathematical models and flow models for the predicates used. For example, for  $N=4 \Rightarrow [[3,1,4,2], [2,4,1,3]]$  (not necessarily in this order).

**D.** Write a Lisp function to substitute all numerical values at any level of a given nonlinear list with a given value **e**. **A MAP function shall be used.**

**Example**, for the list (1 d (2 f (3))), **e**=0 the result is (0 d (0 f (0))).