

Functional and logic programming

- written exam -

Important:

1. Subjects are graded as follows: of - 1p; A – 1.5p; B - 2.5p; C - 2.5p; D - 2.5p.
2. Prolog problems will be resolved using SWI Prolog. The following are required: (1) explanation of the code and of the reasoning behind it; (2) recursive model that solves the problem, for all the predicates used; (3) specification of every predicate (parameters and their meaning, flow model, type of the predicate - deterministic/non-deterministic).
3. Lisp problems will be resolved using Common Lisp. The following are required: (1) explanation of the code and of the reasoning behind it; (2) recursive model that solves the problem, for each function used; (3) specification of every function (parameters and their meaning).

A. Given the following PROLOG predicate definition **f(integer, integer)**, with the flow model (i, o):

f(20, -1):-!.

f(I,Y):-J is I+1, **f(J,V)**, V>0, !, K is J, Y is K.

f(I,Y):-J is I+1, **f(J,V)**, Y is V-1.

Rewrite the definition in order to avoid the recursive call **f(J,V)** in both clauses. Do NOT redefine the predicate. Justify your answer.

B. Given a nonlinear list containing both numerical and non-numerical atoms, write a LISP program that calculates the total number of non-numerical atoms at the superficial level of the sublists (including the original list) having the first numerical atom (at any level) an even number. For example, for the list (A B 12 (5 D (A F (10 B) D (5 F) 1)) C 9) the result will be 7.

C. Write a PROLOG program that generates the list of all subsets with values between the $[a, b]$ interval such that the sum of elements from each subset is an odd value. Write the mathematical models and flow models for the predicates used. For example, for $\mathbf{a}=2$ and $\mathbf{b}=4 \Rightarrow [[2,3],[3,4],[2,3,4]]$ (not necessarily in this order).

D. Write a Lisp function to substitute all numerical values at any level of a given nonlinear list with a given value **e**. **A MAP function shall be used.**

Example, for the list (1 d (2 f (3))), **e**=0 the result is (0 d (0 f (0))).