


2 Chapter 4: Dynamics, Force and Newton's Laws of Motion

1. A 5×10^5 kg rocket is accelerating straight up. The thrusters produce an upward force of 1.25×10^7 N, and the force of air resistance is 4.5×10^6 N downward. (a) Draw a free-body diagram including the weight of the rocket, the thrust, and air resistance. (b) What is the rocket's acceleration?



b) $T - F_g - W = ma_y$

$$a = \frac{1.25 \times 10^7 \text{ N} - 4.5 \times 10^6 \text{ N} - (5 \times 10^5 \text{ kg} \cdot 9.8 \text{ m/s}^2)}{5 \times 10^5 \text{ kg}}$$

$a = 6.20 \text{ m/s}^2$

2. A football player with mass 70 kg pushes on a player with mass 90 kg. (a) According to Newton's 3rd law, if the first player exerts a force of 700 N on the second player, what is the force the second player exerts on the first player?

a) player 1 = 700 N equal & opposite force
 player 2 = -700 N

$F_{12} = 700 \text{ N} \rightarrow F_{21} = -700 \text{ N}$



3. A rocket sled is decelerated at a rate of 200 m/s^2 , and it has a mass of 2000 kg. There is a constant air resistance force of 1000 N. What additional force is required to give the rocket the deceleration?

$a = -200 \text{ m/s}^2$

$m = 2000 \text{ kg}$


air r = 1000 N = -1000 N

$F - 1000 \text{ N} = (200 \text{ m/s}^2)(2000 \text{ kg})$

$F = -4.0 \times 10^5 \text{ N} + 1000 \text{ N}$

$F = -3.99 \times 10^5 \text{ N}$

4. A 76.0-kg person is being pulled away from a burning building as shown in Fig. 1. (a) Draw a free-body diagram including the two tension vectors and the woman's weight. (b) Write down an expression for $F_{net,x}$. (c) Write down an expression for $F_{net,y}$. (d) Assuming $\vec{F}_{net} = 0$, calculate the tension in the two ropes.



$T_1 = \text{Tension 1}$

$T_2 = \text{tension 2}$


$w_f = \text{weight force}$

$w_f = 76 \text{ kg} \times 9.8 \text{ m/s}^2$
 $= 744.8 \text{ N}$

b) $F_{net,x} = ma_x = 76 \text{ kg}(a_x)$

c) $F_{net,y} = ma_y = 76 \text{ kg}(a_y)$

$T_2 = 193.803 \text{ N}$



d. $F_{net} = 0$

x-position: $T_2 \cos 10 - T_1 \sin 15 = 0$

$T_2 = \frac{T_1 \sin 15}{\cos 10} = T_2 = 737.425 \text{ N} (\sin 15)$

$\cos 10$

$= 193.803$

y-position: $T_2 \sin 10 + T_1 \cos 15$

$-T_1 = 744.8 \text{ N} - T_2 \sin 10$

$\cos 15$

$T_1 = 744.8 \text{ N} - \frac{T_2 \sin 10}{\cos 15} (\sin 10)$

$\cos 15$

Figure 1: A person is pulled from a building using tension in a rope.


$T_1 \cos 15 + T_1 \frac{\sin 15}{\cos 10} (\sin 10)$

$1.01 T_1 = \frac{744.8 \text{ N}}{1.01} = 737.43 \text{ N}$

3 Chapter 5: Friction, Drag, and Elasticity

1. Suppose you have a 120-kg wooden crate resting on a wood floor. The coefficients of static and kinetic friction are 0.5 and 0.3, respectively. (a) What maximum force can you exert horizontally on the crate without moving it? (b) If you continue to exert this force once the crate starts to slip, what will the magnitude of its acceleration then be?


a)



weight force
 $= 120 \text{ kg} (9.81) = 1176 \text{ N}$
 $F_s = 0.5 (1176 \text{ N}) = 588 \text{ N}$

The max horizontal force on the crate without moving is $\leq 588 \text{ N}$

b) $F_k = 0.3 (1176 \text{ N}) = 352.8 \text{ N}$
 $F_s - F_k = ma$
 $588 \text{ N} - 352.8 \text{ N} = 120 \text{ kg} a$
 $a = \frac{235.2 \text{ N}}{120 \text{ kg}} = 1.96 \text{ m/s}^2$



Free-body diagram

Figure 2: A skier slides down a slope.

2. Suppose a skier (Fig. 2) is sliding down a slope with an incline of 25 degrees. If the coefficient of kinetic friction is 0.1, what is the skier's acceleration?

$$F = ma \quad w_x = w \sin 25 \quad f_k = 0.1 (w \cos 25) \quad w_x - f_k = ma \quad w = mg$$

$$\sin 25 - (0.1 \cos 25) = \frac{a}{g}$$

$$g (\sin 25 - 0.1 \cos 25) = a \quad a = 9.8 (\sin 25 - 0.1 \cos 25) = 3.25 \text{ m/s}^2$$

3. **Drag Force.** Suppose the skier reaches a top speed of 40 m/s. If his area is 0.75 m², the density of air is 1.225 kg m⁻³, and $C = 0.75$, what is the magnitude of the drag force in Newtons?

$$F_D = \frac{1}{2} \rho v^2 C_D A = \frac{1}{2} (1.225 \text{ kg m}^{-3}) (40 \text{ m/s})^2 (0.75) (0.75 \text{ m}^2)$$

$$F_D = 551.25 \text{ N}$$

4. A mass of 2300 kg is placed on top of a 10.0 m long wooden beam with radius of 4 cm. If the length of the beam decreases by 3 mm, what is the Young's modulus of the wood? Pay attention to the units.

$$\Delta L = \frac{mgL}{YA}$$

$m = 2300 \text{ kg} \quad g = 9.81 \text{ m/s}^2$
 $L_0 = 10 \text{ m} \quad r = 4 \text{ cm} = 0.04 \text{ m}$
 $\Delta L = 0.003 \text{ m}$

$$Y = \frac{2300 \text{ kg} (9.81 \text{ m/s}^2) (10.04)}{\pi (0.04 \text{ m})^2 (0.003 \text{ m})}$$

$$Y = 1.494 \times 10^{10} \text{ N/m}^2$$

4 Chapter 6: Uniform Circular Motion and Gravitation

1. A pitcher in baseball pitches a ball at 144 km per hour, and the ball rotates around his arm at a radius of 0.5 meters. What is the angular velocity of the ball as he throws it, in radians per second?

$$v = r\omega \quad r = 0.5$$

$$\omega = \frac{v}{r} = \frac{40 \text{ m/s}}{0.5} = 80 \text{ rad/sec}$$

$144 \text{ km/hr} \left(\frac{1000 \text{ m}}{1 \text{ km}} \right) \left(\frac{1 \text{ hr}}{3600 \text{ s}} \right) = 40 \text{ m/s}$

2

1.

2. What is the ideal banking angle for a gentle turn of 0.9 km radius on a highway with a 120 km per hour speed limit, assuming everyone travels at the limit?

$$r = .9 \text{ km} \left(\frac{1000 \text{ m}}{1 \text{ km}} \right) = 900 \text{ m}$$

$$v = 120 \text{ km/hr} \left(\frac{1000 \text{ m}}{1 \text{ km}} \right) \left(\frac{1 \text{ hr}}{3600 \text{ s}} \right) = 33.33 \text{ m/s}$$

$$\theta = \tan^{-1} \left(\frac{(33.33 \text{ m/s})^2}{900 \text{ m} (9.81)} \right) = 7.1780$$


a) n

2.

3. Two race car drivers routinely navigate a turn as shown in Fig. 3. (a) Which path may be taken at a higher speed, if both paths correspond to the same force of friction and centripetal force? (b) Suppose path 1 has a radius of curvature of 400 m, and path 2 has a radius of curvature of 800 m. The coefficient of friction is 1.0. If the force of friction balances the centripetal force, what are the tangential velocities of each race car?

a) p

a path 2 because the curve looks more straight & has a larger radius. It looks like the car is traveling faster when turning.



b) path 1: $r = 400 \text{ m}$

path 2: $r = 800 \text{ m}$

$$v = \sqrt{kg r}$$

path 1 $\rightarrow v = \sqrt{1.0 (9.81 \text{ m/s}^2) (400)}$
 $= 62.61 \text{ m/s}$

path 2 $\rightarrow v = \sqrt{1.0 (9.81 \text{ m/s}^2) (800)}$
 $= 88.54 \text{ m/s}$

Figure 3: Two race cars take a turn at different radii of curvature.

4. **Two bonus points:** The existence of the dwarf planet Pluto was proposed based on irregularities in Neptune's orbit. Pluto was subsequently discovered near its predicted position. But it now appears that the discovery was fortuitous, because Pluto is small and the irregularities in Neptune's orbit were not well known. To illustrate that Pluto has a minor effect on the orbit of Neptune compared with the closest planet to Neptune, (a) calculate the acceleration due to gravity at Neptune due to Pluto when they are $4.5 \times 10^{12} \text{ m}$ apart, as they are now. The mass of Pluto is $1.4 \times 10^{22} \text{ kg}$. (b) Now calculate the acceleration due to gravity at Neptune due to Uranus, presently about $2.5 \times 10^{12} \text{ m}$ apart, and compare it with that due to Pluto. The mass of Uranus is $8.62 \times 10^{25} \text{ kg}$.

a) $a_c = \frac{Gm}{r^2}$
 $m = 1.4 \times 10^{22} \text{ kg}$
 $r = 4.5 \times 10^{12} \text{ m}$

$$G = 6.674 \times 10^{-11} \text{ N}$$

b) $a_c = \frac{Gm}{r^2}$
 $m = 8.62 \times 10^{25} \text{ kg}$
 $r = 2.5 \times 10^{12} \text{ m}$

$$a_c = \frac{(6.674 \times 10^{-11}) (1.4 \times 10^{22})}{(4.5 \times 10^{12})^2} = 4.614 \times 10^{-14} \text{ m/s}^2$$

$$a_c = \frac{(6.674 \times 10^{-11}) (8.62 \times 10^{25})}{(2.5 \times 10^{12})^2}$$

$$= 9.204 \times 10^{-10} \text{ m/s}^2$$

Uranus due to Neptune has a greater acceleration than Pluto due to Neptune because Uranus mass is greater.