

Algebra-Based Physics-1: Mechanics, Unit 6

Jordan Hanson

November 4, 2024

Whittier College Department of Physics and Astronomy

Unit 6 Summary

Unit 6 Summary

1. Definition of **momentum**
2. *Conservation of momentum*
 - The proof and the assumptions
 - Examples
3. *Classification of collisions*
 - Elastic
 - Inelastic
 - $1 \rightarrow 1, 1 \rightarrow n, n \rightarrow 1, n \rightarrow n$
 - Lab activity
4. Momentum in multiple dimensions
5. Center of mass
 - Derivation of $\vec{F}_{\text{Net}} = \frac{d\vec{p}_{\text{CM}}}{dt}$
 - Center of mass motion

Definition of Momentum

Definition of momentum

Definition of Momentum

A particle of mass m and velocity $\vec{v}$ has the vector *momentum*:

$$\vec{p} = m\vec{v}$$

Definition of momentum

There is a corollary:

Newton's Second Law with momentum

If a particle has acceleration $\vec{a} = \frac{d\vec{v}}{dt}$, then

$$\vec{F}_{\text{Net}} = \frac{d\vec{p}}{dt}$$

Definition of momentum

An object that has a small mass and an object that has a large mass have the same momentum. Which mass has the largest kinetic energy?

- A: The one with the small mass
- B: The one with the large mass
- C: If the momentum is the same the kinetic energy is the same
- D: Cannot determine the answer

Definition of momentum

An object that has a small mass and an object that has a large mass have the same kinetic energy. Which mass has the largest momentum?

- A: The one with the small mass
- B: The one with the large mass
- C: If the momentum is the same the kinetic energy is the same
- D: Cannot determine the answer

Definition of momentum

The unit of linear momentum is kg m/s . Suppose that a raindrop reaches a terminal velocity of 10 m/s , and the density of water is 1 gram per cm^3 . If a 1 cm^3 water droplet reaches terminal velocity, what is the momentum of the droplet?

- A: 10^{-3} kg m/s
- B: 10^{-2} kg m/s
- C: 10^{-1} kg m/s
- D: 1 kg m/s

Definition of momentum

An small meteor has a mass of 10^3 kg and a velocity of 10 km/s as it enters Earth's atmosphere. What is the initial momentum?

- A: 10^7 kg m/s
- B: 10^6 kg m/s
- C: 10^5 kg m/s
- D: 10^4 kg m/s

Definition of momentum

If it drag of the atmosphere brings the meteor particles to rest, the final momentum is zero. If the average net force on the system is $F_{\text{Net}} = \frac{\Delta p}{\Delta t}$, and the meteorite exists for 1 second, what is the average force on the meteorite?

- A: 10^7 N
- B: -10^7 N
- C: 10^6 N
- D: -10^6 N

Conservation of Momentum

Conservation of Momentum

Conservation of Momentum

The momentum of a system of N particles undergoing no external forces is conserved.

$$\frac{d\vec{P}}{dt} = 0$$

Conservation of Momentum

Suppose two objects with momenta $\vec{p}_1 = m_1\vec{v}_1$ and $\vec{p}_2 = m_2\vec{v}_2$ collide. The new momenta after the collision are $\vec{p}'_1 = m_1\vec{v}'_1$ and $\vec{p}'_2 = m_2\vec{v}'_2$. If $m_2 = 2m_1$ and $\vec{v}_1 = 2\vec{v}_2$, and $\vec{v}'_2$ is observed to equal $\vec{v}_1$, what is $\vec{v}'_1$?

Solve together in groups on boards.

Conservation of Momentum

The proof of conservation of momentum is the combination of two concepts: **Newton's 3rd Law** and **Newton's 2nd Law**. The net forces on two particles by Newton's 3rd Law are

$$\vec{F}_{21} = -\vec{F}_{12} \quad (1)$$

Substituting Newton's 2nd Law for the forces,

$$m_1 \vec{a}_1 = -m_2 \vec{a}_2 \quad (2)$$

Acceleration is defined as the change in velocity, implying

$$m_1 \frac{d\vec{v}_1}{dt} = -m_2 \frac{d\vec{v}_2}{dt} \quad (3)$$

$$\frac{d\vec{p}_1}{dt} = -\frac{d\vec{p}_2}{dt} \quad (4)$$

$$\frac{d\vec{p}_1}{dt} + \frac{d\vec{p}_2}{dt} = 0 \quad (5)$$

$$\frac{d}{dt} (\vec{p}_1 + \vec{p}_2) = 0 \quad (6)$$

Conservation of Momentum

$$\boxed{\frac{d}{dt} (\vec{p}_1 + \vec{p}_2) = 0} \quad (7)$$

Equation 7 states that the total momentum does not change with time. The assumptions hold even if there are more than two particles, for every particle in the system exerts some force on every other particle, even if that force is zero.

$$\frac{d}{dt} \sum_i \vec{p}_i = \frac{d\vec{P}}{dt} = 0 \quad (8)$$

Conservation of Momentum

What two assumptions were necessary in the above proof?

- A: Each mass is constant in time, and the total velocity is zero.
- B: The total velocity is zero, and the net force is $\frac{dP}{dt}$.
- C: The net external force is zero, and each mass is constant in time.
- D: The net external force is zero, and the mass of each particle is zero.

Conservation of Momentum




Figure 1: The assumptions for momentum conservation. (a) Which car exerts more force? (b) At which point are the cars accelerating?

Conservation of Momentum

Colliding Carts

Two carts in a physics lab roll on a level track, with negligible friction. These carts have small magnets at their ends, so that when they collide, they stick together (**Figure 9.16**). The first cart has a mass of 675 grams and is rolling at 0.75 m/s to the right; the second has a mass of 500 grams and is rolling at 1.33 m/s, also to the right. After the collision, what is the velocity of the two joined carts?




Figure 9.16 Two lab carts collide and stick together after the collision.

Figure 2: The laboratory activity for today.

- One cart still, other moving (magnet side)
- One cart still, other moving (velcro side)
- Both carts moving (magnet side)
- Both carts moving (velcro side)

Conservation of Momentum

Test cases from lab:

- One cart still, other moving (magnet side)
- One cart still, other moving (velcro side)
- Both carts moving (magnet side)
- Both carts moving (velcro side)

Answer the following questions:

- In which of the above is the momentum conserved?
- In which of the above is the kinetic energy conserved?
- In which of the above is both the kinetic energy and momentum conserved?
- In which of the above is the final kinetic energy zero?

Conservation of Momentum




Figure 3: Classification of momentum interactions.

Special case of an explosion...

<https://www.youtube.com/watch?v=5zxVQBnmyDA>

Conservation of Momentum




Figure 4: Knowledge of the atom via momentum conservation! To understand the recoil energy, though, we have to understand the different possibilities: elastic or inelastic.

Elastic Collisions

Elastic Collisions




Figure 5: Internal kinetic energy and momentum are conserved if the collision is *elastic*.

Elastic Collisions

Suppose two objects undergo an elastic collision. Given the conditions below, find the quadratic equation for v_1' .

$$m_1 = 0.5\text{kg} \quad (9)$$

$$m_2 = 1.0\text{kg} \quad (10)$$

$$v_1 = 2\text{m/s} \quad (11)$$

$$v_2 = 0\text{m/s} \quad (12)$$

Solve in groups on boards.

Elastic Collisions

Suppose two objects undergo an elastic collision. Given the conditions below, find the quadratic equation for v_1' .

Answer: $\frac{3}{2}v_1'^2 - 2v_1' - 2 = 0$

Which root of this equation is correct, and why?

Inelastic Collisions

Inelastic Collisions




Figure 6: Internal kinetic energy is not conserved, and momentum is conserved if the collision is *inelastic*.

Inelastic Collisions

Suppose two objects undergo an inelastic collision. Given the conditions below,

$$m_1 = 0.5\text{kg} \quad (13)$$

$$m_2 = 1.0\text{kg} \quad (14)$$

$$v_1 = 2\text{m/s} \quad (15)$$

$$v_2 = 0\text{m/s} \quad (16)$$

Solve in groups on boards.

Inelastic Collisions

What is the right answer?

$v' = \frac{2}{3}$ m/s. Should it be positive or negative?

Conclusion

Unit 6 Summary

1. Definition of **momentum**
2. *Conservation of momentum*
 - The proof and the assumptions
 - Examples
3. *Classification of collisions*
 - Elastic
 - Inelastic
 - $1 \rightarrow 1, 1 \rightarrow n, n \rightarrow 1, n \rightarrow n$
 - Lab activity
4. Momentum in multiple dimensions
5. Center of mass
 - Derivation of $\vec{F}_{\text{Net}} = \frac{d\vec{p}_{\text{CM}}}{dt}$
 - Center of mass motion