

Wednesday Warm Up: Unit 0, Review of 135A

Prof. Jordan C. Hanson

January 22, 2025

1 Warm-Up Exercises

1. Suppose a ship sails at 20 km per hour for 3 hours to the West, and then at 20 km per hour for 2 hours to the South. (a) Assuming that the ship starts at the origin of a 2D coordinate system, what is the final location of the ship? (b) What is the average velocity?
2. Suppose an athlete starts a race from rest at $t = 0$ with a constant acceleration of $a = 2.5 \text{ m s}^{-2}$. (a) How long before the speed of the runner is 5 m s^{-1} ? (b) If the runner has a mass of 60 kg, what is the kinetic energy? (c) How much work or energy was required to reach this velocity?
3. Suppose two children each pull a toy in opposite directions. One pulls to the right with a force of 10 N, while the other pulls to the left with a force of 8 N. The toy weighs 0.5 kg. (a) What is the magnitude and direction of the acceleration of the toy? (b) If the toy begins at rest, where is the toy after 2 seconds? (c) If the kids drop this same toy from a height of 10 meters, what will be the final velocity just before it hits the ground?
4. Suppose a physical therapy patient is asked to shove a medicine ball forward off the edge of a table to help rebuild the strength of their shoulders. The medicine ball weighs 7 kg. (a) If the patient is able to give the ball a speed of 1 m s^{-1} , what is the momentum of the ball? (b) If the patient gives the ball the same momentum by rolling it, and it strikes elastically a ball with a mass of 3.5 kg, what will be the velocity of the second ball?
5. Suppose the medicine ball in the previous problem has a mass of 3.5 kg, and a diameter of 10 cm. The moment of inertia for a solid sphere is $I = \frac{2}{5}mr^2$. (a) What is the angular momentum of the ball if it is spun at 1 rotation per second? (b) What is the rotational kinetic energy?
6. The SI unit of **charge** is called a *Coulomb* (1 C). A Coulomb is the amount of charge that flows through a wire in one second, given that the *current* is 1 *Amp*. Define current as $I = \Delta Q / \Delta t$, where Q is the charge as a function of time, and ΔQ is the change in the charge. (a) Suppose we are charging a car battery with an average current of $I = \Delta Q / \Delta t = 2$ Amps, for 5 hours. How many Coulombs of charge did we put in the battery? (b) If a battery has 7 Amp hr of charge, for how long can it provide 0.5 Amps of current?