

Midterm 2 for Calculus-Based Physics-1: Mechanics (PHYS150-O1)

Dr. Jordan Hanson - Whittier College Dept. of Physics and Astronomy

October 16th, 2017

1 Vectors and Newton's Laws

1. Let $\vec{F}_1 = \frac{3}{2}\hat{x} + 2\hat{y}$ N, and $\vec{F}_2 = -2\hat{x} + \frac{3}{2}\hat{y}$ N. a) Give the magnitude of each force. b) What is the net force? c) What is the angle between these two forces?
2. Imagine you are sitting in an airplane that has just lifted off with an acceleration vector 45 degrees with respect to horizontal. Draw a free-body diagram corresponding to you, showing all forces acting on you.
3. Imagine you are riding a skateboard down a hill. Draw a free-body diagram corresponding to you, showing all forces acting on you.

2 Newton's Laws, and Circular Motion

bank.png

Figure 1: Let the weight be \vec{w} , and the total lift be \vec{L} , which may be broken into two components: the turning force (equal to centripetal force \vec{f}_C) and vertical lift (which balances weight).

1. When banking, the free-body diagram of a jet-fighter resembles Fig. ???. To bank while maintaining altitude, the lift force \vec{L} must *both* balance the weight \vec{w} and provide the centripetal force \vec{f}_C . Let the mass of the aircraft be m , the radius of the turn be r , and the angle between \vec{L} and horizontal be θ .

- Show that the angular velocity of the turn, ω , is $\omega = \sqrt{\frac{L \cos \theta}{rm}}$

- If ω is the angular velocity, then the *period* is $T = 2\pi/\omega$. This is the time required to fly in a complete circle. Show that one-half period is $\frac{T}{2} = \pi \sqrt{\frac{rm}{L \cos \theta}}$. This is the time required to turn.

- Let $L = 8 \times 10^5$ N, $m = 2 \times 10^4$ kg, $r = \frac{1}{2}$ km, and $\theta = 60$ degrees. How long does it take the jet fighter to turn?
- What is the speed of the jet fighter?
 - A: 10 m/s
 - B: 20 m/s
 - C: 100 m/s
 - D: 120 m/s

3 Frictional Forces

1. There is a spill of a mystery toxic liquid on a shop floor, and no one wants to touch it. Someone gets the bright idea that they can identify it by the coefficient of kinetic friction and a steel plate. Draw a free body diagram corresponding to a steel plate sliding along the liquid/floor, with friction decelerating it.
2. What is the coefficient of kinetic friction, μ_k , if a steel plate with an initial speed of 5 m/s comes to a stop after 2.5 seconds, assuming $g = 10$ m/s²? (Use the definition of acceleration $\Delta v / \Delta t = a$).
 - 0.1
 - 0.2
 - 0.5
 - 1.2
3. Suppose they get a sample of the mystery liquid in a vile. They assume the drag force is given by Stoke's Law, $F_D = 6\pi r\eta v$, where v is the velocity of a particle moving through the fluid, r is the radius of the particle, and η is the *viscosity*. They drop a bead with $r = 1$ mm and a mass of one gram into the fluid, and observe the bead sink with a constant (terminal) velocity of 1 m/s. What is the viscosity of the fluid? Units: kg/(m s).
 - $5/(3\pi)$ kg/(m s)
 - $5/(30\pi)$ kg/(m s)
 - 10 kg/(m s)
 - 5 kg/(m s)