

Solutions for Homework 2

Dr. Jordan Hanson - Whittier College Dept. of Physics and Astronomy

February 26, 2022

1 Problem 2.5

Find the electric field a distance z above the center of a circular loop of radius r that carries a uniform line charge λ .

Start by filling in the pieces of the Coulomb effect:

$$d\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{dq'}{r^2} \hat{\mathbf{r}} \quad (1)$$

- $\mathbf{r} = z\hat{\mathbf{z}} - R\hat{\mathbf{s}}$
- $r = \sqrt{z^2 + R^2} = z\sqrt{1 + (R/z)^2} = z\sqrt{1 + \epsilon^2}$. (If $\epsilon \rightarrow 0$, this represents the far-field).
- $r^2 = z^2 + R^2$
- $\hat{\mathbf{r}} = (\hat{\mathbf{z}} - \epsilon\hat{\mathbf{s}})/(1 + \epsilon^2)^{1/2}$
- $dq' = \lambda R d\phi'$, because cylindrical coordinates are the correct choice here.
- Note that $\epsilon = R/z$, so if $z = 0$ then $\epsilon \rightarrow \infty$, and $\epsilon \rightarrow 0$ if $z \gg R$.
- $Q = 2\pi R\lambda$, the total charge.

Integrate to add up the $d\mathbf{E}$ to find \mathbf{E} :

$$\mathbf{E} = \frac{\lambda R}{4\pi\epsilon_0 z^2 (1 + \epsilon^2)^{3/2}} \int_0^{2\pi} d\phi' (\hat{\mathbf{z}} - \epsilon\hat{\mathbf{s}}) \quad (2)$$

By symmetry, the $\hat{\mathbf{s}}$ term evaluates to zero. The result is

$$\mathbf{E} = \frac{2\pi\lambda R\hat{\mathbf{z}}}{4\pi\epsilon_0 z^2 (1 + \epsilon^2)^{3/2}} = \frac{Q\hat{\mathbf{z}}}{4\pi\epsilon_0 z^2 (1 + \epsilon^2)^{3/2}} \quad (3)$$

Checks: $\mathbf{E} = 0$ if $z = 0$ because $\epsilon \rightarrow \infty$. Also, we find the far-field effect if $z \ll R$, because $\epsilon \rightarrow 0$. The units also check out.

2 Problem 2.6

Find the electric field a distance z above the center of a flat circular disc of radius R that carries a uniform surface charge σ . What does your formula give in the limit $R \rightarrow \infty$? Also check the case $z \gg R$.

Start by filling in the pieces of the Coulomb effect:

$$d\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{dq'}{r^2} \hat{\mathbf{r}} \quad (4)$$

- $\mathbf{r} = z\hat{\mathbf{z}} - s\hat{\mathbf{s}}$. As in Problem 2.5, the $\hat{\mathbf{s}}$ -component will vanish upon integration.
- $r^2 = z^2 + s^2$
- $\hat{\mathbf{r}} = (z\hat{\mathbf{z}} - s\hat{\mathbf{s}})/(z^2 + s^2)^{1/2}$
- $dq' = \sigma da' = s ds d\phi$, because cylindrical coordinates work best here.
- $Q = \pi R^2 \sigma$ is the total charge.
- Let $z \tan \theta = s$, so that $ds = z \sec^2 \theta d\theta$, and $\theta_0 = \tan^{-1}(R/z)$

Integrate to add up the $d\mathbf{E}$ to find \mathbf{E} :

$$\mathbf{E} = \frac{\sigma}{2\epsilon_0} z \hat{z} \int_0^R \frac{s ds}{(s^2 + z^2)^{3/2}} \quad (5)$$

$$\mathbf{E} = \frac{\sigma}{2\epsilon_0} \cos \theta \Big|_0^{\theta_0} \hat{z} = \frac{\sigma}{2\epsilon_0} (1 - \cos \theta_0) \hat{z} \quad (6)$$

We know what is $\tan \theta = R/z$, but what is $\cos \theta_0$? (*Hint: draw the triangle and then find the hypotenuse*). The result is

$$\mathbf{E} = \frac{\sigma \hat{z}}{2\epsilon_0} \left(1 - \frac{z}{\sqrt{z^2 + R^2}} \right) \quad (7)$$

The units check automatically because of the units of σ (Coulombs per meter squared), and the ϵ_0 in the denominator. If $R \rightarrow \infty$, then the second term vanishes and the field is

$$\mathbf{E} \rightarrow \frac{\sigma \hat{z}}{2\epsilon_0} \quad (8)$$

This form of the field is the boundary condition near a charged surface. To check the limit that $z \gg R$, first factor a z :

$$\mathbf{E} = \frac{\sigma \hat{z}}{2\epsilon_0} z \left(z^{-1} - (z^2 + R^2)^{-1/2} \right) = \frac{\sigma \hat{z}}{2\epsilon_0} z \left(z^{-1} - z^{-1} (1 + (R/z)^2)^{-1/2} \right) \quad (9)$$

Now replace $(1 + (R/z)^2)^{-1/2} \approx (1 - (1/2)(R/z)^2)$, and notice the $1/z$ terms cancel:

$$\mathbf{E} = \frac{\sigma \hat{z}}{2\epsilon_0} z \left(\frac{1}{2z} \left(\frac{R}{z} \right)^2 \right) \quad (10)$$

Multiply top and bottom by π , and recall that $Q = \pi R^2$ to find the point-charge field:

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{Q \hat{z}}{z^2} \quad (11)$$