


# INORGANIC CHEMISTRY

ENTHUSIAST | LEADER | ACHIEVER



**STUDY MATERIAL**

Hydrogen & It's Compounds

ENGLISH MEDIUM

## Copyright Statement

All rights including trademark and copyrights and rights of translation etc. reserved and vested exclusively with ALLEN Career Institute Private Limited. (ALLEN)

No part of this work may be copied, reproduced, adapted, abridged or translated, transcribed, transmitted, stored or distributed in any form retrieval system, computer system, photographic or other system or transmitted in any form or by any means whether electronic, magnetic, chemical or manual, mechanical, digital, optical, photocopying, recording or otherwise, or stored in any retrieval system of any nature without the written permission of the Allen Career Institute Private Limited. Any breach will entail legal action and prosecution without further notice.

This work is sold/distributed by Allen Career Institute Private Limited subject to the condition and undertaking given by the student that all proprietary rights (under the Trademark Act, 1999 and Copyright Act, 1957) of the work shall be exclusively belong to ALLEN Career Institute Private Limited. Neither the Study Materials and/or Test Series and/or the contents nor any part thereof i.e. work shall be reproduced, modify, re-publish, sub-license, upload on website, broadcast, post, transmit, disseminate, distribute, sell in market, stored in a retrieval system or transmitted in any form or by any means for reproducing or making multiple copies of it.

Any person who does any unauthorised act in relation to this work may be liable to criminal prosecution and civil claims for damages. Any violation or infringement of the propriety rights of Allen shall be punishable under Section- 29 & 52 of the Trademark Act, 1999 and under Section- 51, 58 & 63 of the Copyright Act, 1957 and any other Act applicable in India. All disputes are subjected to the exclusive jurisdiction of courts, tribunals and forums at Kota, Rajasthan only.

**Note:-** This publication is meant for educational and learning purposes. All reasonable care and diligence have been taken while editing and printing this publication. ALLEN Career Institute Private Limited shall not hold any responsibility for any error that may have inadvertently crept in. ALLEN Career Institute Private Limited is not responsible for the consequences of any action taken on the basis of this publication.

## HYDROGEN AND ITS COMPOUNDS

### 4.0 INTRODUCTION :

Hydrogen is the lightest element and also the lightest gas in the periodic table.

#### Order of Abundance of H :

Hydrogen is the most abundant element in the universe (70% of the total mass of the universe.)

**Order of abundance of H :** Universe > Sun > Earth

The planets Jupiter and Saturn consist mainly of H<sub>2</sub>. Similarly about half the mass of the sun and some other stars is made up of hydrogen.

In Sun's atmosphere and in universe, it is found in atomic form. While in earth it is generally found in molecular form. At Sun, the stratosphere is made up of H (atomic hydrogen) and they undergoes fusion and converted into He nuclei and this reaction is exothermic.

It is the ninth element on earth in order of abundance.

Earth does not possess enough gravitational force to retain live hydrogen molecule i.e. why it is not found in earth atmosphere in atomic form.

Hydrogen is the most reactive element in atomic form but it is less reactive in molecular form because of very high bond dissociation energy due to 1s-1s overlapping.

### 4.1 ISOTOPIC EFFECT :


The effect which can change the physical and chemical properties of isotopes is called isotopic effect.

It is because of difference in mass.

In isotopic effect maximum changes occurs in physical properties like melting point, boiling point, bond energy, while minimum changes occurs in chemical properties like state of chemical reaction etc.

**Imp.** Isotopic effect is found only in hydrogen isotopes. Because there is a large difference in mass.

**Ex.** Which of the following reaction is fast and why ?





**Ans.** (i) because C-H bond energy is less in comparison to C-D bond energy.


### 4.2 METHOD OF PREPARATION :

#### (a) From acids :

The metal which are placed above H in electrochemical series react with dil acids to liberate H<sub>2</sub>.




**Lab preparation :** When impure Zn reacts with dil. H<sub>2</sub>SO<sub>4</sub> it forms H<sub>2</sub>




**Ex.** Why we use impure Zn.

**Ans.** Because the rate of reaction with pure Zn is very slow.


- (b) **By alkalis :** Only (Be, Zn, Al, Sn, Pb, Si) (Amphoteric metal) react with boiling NaOH or KOH to evolve  $H_2$ .




(sodium zincate)




(sodium meta aluminate)




(sodium stannate)



(sodium plumbate)



(sodium silicate)



(sodium beryllate)

- (c) **With water :**

(i) With cold water ( $7^\circ\text{C}$ – $25^\circ\text{C}$ ) : Li, K, Ba, Ca, Na, Sr


(ii) With hot water ( $25^\circ\text{C}$ – $90^\circ\text{C}$ ) : Mg, Al, Cr, Mn, Zn

(iii) With steam (greater than  $100^\circ\text{C}$ ) : Fe, Cd, Ni, Sn, Pb

- (d) **Commercial or industrial method to prepare  $H_2$  :**



The commonly used processes are outlined below:

- (i) Electrolysis of acidified water using platinum electrodes gives hydrogen.





- (ii) High purity ( $>99.95\%$ ) dihydrogen is obtained by electrolysis of warm aqueous barium hydroxide solution between nickel electrodes.

- (iii) It is obtained as a by product in the manufacture of sodium hydroxide and chlorine by the electrolysis of brine solution. During electrolysis, the reactions that take place are:




- (iv) **From hydrocarbons :** Reaction of steam on hydrocarbons or coke at high temperatures in the presence of catalyst yields hydrogen.




The mixture of CO and  $H_2$  is called *water gas*. As this mixture of CO and  $H_2$  is used for the synthesis of methanol and a number of hydrocarbons, it is also called *synthesis gas* or '**syngas**'. Now days 'syngas' is produced from sewage, saw-dust, scrap wood, newspapers etc. The process of producing 'syngas' from coal is called '*coal gasification*'.

- (v) **Bosch process :**




The production of dihydrogen can be increased by reacting carbon monoxide of syngas mixtures with steam in the presence of iron chromate as catalyst.



This is called *water-gas shift reaction*.

- (vi) **Lane's process :**



### 4.3 PHYSICAL PROPERTIES OF HYDROGEN :

- Hydrogen is the lightest, colorless, odourless and tasteless gas. It is sparingly soluble in water. It is inflammable and less reactive gas.
- Its m.p. ( $-259.2^{\circ}\text{C}$ ) and b.p. ( $-252^{\circ}\text{C}$ ) are very low indicating less intermolecular attraction. Due to low m.p. liquid hydrogen is used as a **cryogenic fluid** (to produce low temperature).
- H—H bond energy is  $104 \text{ Kcal mol}^{-1}$  or  $436 \text{ KJ/mol}$
- H—H bond length is  $74 \text{ pm}$  so  $\text{H}_2$  is less reactive and require high temp for reaction.

### 4.4 USES OF HYDROGEN :

- Hydrogenation of vegetable oil to form solid fats i.e. vanaspati ghee.
- In liquid form as a rocket fuel. (Liquid  $\text{H}_2$  + Liquid  $\text{O}_2$ )
- In a air ship of balloons as a mixture of Hydrogen & Helium [ $15\% \text{H}_2$  +  $85\% \text{He}$ ]
- Formation of different compounds.

Like  $\rightarrow \text{NH}_3$ , (Haber process) alkane, alcohol and other hydrocarbons

### BEGINNER'S BOX-1

- Which of the following produces hydrolith with dihydrogen ?  
 (1) Mg (2) Al (3) Cu (4) Ca
- Hydrogen combines with other elements by :-  
 (1) Losing an electron (2) Gaining an electron  
 (3) Sharing an electron (4) Losing, gaining or sharing electron
- Hydrogen readily combines with non-metals and thus it shows its :-  
 (1) Electronegativity character (2) Electropositive character  
 (3) Both (1) and (2) (4) None of these
- The oxidation states shown by hydrogen are :-  
 (1)  $-1$  only (2) Zero only (3)  $+1, -1, 0$  (4)  $+1$  only
- Hydrogen readily combines with metals and thus shows its :-  
 (1) Electropositive character (2) Electronegative character  
 (3) Both (1) and (2) (4) None of these
- In which of the compounds does hydrogen have an oxidation state of  $-1$  :-  
 (1)  $\text{CH}_4$  (2)  $\text{NH}_3$  (3)  $\text{HCl}$  (4)  $\text{CaH}_2$
- Match List I (Fuels) with List II (composition) and select the correct answer using the codes given below the lists :-

#### List I (Fuels)

- Water gas
- Producer gas
- Coal gas
- Natural gas

	A	B	C	D
(1)	iii	i	iv	ii
(2)	iii	i	ii	iv
(3)	i	iii	iv	ii
(4)	iii	ii	iv	i

#### List II (Composition)

- A mixture of CO and  $\text{N}_2$
- Methane
- A mixture of CO and  $\text{H}_2$
- A mixture of CO,  $\text{H}_2$ ,  $\text{CH}_4$  and  $\text{CO}_2$

- $\text{H}_2$  react with halogen in presence of catalyst to form hydrogen halide HX. X (Halogen) is :-  
 (1)  $\text{Cl}_2$  (2)  $\text{Br}_2$  (3)  $\text{F}_2$  (4)  $\text{I}_2$
- Which of the following is a method of preparation of  $\text{H}_2$  ?  
 (1) reaction of granulated Zn with dil. HCl (2) reaction of Zn with aqueous Alkali.  
 (3) by electrolysis of Brine solution (4) All of the above

## 4.5 HYDRIDES

The binary compounds of hydrogen with different elements are called hydrides.

These are of three types :


### (a) Ionic/Salt like/Saline hydrides :

- ◆ Compounds of hydrogen with s-block elements except beryllium & magnesium are called ionic hydrides.




$\text{BeH}_2, \text{MgH}_2$  are covalent polymeric hydride.

- ◆ Structure of these hydrides are similar to rock salt, so they are also called salt like/saline hydrides.
- ◆ Down the group, atomic size  $\uparrow$  Lattice energy  $\downarrow$  stability  $\downarrow$  Melting point  $\downarrow$  Boiling point  $\downarrow$
- ◆ On electrolysis of these hydrides, hydrogen is liberated at anode.
- ◆ On reaction with water these hydrides will form hydrogen



- ◆ These hydrides form complex hydrides which are very good reducing agents.





$\text{LiAlH}_4 \longrightarrow$  Lithium aluminium hydride

$\text{NaBH}_4 \longrightarrow$  Sodium borohydride

### (b) Metallic / Interstitial hydrides :

- ◆ They are the compounds of d & f-block elements. In these hydrides hydrogen occupies interstitial sites present in metallic lattice, so they are called interstitial hydrides.
- ◆ Properties of these hydrides are similar to parent metals, so they are also known as metallic hydrides.
- ◆ These hydrides are non stoichiometric in nature (i.e. having variable composition)



- ◆ Metals of group 7,8,9 don't form any hydrides so this particular part of periodic table is known as hydride gap.
- ◆ In group-6 only one hydride  $\text{CrH}$  is formed.

### (c) Covalent/Molecular hydrides

- ◆ They are the compounds of hydrogen with p-block elements  $\text{CH}_4, \text{NH}_3, \text{H}_2\text{O}, \text{HF}$ , etc.
- ◆ These hydrides exist as molecules, so they are also known as molecular hydrides. These hydrides are non-conductor of electricity.

These hydrides are again divides into 3 categories.

#### (a) Electron deficient hydrides :

They are the hydrides of group 13 elements.

e.g.  $\text{BH}_3, \text{AlH}_3, \text{GaH}_3$  – In these hydrides central element does not have complete octet. i.e. why they are called electron deficient compounds.

#### (b) Electron precise hydrides :

They are the hydrides of group 14 element.

e.g.  $\text{CH}_4, \text{SiH}_4, \text{GeH}_4$  – In these type of hydrides central element has  $8e^-$  in its outer most shell.

#### (c) Electron rich hydrides :


These are the hydrides of group 15, 16 and 17

e.g.  $\text{NH}_3, \text{H}_2\text{O}, \text{HF}$  – In these hydrides lone pair are present on central element which can be given to others. So they are called electron rich hydrides.






- (iii) **Permutit process** : Permutit is hydrated sodium alumino silicate




Permutit is also known as sodium zeolite ( $\text{Na}_2\text{Z}$ ) means Zeolite is  $\text{Al}_2\text{Si}_2\text{O}_8 \cdot x\text{H}_2\text{O}$ .

In this process when hard water is poured into chamber, it may contain organic impurities like plant. These impurities can be removed by gravel. On moving upwards hard water will react with  $\text{Na}_2\text{Z}$  during this reaction  $\text{Na}^+$  ions of  $\text{Na}_2\text{Z}$  will replace  $\text{Mg}^{+2}$  &  $\text{Ca}^{+2}$  of impurities.



$\text{NaCl}$  is dissolved in water & water becomes soft.

After some time when  $\text{Na}_2\text{Z}$  is completely converted into  $\text{CaZ}$ . Process is stopped and for regeneration of  $\text{Na}_2\text{Z}$ . We use  $\text{NaCl}$  solution.






- (iv) **Ion exchange resin** : This process removes both temporary and permanent hardness. Also, by this process we can remove both cation & anion of hardness.


This process contain two chambers.

- Cation exchange resin : This resin contains granular insoluble organic acid having giant molecules with  $\text{RCOOH}^+$  group.
- Anion exchange resin : This resin contain giant organic molecules with basic groups derived from amines with  $\text{RNH}_3^+\text{OH}^-$

**Process** : When hard water is poured into first chamber the cation of hardness ( $\text{Mg}^{+2}, \text{Ca}^{+2}$ ) removed by  $\text{H}^+$  ions of organic acid.



This water becomes soften but not used for drinking purpose because this water contain the impurity of acid. To remove anion of hardness this acidic water then passed through another bed containing anion exchanger. This exchanger removes anion like  $\text{Cl}^-$ ,  $\text{SO}_4^{2-}$  &  $\text{HCO}_3^-$ .



This water is free from impurities & can be used for drinking purpose.


After sometime when both resin gets exhausted process is stopped.

#### Regeneration of resin :

- Cation exchange resin : We use dil acid.  

$$2\text{H}^+\text{Cl}^- + (\text{RCOO})_2\text{Ca} \longrightarrow 2\text{RCOOH}^+ + \text{CaCl}_2$$
- Anion exchange resin : We use dil  $\text{NaOH}$  solution  

$$\text{RNH}_3^+\text{Cl}^- + \text{Na}^+\text{OH}^- \longrightarrow \text{Na}^+\text{Cl}^- + \text{RNH}_3^+\text{OH}^-$$







#### 4.7 HEAVY WATER (D<sub>2</sub>O)

##### Method of preparation :

**Repeated electrolysis of H<sub>2</sub>O :** On electrolysis of water (impure) H<sub>2</sub>O dissociate into H<sup>+</sup> & OH<sup>-</sup> while a fractional part of D<sub>2</sub>O will dissociate into D<sup>+</sup> & OD<sup>-</sup>



D<sup>+</sup> & OD<sup>-</sup> due to more mass have less mobility i.e. why H<sup>+</sup> & OH<sup>-</sup> will move towards cathode & anode respectively while D<sup>+</sup> & OD<sup>-</sup> will be in solution.

This process is repeated six times.

##### Properties of Heavy water :

##### Physical properties :

- Heavy water is a colourless, odourless and tasteless mobile liquid.
- Nearly all the physical constants are higher than the corresponding values of ordinary water.  
(dielectric constant order H<sub>2</sub>O > D<sub>2</sub>O).

##### Chemical properties :

- Heavy water is chemically similar to ordinary water. However, D<sub>2</sub>O reacts more slowly than H<sub>2</sub>O in chemical reactions.
- All chemical properties are same, but reactions are slow.

**Uses : As a neutron moderator :** Fission in uranium-235 is brought by slow speed neutrons. The substances which are used for slowing down the speed of neutrons are called moderators. Heavy water is used for this purpose in nuclear reactors.


#### 4.8 H<sub>2</sub>O<sub>2</sub> (HYDROGEN PEROXIDE)

##### Method of preparation (According to NCERT)





- Acidifying barium peroxide and removing excess water by evaporation under reduced pressure gives hydrogen peroxide.



- Industrial Method : Auto oxidation of 2 ethyl anthraquinol ( cyclic process ) :





- Electrolytic Process : (Used 50%) H<sub>2</sub>SO<sub>4</sub> in electrolytic cell using Pt as anode and graphite as cathode.




**Physical properties :**

- (i) Pure  $\text{H}_2\text{O}_2$  is colourless, odourless liquid and impure with bluish layer.
- (ii) It has more Hydrogen bonding than  $\text{H}_2\text{O}$ . So, order of boiling point is  $\text{H}_2\text{O}_2 > \text{D}_2\text{O} > \text{H}_2\text{O}$   
 Order of melting point =  $\left[ \begin{matrix} \text{H}_2\text{O}_2 < \text{H}_2\text{O} \\ -0.35^\circ\text{C} & 0^\circ\text{C} \end{matrix} \right]$
- (iii) It is soluble in  $\text{H}_2\text{O}$ , alcohol and ether.
- (iv) It has bitter taste and harmful for skin.
- (v) It is a dibasic weak acid.
- (vi) It has oxidising as well as reducing property.
- (vii)  $\text{H}_2\text{O}_2$  easily decompose in presence of light and temperature, so  $\text{H}_2\text{O}_2$  always kept in dark bottles and kept at cool places and small amount of inhibitor is added like R-OH, glycerol and Acetone.
- (viii) 30% solution of  $\text{H}_2\text{O}_2$  is called **Perhydrol**.
- (ix)  $\text{H}_2\text{O}_2$  decomposes at its boiling point hence its distillation is carried out under reduced pressure.

**Chemical properties :**
**(A) Oxidising and Reducing behaviour of  $\text{H}_2\text{O}_2$** 

**Structure of hydrogen peroxide**


(a) Gas phase






(a) Solid phase

**Uses**

- |  |  |
|--|--|
| (i) Bleaching agent  | (ii) Hair dying                                      |
| (iii) $\text{H}_2\text{O}_2 + \text{N}_2\text{H}_4$ as Rocket propellant | (iv) $\text{H}_2\text{O}_2$ as oxidant and reductant |
| (v) Antiseptic   |  |

## BEGINNER'S BOX-2

1. The structure of  $\text{H}_2\text{O}_2$  is
- (1)  (2)  (3)  $\text{H}-\text{O}-\text{O}-\text{H}$  (4) 
2. Temporary hardness may be removed from water by adding
- (1)  $\text{CaCO}_3$  (2)  $\text{Ca}(\text{OH})_2$  (3)  $\text{CaSO}_4$  (4)  $\text{HCl}$
3. Temporary hardness of water can be removed by
- (1) Addition of potassium permanganate (2) Boiling  
(3) Filtration (4) Addition of chlorine
4. When zeolite (Hydrated sodium aluminium silicate) is treated with hard water the sodium ions are exchanged with
- (1)  $\text{OH}^-$  ions (2)  $\text{SO}_4^{2-}$  ions (3)  $\text{Ca}^{2+}$  ions (4)  $\text{H}^+$  ions
5. When temporary hard water containing  $\text{Mg}(\text{HCO}_3)_2$  is boiled the ppt. formed is of
- (1)  $\text{MgCO}_3$  (2)  $\text{MgO}$  (3)  $\text{Mg}(\text{OH})_2$  (4) None of these
6. Which of the following can effectively remove all types of hardness of water
- (1) Soap (2) Washing soda (3) Slaked lime (4) None of these
7. Impurity of  $\text{Mg}(\text{HCO}_3)_2$  is removed from water on boiling. It removes in the form of
- (1)  $\text{MgCO}_3$  (2)  $\text{Mg}(\text{OH})_2$  (3)  $\text{Mg}(\text{HCO}_3)_2$  (4) None of these
8. In cation exchange process for permanent hardness,  $\text{H}^+$  exchange for  $\text{Na}^+$ ,  $\text{Mg}^{+2}$ ,  $\text{Ca}^{+2}$  and other cation present in water. This process result in proton release and thus makes the nature of water :-
- (1) Acidic (2) Basic (3) Neutral (4) Amphoteric
9. Which of the following option is correct ?  
[Statement T for true & F for false]
- (a)  $\text{H}_2\text{O}_2$  is almost colourless (very pale blue) liquid  
(b)  $\text{H}_2\text{O}_2$  is miscible with water in all proportion and forms hydrated  $\text{H}_2\text{O}_2 \cdot \text{H}_2\text{O}$   
(c) 1%  $\text{H}_2\text{O}_2$  is concentrated to 30% by distillation under reduced pressure  
(d) 1 ml of 30 %  $\text{H}_2\text{O}_2$  solution will give 100 ml of oxygen at STP
- (1) TTTT (2) FTTT (3) FTFT (4) FFTT
10. In a following chemical reaction
- $$\text{MnO}_4^- + \text{H}^+ + \text{H}_2\text{O}_2 \rightarrow$$
- The incorrect option is
- (1)  $\text{O}_2$  is release (2)  $\text{Mn}^{+7}$  reduced into  $\text{Mn}^{+2}$   
(3)  $\text{H}_2\text{O}_2$  acts as an oxidising agent (4) All of these

## ANSWER'S KEY

BEGINNER'S BOX-1	Que.	1	2	3	4	5	6	7	8	9	
	Ans.	4	4	2	3	2	4	1	4	4	
BEGINNER'S BOX-2	Que.	1	2	3	4	5	6	7	8	9	10
	Ans.	2	2	2	3	3	2	2	1	1	3