


PHYSICAL CHEMISTRY

ENTHUSIAST | LEADER | ACHIEVER



EXERCISE

Electrochemistry

ENGLISH MEDIUM

EXERCISE-I (Conceptual Questions)
Build Up Your Understanding
ELECTROLYTIC CONDUCTANCE

1. Strong electrolytes are those which :
- (1) dissolve readily in water
 - (2) conduct electricity
 - (3) dissociate into ions even at high concentration
 - (4) dissociate into ions at high dilution.

EC0001

2. Molten sodium chloride conducts electricity due to the presence of :
- (1) free electrons
 - (2) free ions
 - (3) free molecules
 - (4) free atoms of Na and Cl

EC0002

3. Electrolytic conduction is due to the movement of :
- (1) molecules
 - (2) atoms
 - (3) ions
 - (4) electrons

EC0003

4. Which of the following solutions of KCl has the lowest value of equivalent conductance ?
- (1) 1 M
 - (2) 0.1 M
 - (3) 0.01 M
 - (4) 0.001 M

EC0004

5. If the specific resistance of a solution of concentration $C \text{ g eq L}^{-1}$ is R , then its equivalent conductance is :

- (1) $\frac{100R}{C}$
- (2) $\frac{RC}{1000}$
- (3) $\frac{1000}{RC}$
- (4) $\frac{C}{1000R}$

EC0006

6. The specific conductances in $\text{ohm}^{-1} \text{ cm}^{-1}$ of four electrolytes P, Q, R and S are given in brackets :
- P (5.0×10^{-5}) Q (7.0×10^{-8})
 R (1.0×10^{-10}) S (9.2×10^{-3})
- The one that offers highest resistance to the passage of electric current is
- (1) P
 - (2) S
 - (3) R
 - (4) Q

EC0007

7. The specific conductance of a salt of 0.01 M concentration is $1.061 \times 10^{-4} \text{ S cm}^{-1}$. Molar conductance of the same solution will be :
- (1) $1.061 \times 10^{-4} \text{ S cm}^2 \text{ mol}^{-1}$
 - (2) $1.061 \text{ S cm}^2 \text{ mol}^{-1}$
 - (3) $10.61 \text{ S cm}^2 \text{ mol}^{-1}$
 - (4) $106.1 \text{ S cm}^2 \text{ mol}^{-1}$

EC0008

8. Which statement is not correct :-
- (1) Conductance of an electrolytic solution increases with dilution
 - (2) Conductance of an electrolytic solution decreases with dilution
 - (3) Specific conductance of an electrolytic solution decreases with dilution
 - (4) Equivalent conductance of an electrolytic solution increases with dilution.

EC0011

9. The resistance of 0.01 N solution of an electrolyte was found to be 210 ohm at 298 K using a conductivity cell of cell constant 0.66 cm^{-1} . The equivalent conductance of solution is :-

- (1) $314.28 \text{ mho cm}^2 \text{ eq}^{-1}$
- (2) $3.14 \text{ mho cm}^2 \text{ eq}^{-1}$
- (3) $314.28 \text{ mho}^{-1} \text{ cm}^2 \text{ eq}^{-1}$
- (4) $3.14 \text{ mho}^{-1} \text{ cm}^2 \text{ eq}^{-1}$

EC0012

10. Electrolytic conduction differs from metallic conduction from the fact that in the former

- (1) The resistance increases with increasing temperature
- (2) The resistance decreases with increasing temperature
- (3) The resistance remains constant with increasing temperature
- (4) The resistance is independent of the length of the conductor

EC0013

11. The specific conductance of a 0.01 M solution of KCl is $0.0014 \text{ ohm}^{-1} \text{ cm}^{-1}$ at 25° C . Its equivalent conductance ($\text{cm}^2 \text{ ohm}^{-1} \text{ eq}^{-1}$) is :-

- (1) 140
- (2) 14
- (3) 1.4
- (4) 0.14

EC0014

12. Which one of the following is wrong :-

- (1) Specific conductance increases on dilution.
- (2) Specific conductance decreases on dilution.
- (3) Equivalent conductance increases on dilution.
- (4) Molar conductance increases on dilution.

EC0015

KOHLRAUSCH LAW

13. At infinite dilution, the equivalent conductances of CH_3COONa , HCl and CH_3COOH are 91, 426 and $391 \text{ mho cm}^2 \text{ eq}^{-1}$ respectively at 25°C . The eq. conductance of NaCl at infinite dilution will be :

- (1) 126 (2) 209 (3) 391 (4) 908

EC0016

14. For HCl solution at 25°C , equivalent conductance at infinite dilution is $425 \text{ ohm}^{-1} \text{ cm}^2 \text{ eq}^{-1}$. The specific conductance of a solution of HCl is $3.825 \text{ ohm}^{-1} \text{ cm}^{-1}$. If the degree of dissociation is 90%, the normality of the solution is :-

- (1) 0.90 N (2) 1.0 N
(3) 10 N (4) 1.2 N

EC0019

15. The molar conductivities Λ_{NaOAc}^0 and Λ_{HCl}^0 at infinite dilution in water at 25°C are 91.0 and $426.2 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. To calculate Λ_{HOAc}^0 the additional value required is :

- (1) Λ_{NaCl}^0 (2) $\Lambda_{\text{H}_2\text{O}}^0$ (3) Λ_{KCl}^0 (4) Λ_{NaOH}^0

EC0020

16. The molar conductance of AgNO_3 , AgCl and NaCl at infinite dilution are 116.5, 121.6 and $110.3 \text{ S cm}^2 \text{ mol}^{-1}$ respectively. The molar conductance of NaNO_3 is :

- (1) $111.4 \text{ S cm}^2 \text{ mol}^{-1}$ (2) $105.2 \text{ S cm}^2 \text{ mol}^{-1}$
(3) $130.6 \text{ S cm}^2 \text{ mol}^{-1}$ (4) $150.2 \text{ S cm}^2 \text{ mol}^{-1}$

EC0021

17. The conductivity of a saturated solution of BaSO_4 is $3.06 \times 10^{-6} \text{ ohm}^{-1} \text{ cm}^{-1}$ and its molar conductance is $1.53 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$. The K_{sp} of BaSO_4 will be

- (1) 4×10^{-12} (2) 2.5×10^{-9}
(3) 2.5×10^{-13} (4) 4×10^{-6}

EC0022

18. Kohlrausch's law states that at :-

- (1) Infinite dilution, each ion makes definite contribution to conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
- (2) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
- (3) Finite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.

- (4) Infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte depending on the nature of the other ion of electrolyte.

EC0099**GALVANIC CELL**

19. In the galvanic cell



the electrons will travel in the external circuit :

- (1) from Ag to Cu
- (2) from Cu to Ag
- (3) electrons do not travel in the external circuit
- (4) in any direction

EC0023

20. The direction of current in the Daniell cell when Zn and Cu electrodes are connected is :

- (1) from Cu to Zn in the cell
- (2) from Cu to Zn out side the cell
- (3) from Zn to Cu outside the cell
- (4) in any direction in the cell

EC0024

21. The equation representing the process by which standard reduction potential of zinc can be defined is

- (1) $\text{Zn}^{2+} (\text{s}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$
- (2) $\text{Zn} (\text{g}) \longrightarrow \text{Zn}^{2+} (\text{g}) + 2\text{e}^-$
- (3) $\text{Zn}^{2+} (\text{g}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$
- (4) $\text{Zn}^{2+} (\text{aq.}) + 2\text{e}^- \longrightarrow \text{Zn} (\text{s})$

EC0025

22. A standard hydrogen electrode has zero electrode potential because :

- (1) Hydrogen is easiest to oxidize.
- (2) This electrode potential is assumed to be zero.
- (3) Hydrogen atom has only one electron.
- (4) Hydrogen is the lightest element.

EC0026

23. Which is not true for a standard hydrogen electrode ?

- (1) The hydrogen ion concentration is 1 M.
- (2) Temperature is 25°C .
- (3) Pressure of hydrogen is 1 bar.
- (4) It contains a metallic conductor which does not adsorb hydrogen.

EC0027

24. E° for the half cell

$\text{Zn}^{2+} | \text{Zn}$ is -0.76 V . E.M.F. of the cell
 $\text{Zn} | \text{Zn}^{2+} (1\text{M}) || 2\text{H}^+ (1\text{M}) | \text{H}_2 (1 \text{ atm})$ is :

- (1) -0.76 V (2) $+0.76 \text{ V}$
 (3) -0.38 V (4) $+0.38 \text{ V}$

EC0028

25. $\text{Cu(s)} | \text{Cu}^{+2} (1 \text{ M}) || \text{Zn}^{+2} (1 \text{ M}) | \text{Zn(s)}$

A cell represented above should have emf.

- (1) Positive
 (2) Negative
 (3) Zero
 (4) Cannot be predicted

EC0030

26. Given electrode potentials :

$\text{Fe}^{3+} + e \longrightarrow \text{Fe}^{2+}$; $E^\circ = 0.771 \text{ V}$

$\text{I}_2 + 2e \longrightarrow 2\text{I}^-$; $E^\circ = 0.536 \text{ V}$

E°_{cell} for the cell reaction

$2\text{Fe}^{3+} + 2\text{I}^- \rightarrow 2\text{Fe}^{2+} + \text{I}_2$ is -

- (1) $(2 \times 0.771 - 0.536) = 1.006 \text{ V}$
 (2) $(0.771 - 0.5 \times 0.536) = 0.503 \text{ V}$
 (3) $0.771 - 0.536 = 0.235 \text{ V}$
 (4) $0.536 - 0.771 = -0.235 \text{ V}$

EC0031

27. Which of the following is not an anodic half reaction:-

- (1) $\text{Ag}^+ \rightarrow \text{Ag} - e^-$
 (2) $\text{Cu} \rightarrow \text{Cu}^{2+} + 2e^-$
 (3) $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$
 (4) $4\text{OH}^- \rightarrow 2\text{H}_2\text{O} + \text{O}_2 + 4e^-$

EC0032

28. Which of the following statements is correct :-

- (1) Oxidation occurs at anode in both galvanic and electrolytic cell.
 (2) Reduction occurs at anode in both galvanic and electrolytic cell
 (3) Reduction occurs at anode in electrolytic cell where as oxidation occurs at cathode in galvanic cell
 (4) Oxidation occurs at anode in electrolytic cell where as reduction occurs at anode in a galvanic cell,

EC0033

29. Other things being equal, the life of a Daniell cell may be increased by :-

- (1) Keeping low temperature
 (2) Using large copper electrode
 (3) Decreasing concentration of copper ions
 (4) Using large zinc electrodes

EC0034

ELECTROCHEMICAL SERIES

30. Zn can not displace following ions from their aqueous solution :-

- (1) Ag^+ (2) Cu^{2+} (3) Fe^{2+} (4) Na^+

EC0035

31. The standard reduction potentials at 25°C for the following half reactions are given against each :-

$\text{Zn}^{2+} (\text{aq}) + 2e^- \rightleftharpoons \text{Zn(s)}, -0.762 \text{ V}$

$\text{Cr}^{3+} (\text{aq}) + 3e^- \rightleftharpoons \text{Cr(s)}, -0.740 \text{ V}$

$2\text{H}^+ + 2e^- \rightleftharpoons \text{H}_2(\text{g}), 0.00 \text{ V}$

$\text{Fe}^{3+} + 2e^- \rightleftharpoons \text{Fe}^{2+}, 0.77 \text{ V}$

Which is the strongest reducing agent ?

- (1) Zn (2) Cr
 (3) $\text{H}_2(\text{g})$ (4) $\text{Fe}^{2+} (\text{aq})$

EC0036

32. Using the standard electrode potential values given below, decide which of the statements, I, II, III and IV are correct. Choose the right answer from (1), (2), (3) and (4).

$\text{Fe}^{2+} + 2e^- \rightleftharpoons \text{Fe}$; $E^\circ = -0.44 \text{ V}$

$\text{Cu}^{2+} + 2e^- \rightleftharpoons \text{Cu}$; $E^\circ = +0.34 \text{ V}$

$\text{Ag}^+ + e^- \rightleftharpoons \text{Ag}$; $E^\circ = +0.80 \text{ V}$

I. Copper can displace iron from FeSO_4 solution.

II. Iron can displace copper from CuSO_4 solution.

III. Silver can displace copper from CuSO_4 solution.

IV. Iron can displace silver from AgNO_3 solution.

- (1) I and II (2) II and III
 (3) II and IV (4) I and IV

EC0041

33. The standard electrode potentials for the elements A, B and C are 0.68 , -2.50 and 0.50 V respectively. The order of their reducing power is :

- (1) $A > B > C$ (2) $A > C > B$
 (3) $C > B > A$ (4) $B > C > A$

EC0042

34. The oxidation potential of Zn, Cu, Ag, H_2 and Ni electrodes are 0.76 V , -0.34 V , -0.80 V , 0 V , 0.55 V respectively. Which of the following reaction will provide maximum voltage ?

- (1) $\text{Zn} + \text{Cu}^{2+} \longrightarrow \text{Cu} + \text{Zn}^{2+}$
 (2) $\text{Zn} + 2\text{Ag}^+ \longrightarrow 2\text{Ag} + \text{Zn}^{2+}$
 (3) $\text{H}_2 + \text{Cu}^{2+} \longrightarrow 2\text{H}^+ + \text{Cu}$
 (4) $\text{H}_2 + \text{Ni}^{2+} \longrightarrow 2\text{H}^+ + \text{Ni}$

EC0043

35. The standard reduction potential at 25 °C of Li^+ / Li , $\text{Ba}^{2+} / \text{Ba}$, Na^+ / Na and $\text{Mg}^{2+} / \text{Mg}$ are -3.05 V , -2.73 V , -2.71 V and -2.37 V respectively. Which one of the following is the strongest oxidising agent ?

(1) Na^+ (2) Li^+ (3) Ba^{2+} (4) Mg^{2+}

EC0045

36. A gas X at 1 atm is bubbled through a solution containing a mixture of 1 M Y^- and 1 M Z^- at 25 °C. If the reduction potential of $(\text{Z}/\text{Z}^-) > (\text{Y}/\text{Y}^-) > (\text{X}/\text{X}^-)$ then :

(1) Y will oxidise X^- and not Z^-
 (2) Y will oxidise Z^- and not X^-
 (3) Y will oxidise both X^- and Z^-
 (4) Y will reduce both X^- and Z^-

EC0046

37. The standard electrode potential of Zn, Ag and Cu electrodes are -0.76 V , 0.80 V and 0.34 V respectively, then :

(1) Ag can oxidise Zn and Cu
 (2) Ag can reduce Zn^{2+} and Cu^{2+}
 (3) Zn can reduce Ag^+ and Cu^{2+}
 (4) Cu can oxidise Zn and Ag

EC0047

38. Standard reduction potentials of four metal electrodes are

A = -0.250 V , B = -0.140 V
 C = -0.126 V , D = -0.402 V

The metal that displaces A from aqueous solution of its compounds is :-

(1) B (2) C
 (3) D (4) None of the above

EC0049

39. The following four colourless salt solutions are placed in separate test tubes and a strip of Cu is placed in each solution. Which solution finally turns blue :-

(1) $\text{Zn}(\text{NO}_3)_2$ (2) $\text{Mg}(\text{NO}_3)_2$
 (3) KNO_3 (4) AgNO_3

EC0050

40. Which of the following displacement does not occur

(1) $\text{Zn} + 2\text{H}^+ \rightarrow \text{Zn}^{2+} + \text{H}_2 \uparrow$
 (2) $\text{Fe} + 2\text{Ag}^+ \rightarrow \text{Fe}^{2+} + \text{Ag} \downarrow$
 (3) $\text{Cu} + \text{Fe}^{2+} \rightarrow \text{Cu}^{2+} + \text{Fe} \downarrow$
 (4) $\text{Zn} + \text{Pb}^{2+} \rightarrow \text{Zn}^{2+} + \text{Pb} \downarrow$

EC0051

41. On the basis of the following E° values, the strongest oxidizing agent is :-

$[\text{Fe}(\text{CN})_6]^{4-} \rightarrow [\text{Fe}(\text{CN})_6]^{3-} + e^-$, $E^\circ = -0.35 \text{ V}$

$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$; $E^\circ = -0.77 \text{ V}$

(1) Fe^{3+}
 (2) $[\text{Fe}(\text{CN})_6]^{3-}$
 (3) $[\text{Fe}(\text{CN})_6]^{4-}$
 (4) Fe^{2+}

EC0098

NERNST EQUATION

42. $E^\circ(\text{Ni}^{2+}/\text{Ni}) = -0.25 \text{ V}$

$E^\circ(\text{Au}^{3+}/\text{Au}) = 1.50 \text{ V}$

The emf of the voltaic cell.

$\text{Ni} | \text{Ni}^{2+} (1.0 \text{ M}) || \text{Au}^{3+} (1.0 \text{ M}) | \text{Au}$ is :

(1) 1.25 V (2) -1.75 V
 (3) 1.75 V (4) 4.0 V

EC0052

43. The emf of the cell

$\text{Ti(s)} | \text{Ti}^+ (0.0001 \text{ M}) || \text{Cu}^{2+} (0.01 \text{ M}) | \text{Cu(s)}$ is 0.83 V

The emf of this cell will be increased by :-

(1) Increasing the concentration of Cu^{+2} ions
 (2) Decreasing the concentration of Ti^+
 (3) Increasing the concentration of both
 (4) (1) & (2) both

EC0054

44. Which of the following represents the electrode potential of silver electrode dipped into 0.1 M AgNO_3 solution at 25°C ?

(1) E°_{red} (2) $(E^\circ_{\text{red}} + 0.059)$
 (3) $(E^\circ_{\text{ox}} - 0.059)$ (4) $(E^\circ_{\text{red}} - 0.059)$


EC0056

45. The electrode potential of a hydrogen electrode dipped in solution of $\text{pH} = 1$ is

(1) 0.059 V (2) 0.00 V
 (3) -0.059 V (4) 0.59 V

EC0057

46. Consider the reaction




The emf of the cell when

$[\text{Cl}^-] = [\text{Br}^-] = 0.01 \text{ M}$ and Cl_2 gas at 0.25 atm pressure will be (E° for the above reaction is $= 0.29 \text{ V}$)

(1) 0.54 V (2) 0.272 V
 (3) 0.29 V (4) -0.29 V

EC0058

47. The standard emf for the cell reaction



The emf for the cell reaction when 0.1 M Cu^{2+} and 0.1 M Zn^{2+} solution are used at 25°C is :

(1) 1.10 V (2) 0.110 V
 (3) -1.10 V (4) -0.110 V

EC0059

48. E° for $F_2 + 2e^- \rightarrow 2F^-$ is 2.8 V,
 E° for $\frac{1}{2}F_2 + e^- \rightarrow F^-$ is ?
 (1) 2.8 V (2) 1.4 V (3) -2.8 V (4) -1.4 V
EC0060
49. How much will the potential of $Zn | Zn^{2+}$ change if the solution of Zn^{2+} is diluted 10 times
 (1) increases by 0.03 V
 (2) decreases by 0.03 V
 (3) increases by 0.059 V
 (4) decreases by 0.059 V
EC0062
50. How much will the potential of a hydrogen electrode change when its solution initially at pH = 0 is neutralised to pH = 7 ?
 (1) increases by 0.059 V
 (2) decreases by 0.059 V
 (3) increases by 0.41 V
 (4) decreases by 0.41 V
EC0063
51. Which of the following will increase the voltage of the cell with following cell reaction
 $Sn(s) + 2Ag^+(aq) \rightarrow Sn^{2+}(aq) + 2Ag(s)$
 (1) Increase in the size of silver rod
 (2) Increase in the concentration of Sn^{2+} ions
 (3) Increase in the concentration of Ag^+ ions
 (4) Decrease in the concentration of Ag^+ ions
EC0064
52. E° for the reaction $Fe + Zn^{2+} \rightarrow Zn + Fe^{2+}$ is -0.35 V. The given cell reaction is :
 (1) spontaneous (2) non-spontaneous
 (3) in equilibrium (4) can't say anything
EC0065
53. For a reaction - $A(s) + 2B^+ \rightleftharpoons A^{2+} + 2B(s)$
 K_c has been found to be 10^{12} . The E°_{cell} is:
 (1) 0.354 V (2) 0.708 V
 (3) 0.0098 V (4) 1.36 V
EC0066
54. The standard electrode potential (E°) for $OC\bar{I}^- / Cl^-$ and $Cl^- / \frac{1}{2}Cl_2$ respectively are 0.94 V and -1.36 V. The E° value of $OC\bar{I}^- / \frac{1}{2}Cl_2$ will be :
 (1) -2.20 V (2) -0.42 V (3) 0.52 V (4) 1.04 V
EC0067
55. The hydrogen electrode is dipped in a solution of pH = 3 at 25 °C. The electrode potential of the half cell would be :
 (1) 0.177 V (2) -0.177 V
 (3) 0.087 V (4) 0.059 V
EC0069
56. What is the potential of the cell containing two hydrogen electrodes as represented below
 $Pt; H_2(g) | H^+(10^{-8}M) || H^+(0.001 M) | H_2(g).Pt$;
 (1) -0.295 V (2) -0.0591 V
 (3) 0.295 V (4) 0.0591 V
EC0070
57. Consider the cell $Cu | Cu^{+2} || Ag^+ | Ag$. If the concentration of Cu^{+2} and Ag^+ ions becomes ten times, then the emf of the cell will :-
 (1) Becomes 10 times
 (2) Remains same
 (3) Increases by 0.0295 V
 (4) Decreases by 0.0295 V
EC0071
58. The emf of the cell
 $Ni | Ni^{+2}(1.0 M) || Au^{+3}(0.1M) | Au$
 $[E^\circ \text{ for } Ni^{+2} | Ni = -0.25 V,$
 $E^\circ \text{ for } Au^{+3} | Au = 1.50 V]$ is given as:-
 (1) 1.25 V (2) -1.75 V
 (3) 1.78 V (4) 1.73 V
EC0072
59. The equilibrium constant (in approx) of the cell reaction :
 $Cu(s) + 2Ag^+(aq.) \rightleftharpoons Cu^{2+}(aq.) + 2Ag(s)$
 if $E^\circ_{cell} = 0.465 V$ at 298 K is :-
 (1) 2.0×10^{10} (2) 3.16×10^{12}
 (3) 3.16×10^{15} (4) 4×10^{10}
EC0097
60. The emf of the cell in which the following reaction
 $Zn(s) + Ni^{2+}(a = 0.1) \rightleftharpoons Zn^{2+}(a = 1.0) + Ni(s)$
 occurs, is found to be 0.5105 V at 298 K. The standard e.m.f. of the cell is :-
 (1) -0.5105 V (2) 0.5400 V
 (3) 0.4810 V (4) 0.5696 V
EC0053
61. If $E^\circ_{Fe^{+2}/Fe} = -0.441V$ and $E^\circ_{Fe^{+3}/Fe^{+2}} = 0.771V$ the standard EMF of the reaction
 $Fe + 2Fe^{+3} \rightarrow 3Fe^{+2}$ will be :
 (1) 0.330 V (2) 1.653 V
 (3) 1.212 V (4) 0.111 V
EC0093

ELECTROLYSIS

- 62.** When an electric current is passed through acidified water, 112 mL of hydrogen gas at STP collects at the cathode in 965 s. The current passed in ampere is :
 (1) 1.0 (2) 0.5
 (3) 0.1 (4) 2.0
EC0073
- 63.** Two electrolytic cells one containing acidified ferrous chloride and another acidified ferric chloride are connected in series. The ratio of iron deposited at cathodes in the two cells when electricity is passed through the cells will be :
 (1) 3 : 1 (2) 2 : 1 (3) 1 : 1 (4) 3 : 2
EC0074
- 64.** A current of 9.65 A flowing for 10 minute deposits 3.0 g of a metal. The equivalent weight of the metal is :
 (1) 10 (2) 30 (3) 50 (4) 96.5
EC0075
- 65.** How many coulombs of electric charge are required for the oxidation of 1 mol of H_2O to O_2 ?
 (1) $9.65 \times 10^4 \text{ C}$ (2) $4.825 \times 10^5 \text{ C}$
 (3) $1.93 \times 10^5 \text{ C}$ (4) $1.93 \times 10^4 \text{ C}$
EC0077
- 66.** On passing 10800 C through electrolytic solution, 2.977 g of metal (atomic mass 106.4 g mol^{-1}) was deposited, the charge on the metal cation is -
 (1) +4 (2) +3 (3) +2 (4) +1
EC0079
- 67.** On passing electricity through dilute H_2SO_4 solution the amount of substance liberated at the cathode and anode are in the ratio :
 (1) 1 : 8 (2) 8 : 1
 (3) 16 : 1 (4) 1 : 16
EC0080
- 68.** During electrolysis of fused calcium hydride, the hydrogen is produced at :
 (1) Cathode
 (2) Anode
 (3) Hydrogen is not liberated at all
 (4) H_2 produced reacts with oxygen to form water
EC0081
- 69.** A silver cup is plated with silver by passing 965 A current for one second, the mass of Ag deposited is :-(At. wt. of Ag = 107.87)
 (1) 9.89 g. (2) 107.87 g.
 (3) 1.0787 g. (4) 100.2 g.
EC0082
- 70.** When electricity is passed through molten AlCl_3 , 13.5 g Al is deposited. The number of Faradays must be :-
 (1) 5.0 (2) 1.0 (3) 1.5 (4) 3.0
EC0083
- 71.** A solution of sodium sulphate in water is electrolysed using inert electrodes. The product at the cathode and anode are respectively :-
 (1) H_2 , SO_2 (2) O_2 , H_2
 (3) O_2 , Na (4) H_2 , O_2
EC0084
- 72.** One Faraday of electricity will liberate one mole of the metal from the solution of
 (1) Auric chloride (AuCl_3) (2) Silver nitrate
 (3) Calcium chloride (4) Copper sulphate
EC0085
- 73.** When 96500 C of electricity are passed through molten barium chloride solution, the amount of barium deposited will be :-
 (1) 0.5 mol (2) 1.0 mol
 (3) 1.5 mol (4) 2.0 mol
EC0086
- 74.** A factory produces 40 kg of calcium in two hours by electrolysis. How much aluminium can be produced by the same current in two hours :-
 (1) 22 kg (2) 18 kg (3) 9 kg (4) 27 kg
EC0087
- 75.** What would be the ratio of moles of Ag, Cu, Fe that would be deposited by passage of same quantity of electricity through solutions of salts containing Ag^+ , Cu^{+2} , Fe^{+3} :-
 (1) 1 : 1 : 1 (2) $1 : \frac{1}{2} : \frac{1}{3}$
 (3) $\frac{1}{3} : \frac{1}{2} : 1$ (4) 1 : 2 : 3
EC0088
- 76.** Electrolysis of aq. CuSO_4 causes :-
 (1) An increase in pH
 (2) A decrease in pH
 (3) Either decrease or increase in pH
 (4) None
EC0089

77. The passage of current liberates H_2 at cathode and Cl_2 at anode. The solution is :-
 (1) $CuSO_4$ (aq) (2) $CuCl_2$ (aq.)
 (3) $NaCl$ (aq.) (4) Water

EC0090
COMMERCIAL CELLS

78. When lead accumulator is charged, it acts as :
 (1) an electrolytic cell (2) a galvanic cell
 (3) a Daniel cell (4) none of the above

EC0091

79. The thermodynamic efficiency of cell is given by -

- (1) $\frac{\Delta H}{\Delta G} \times 100$ (2) $\frac{nFE}{\Delta G} \times 100$
 (3) $-\frac{nFE}{\Delta H} \times 100$ (4) Zero

EC0169
EXERCISE-I (Conceptual Questions)
ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	3	2	3	1	3	3	3	2	1	2	1	1	1	3	1
Que.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	2	4	2	2	2	4	2	4	2	2	3	1	1	4	4
Que.	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
Ans.	1	3	4	2	4	1	3	3	4	3	1	3	4	4	3
Que.	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans.	2	1	1	1	4	3	2	1	3	2	3	3	4	3	2
Que.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75
Ans.	3	1	4	3	3	1	1	2	3	3	4	2	1	2	2
Que.	76	77	78	79											
Ans.	2	3	1	3											

EXERCISE-II (Previous Year Questions)

AIPMT/NEET

AIPMT 2009

1. Given :
 (i) $\text{Cu}^{2+} + 2e^- \rightarrow \text{Cu}$, $E^\circ = 0.337 \text{ V}$
 (ii) $\text{Cu}^{2+} + e^- \rightarrow \text{Cu}^+$, $E^\circ = 0.153 \text{ V}$
 Electrode potential, E° for the reaction, $\text{Cu}^+ + e^- \rightarrow \text{Cu}$, will be :-
 (1) 0.38 V (2) 0.52 V
 (3) 0.90 V (4) 0.30 V

EC0100

2. The equivalent conductance of $\frac{M}{32}$ solution of a weak monobasic acid is $8.0 \text{ mho cm}^2 \text{ eq}^{-1}$ and at infinite dilution is $400 \text{ mho cm}^2 \text{ eq}^{-1}$. The dissociation constant of this acid is :-
 (1) 1.25×10^{-4} (2) 1.25×10^{-5}
 (3) 1.25×10^{-6} (4) 6.25×10^{-4}

EC0101

3. Al_2O_3 is reduced by electrolysis at low potential and high current. If $4.0 \times 10^4 \text{ A}$ of current is passed through molten Al_2O_3 for 6 hours, what mass of aluminium is produced? (Assume 100% current efficiency, At. mass of Al = 27 g mol^{-1})
 (1) $1.3 \times 10^4 \text{ g}$ (2) $9.0 \times 10^3 \text{ g}$
 (3) $8.1 \times 10^4 \text{ g}$ (4) $2.4 \times 10^5 \text{ g}$

EC0102

AIPMT 2010

4. An increase in equivalent conductance of a strong electrolyte with dilution is mainly due to:-
 (1) Increase in number of ions.
 (2) Increase in ionic mobility of ions.
 (3) 100% ionisation of electrolyte at normal dilution.
 (4) Increase in both i.e. number of ions and ionic mobility of ions.

EC0103

5. Consider the following relations for emf of an electrochemical cell :
 (a) emf of cell = (Oxidation potential of anode) - (Reduction potential of cathode)
 (b) emf of cell = (Oxidation potential of anode) + (Reduction potential of cathode)
 (c) emf of cell = (Reduction potential of anode) + (Reduction potential of cathode)
 (d) emf of cell = (Oxidation potential of anode) - (Oxidation potential of cathode)

Which of the above relations are correct :

- (1) (a) and (b) (2) (c) and (d)
 (3) (b) and (d) (4) (c) and (a)

EC0104

6. Which of the following expressions correctly represents the equivalent conductance of $\text{Al}_2(\text{SO}_4)_3$ at infinite dilution. Given that $\Lambda^\circ_{\text{Al}^{3+}}$ and $\Lambda^\circ_{\text{SO}_4^{2-}}$ are the equivalent conductances at infinite dilution of the respective ions :-

- (1) $\Lambda^\circ_{\text{Al}^{3+}} + \Lambda^\circ_{\text{SO}_4^{2-}}$
 (2) $\left(\Lambda^\circ_{\text{Al}^{3+}} + \Lambda^\circ_{\text{SO}_4^{2-}} \right) \times 6$
 (3) $\frac{1}{3} \Lambda^\circ_{\text{Al}^{3+}} + \frac{1}{2} \Lambda^\circ_{\text{SO}_4^{2-}}$
 (4) $2\Lambda^\circ_{\text{Al}^{3+}} + 3\Lambda^\circ_{\text{SO}_4^{2-}}$

EC0105

7. For the reduction of silver ions with copper metals, the standard cell potential was found to be $+0.46 \text{ V}$ at 25°C . The value of standard Gibbs energy, ΔG° will be ($F = 96500 \text{ C mol}^{-1}$)
 (1) -98.0 kJ (2) -89.0 kJ
 (3) -89.0 J (4) -44.5 kJ



EC0106

AIPMT Pre. 2011

8. Standard electrode potential of three metals X, Y and Z are -1.2 V , $+0.5 \text{ V}$ and -3.0 V respectively. The reducing power of these metals will be :-
 (1) $Y > Z > X$ (2) $Y > X > Z$
 (3) $Z > X > Y$ (4) $X > Y > Z$

EC0109

9. The electrode potentials for



are $+0.15 \text{ V}$ and $+0.50 \text{ V}$ respectively. The value of $E^\circ_{\text{Cu}^{2+}/\text{Cu}}$ will be :-

- (1) 0.500 V (2) 0.325 V
 (3) 0.650 V (4) 0.150 V

EC0110

10. Standard electrode potential for $\text{Sn}^{4+}/\text{Sn}^{2+}$ couple is $+0.15 \text{ V}$ and that for the Cr^{3+}/Cr couples is -0.74 V . These two couples in their standard state are connected to make a cell. The standard cell potential will be :-

- (1) $+1.19 \text{ V}$ (2) $+0.89 \text{ V}$
 (3) $+0.18 \text{ V}$ (4) $+1.83 \text{ V}$

EC0111

11. If the E°_{cell} for a given reaction has a negative value, then which of the following gives the correct relationship for the values of ΔG° and K_{eq} ?

- (1) $\Delta G^\circ > 0$; $K_{\text{eq}} > 1$ (2) $\Delta G^\circ < 0$; $K_{\text{eq}} > 1$
(3) $\Delta G^\circ < 0$; $K_{\text{eq}} < 1$ (4) $\Delta G^\circ > 0$; $K_{\text{eq}} < 1$

EC0112

AIPMT Mains 2011

12. A solution contains Fe^{2+} , Fe^{3+} and I^- ions. This solution was treated with iodine at 35 °C. E° for $\text{Fe}^{3+} | \text{Fe}^{2+}$ is +0.77 V and E° for $\text{I}_2 | 2\text{I}^- = 0.536$ V. The favourable redox reaction is :-

- (1) Fe^{2+} will be oxidised to Fe^{3+}
(2) I_2 will be reduced to I^-
(3) There will be no redox reaction
(4) I^- will be oxidised to I_2

EC0113

AIPMT Pre. 2012

13. Limiting molar conductivity of NH_4OH

(i.e. $\Lambda_m^\circ(\text{NH}_4\text{OH})$) is equal to:-

- (1) $\Lambda_m^\circ(\text{NH}_4\text{OH}) + \Lambda_m^\circ(\text{NH}_4\text{Cl}) - \Lambda_m^\circ(\text{HCl})$
(2) $\Lambda_m^\circ(\text{NH}_4\text{Cl}) + \Lambda_m^\circ(\text{NaOH}) - \Lambda_m^\circ(\text{NaCl})$
(3) $\Lambda_m^\circ(\text{NH}_4\text{Cl}) + \Lambda_m^\circ(\text{NaCl}) - \Lambda_m^\circ(\text{NaOH})$
(4) $\Lambda_m^\circ(\text{NaOH}) + \Lambda_m^\circ(\text{NaCl}) - \Lambda_m^\circ(\text{NH}_4\text{Cl})$

EC0114

AIPMT Mains 2012

14. Molar conductivities (Λ_m°) at infinite dilution of NaCl , HCl and CH_3COONa are 126.4, 425.9 and 91.0 $\text{S cm}^2 \text{mol}^{-1}$ respectively. Λ_m° for CH_3COOH will be :-

- (1) 290.8 $\text{S cm}^2 \text{mol}^{-1}$
(2) 390.5 $\text{S cm}^2 \text{mol}^{-1}$
(3) 425.5 $\text{S cm}^2 \text{mol}^{-1}$
(4) 180.5 $\text{S cm}^2 \text{mol}^{-1}$

EC0115

NEET-UG 2013

15. At 25 °C molar conductance of 0.1 molar aqueous solution of ammonium hydroxide is 9.54 $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$ and at infinite dilution its molar conductance is 238 $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$. The degree of ionisation of ammonium hydroxide at the same concentration and temperature is :-

- (1) 40.800 % (2) 2.080 %
(3) 20.800 % (4) 4.008 %


EC0117

16. A hydrogen gas electrode is made by dipping platinum wire in a solution of HCl of pH = 10 and by passing hydrogen gas around the platinum wire at 1 atm pressure. The oxidation potential of electrode would be ?



- (1) 1.18 V (2) 0.059 V
(3) 0.59 V (4) 0.118 V

EC0118

17. A button cell used in watches function as following



If half cell potentials are



$$E^\circ = 0.34 \text{ V}$$

The standard cell potential will be :-

- (1) 1.34 V (2) 1.10 V
(3) 0.42 V (4) 0.84 V

EC0119

AIPMT 2014

18. When 0.1 mol MnO_4^{2-} is oxidised the quantity of electricity required to completely oxidise MnO_4^{2-} to MnO_4^- is :-

- (1) 96500 C (2) 2×96500 C
(3) 9650 C (4) 96.50 C

EC0121

19. The weight of silver (at wt. = 108) displaced by a quantity of electricity which displaces 5600 mL of O_2 at STP will be :-

- (1) 5.4 g (2) 10.8 g
(3) 54.0 g (4) 108.0 g

EC0122

AIPMT 2015

20. A device that converts energy of combustion of fuels like hydrogen and methane, directly into electrical energy is known as :-

- (1) Electrolytic cell (2) Dynamo
(3) Ni-Cd cell (4) Fuel Cell

EC0124

NEET-I 2016

21. The pressure of H_2 required to make the potential of H_2 -electrode zero in pure water at 298 K is :-

- (1) 10^{-14} atm (2) 10^{-12} atm
(3) 10^{-10} atm (4) 10^{-4} atm

EC0127

NEET-II 2016

22. The molar conductivity of a 0.5 mol/dm^3 solution of AgNO_3 with electrolytic conductivity of $5.76 \times 10^{-3} \text{ S cm}^{-1}$ at 298 K is

- (1) $0.086 \text{ S cm}^2/\text{mol}$ (2) $28.8 \text{ S cm}^2/\text{mol}$
(3) $2.88 \text{ S cm}^2/\text{mol}$ (4) $11.52 \text{ S cm}^2/\text{mol}$

EC0128

23. During the electrolysis of molten sodium chloride, the time required to produce 0.10 mol of chlorine gas using a current of 3 A is

- (1) 220 minutes (2) 330 minutes
(3) 55 minutes (4) 110 minutes

EC0129

24. If the E°_{cell} for a given reaction has a negative value, which of the following gives the **correct** relationships for the values of ΔG° and K_{eq} ?

- (1) $\Delta G^\circ < 0$; $K_{eq} > 1$ (2) $\Delta G^\circ < 0$; $K_{eq} < 1$
(3) $\Delta G^\circ > 0$; $K_{eq} < 1$ (4) $\Delta G^\circ > 0$; $K_{eq} > 1$

EC0130

25. The number of electrons delivered at the cathode during electrolysis by a current of 1 A in 60 s is (charge on electron = 1.60×10^{-19} C)

- (1) 3.75×10^{20} (2) 7.48×10^{23}
(3) 6×10^{23} (4) 6×10^{20}

EC0131

NEET(UG) 2017

26. In the electrochemical cell :-

Zn | ZnSO₄(0.01M) || CuSO₄(1.0 M) | Cu, the emf of this Daniel cell is E₁. When the concentration of ZnSO₄ is changed to 1.0M and that of CuSO₄ changed to 0.01M, the emf changes to E₂.

Which one of the relationship is correct between E_1 and E_2 ?


(Given, $\frac{RT}{F} = 0.059$)

- (1) $E_1 < E_2$ (2) $E_1 > E_2$
(3) $E_2 = 0 \neq E_1$ (4) $E_1 = E_2$

EC0133

NEET(UG) 2018

27. Consider the change in oxidation state of Bromine corresponding to different EMF values as shown in the diagram below:



Then the species undergoing disproportionation is:-

- (1) BrO_3^- (2) BrO_4^-
(3) Br_2 (4) HBrO

EC0135

NEET(UG) 2019


28. For a cell involving one electron $E^\circ_{\text{cell}} = 0.59\text{V}$ at 298 K, the equilibrium constant for the cell reaction is :-

$$\left[\text{Given that } \frac{2.303RT}{F} = 0.059V \text{ at } T = 298K \right]$$

- (1) 1.0×10^2 (2) 1.0×10^5
(3) 1.0×10^{10} (4) 1.0×10^{30}

EC0192

29. For the cell reaction



$E_{\text{cell}}^{\circ} = 0.24\text{V}$ at 298 K. The standard Gibbs energy ($\Delta_r G^{\circ}$) of the cell reaction is :

[Given that Faraday constant $F = 96500 \text{ C mol}^{-1}$]

- (1) $-46.32 \text{ kJ mol}^{-1}$ (2) $-23.16 \text{ kJ mol}^{-1}$
(3) $46.32 \text{ kJ mol}^{-1}$ (4) $23.16 \text{ kJ mol}^{-1}$

EC0193

NEET(UG) (Odisha) 2019

- 30.** Following limiting molar conductivities are given as

$$\lambda_{m(H_2SO_4)}^0 = x \text{ S cm}^2 \text{ mol}^{-1}$$

$$\lambda_{m(K_2SO_4)}^0 = y \text{ S cm}^2 \text{ mol}^{-1}$$

$$\lambda_{m(CH_3COOK)}^0 = z \text{ S cm}^2 \text{ mol}^{-1}$$

λ_m^0 (in $\text{S cm}^2 \text{ mol}^{-1}$) for CH_3COOH will be-

- (1) $x - y + 2z$ (2) $x + y - z$
(3) $x - y + z$ (4) $\frac{(x - y)}{2} + z$

EC0194

- 31.** The standard electrode potential (E°) values of $\text{Al}^{3+}|\text{Al}$, $\text{Ag}^+|\text{Ag}$, $\text{K}^+|\text{K}$ and $\text{Cr}^{3+}|\text{Cr}$ are -1.66 V , 0.80 V , -2.93 V and -0.74 V respectively. The correct decreasing order of reducing power of the metal is :

- (1) $\text{Ag} > \text{Cr} > \text{Al} > \text{K}$ (2) $\text{K} > \text{Al} > \text{Cr} > \text{Ag}$
(3) $\text{K} > \text{Al} > \text{Ag} > \text{Cr}$ (4) $\text{Al} > \text{K} > \text{Ag} > \text{Cr}$

EC0195

NEET (UG) 2020

- 32.** The number of Faradays(F) required to produce 20 g of calcium from molten CaCl_2 (Atomic mass of $\text{Ca} = 40 \text{ g mol}^{-1}$) is :

- (1) 4 (2) 1 (3) 2 (4) 3

EC0205

- 33.** On electrolysis of dil. sulphuric acid using Platinum (Pt) electrode, the product obtained at anode will be:

- (1) SO_2 gas (2) Hydrogen gas
(3) Oxygen gas (4) H_2S gas

EC0206

NEET (UG) 2020 (COVID-19)

- 34.** Identify the reaction from following having top position in EMF series (Std. red. potential) according to their electrode potential at 298 K.

- (1) $\text{Mg}^{2+} + 2e^- \rightarrow \text{Mg(s)}$ (2) $\text{Fe}^{2+} + 2e^- \rightarrow \text{Fe(s)}$
(3) $\text{Au}^{3+} + 3e^- \rightarrow \text{Au(s)}$ (4) $\text{K}^+ + 1e^- \rightarrow \text{K(s)}$

EC0207

- 35.** In a typical fuel cell, the reactants (R) and product (P) are :-

- (1) $\text{R} = \text{H}_{2(g)}$, $\text{O}_{2(g)}$; $\text{P} = \text{H}_2\text{O}_{2(l)}$
(2) $\text{R} = \text{H}_{2(g)}$, $\text{O}_{2(g)}$; $\text{P} = \text{H}_2\text{O}_{(l)}$
(3) $\text{R} = \text{H}_{2(g)}$, $\text{O}_{2(g)}$, $\text{Cl}_{2(g)}$; $\text{P} = \text{HClO}_{4(aq)}$
(4) $\text{R} = \text{H}_{2(g)}$, $\text{N}_{2(g)}$; $\text{P} = \text{NH}_{3(aq)}$

EC0208

NEET (UG) 2021

- 36.** The molar conductance of NaCl , HCl and CH_3COONa at infinite dilution are 126.45, 426.16 and 91.0 $\text{S cm}^2 \text{ mol}^{-1}$ respectively. The molar conductance of CH_3COOH at infinite dilution is.

Choose the right option for your answer.

- (1) 201.28 $\text{S cm}^2 \text{ mol}^{-1}$
(2) 390.71 $\text{S cm}^2 \text{ mol}^{-1}$
(3) 698.28 $\text{S cm}^2 \text{ mol}^{-1}$
(4) 540.48 $\text{S cm}^2 \text{ mol}^{-1}$

EC0209

- 37.** The molar conductivity of 0.007 M acetic acid is 20 $\text{S cm}^2 \text{ mol}^{-1}$. What is the dissociation constant of acetic acid ? Choose the correct option.

$$\left[\begin{array}{l} \Lambda_{H^+}^\circ = 350 \text{ S cm}^2 \text{ mol}^{-1} \\ \Lambda_{CH_3COO^-}^\circ = 50 \text{ S cm}^2 \text{ mol}^{-1} \end{array} \right]$$

- (1) $1.75 \times 10^{-4} \text{ mol L}^{-1}$
(2) $2.50 \times 10^{-4} \text{ mol L}^{-1}$
(3) $1.75 \times 10^{-5} \text{ mol L}^{-1}$
(4) $2.50 \times 10^{-5} \text{ mol L}^{-1}$

EC0210

NEET (UG) 2021 (Paper-2)

- 38.** An electrolytic cell contains a solution of Ag_2SO_4 and has platinum electrodes. A current is passed until 1.6g of O_2 has been liberated at anode. The amount of silver deposited at cathode will be

- (1) 108 g (2) 1.6 g
(3) 0.8 g (4) 21.60 g

EC0211

- 39.** The specific conductance of a saturated solution of silver bromide is $k \text{ S cm}^{-1}$. The limiting ionic conductivity of Ag^+ and Br^- ions are x and y , respectively. The solubility of silver bromide in g L^{-1} is : (Molar mass of $\text{AgBr} = 188$)

- (1) $\frac{k \times 1000}{x - y}$ (2) $\frac{k}{x + y} \times 188$
(3) $\frac{k \times 1000 \times 188}{x + y}$ (4) $\frac{x + y}{k} \times \frac{1000}{188}$

EC0212

NEET (UG) 2022


40. At 298 K, the standard electrode potentials of Cu^{2+}/Cu , Zn^{2+}/Zn , Fe^{2+}/Fe and Ag^+/Ag are 0.34V, - 0.76 V, - 0.44 V and 0.80 V, respectively.

On the basis of standard electrode potential, predict which of the following reaction can not occur ?


- (1) $\text{CuSO}_4(\text{aq}) + \text{Fe}(\text{s}) \rightarrow \text{FeSO}_4(\text{aq}) + \text{Cu}(\text{s})$
- (2) $\text{FeSO}_4(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Fe}(\text{s})$
- (3) $2\text{CuSO}_4(\text{aq}) + 2\text{Ag}(\text{s}) \rightarrow 2\text{Cu}(\text{s}) + \text{Ag}_2\text{SO}_4(\text{aq})$
- (4) $\text{CuSO}_4(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Cu}(\text{s})$

EC0213

41. Given below are half cell reactions :



$$E_{\text{Mn}^{2+}/\text{MnO}_4^-}^\circ = -1.510\text{V}$$




$$E_{\text{O}_2/\text{H}_2\text{O}}^\circ = +1.223\text{V}$$

Will the permanganate ion, MnO_4^- liberate O_2 from water in the presence of an acid ?

- (1) No, because $E_{\text{cell}}^\circ = -0.287\text{V}$
- (2) Yes, because $E_{\text{cell}}^\circ = +2.733\text{V}$
- (3) No, because $E_{\text{cell}}^\circ = -2.733\text{V}$
- (4) Yes, because $E_{\text{cell}}^\circ = +0.287\text{V}$

EC0214

42. Find the emf of the cell in which the following reaction takes place at 298 K



(Given that $E_{\text{cell}}^\circ = 10.5\text{V}$, $\frac{2.303RT}{F} = 0.059$ at 298 K)




- (1) 1.385 V
- (2) 0.9615 V
- (3) 1.05 V
- (4) 1.0385 V

EC0215

NEET (UG) 2022 (OVERSEAS)





43. The three cells with their $E_{\text{(cell)}}^\circ$ values are given below :

Cells $E_{\text{(cell)}}^\circ / \text{V}$



The standard Gibbs free energy change values for three cells are, respectively

(F represents charge on 1 mole of electrons)



EC0216



44. Λ_m° for NaCl, HCl and CH_3COONa are 126.4, 425.9 and $91.05\text{ S cm}^2 \text{ mol}^{-1}$ respectively. If conductivity of $0.001028\text{ mol L}^{-1}$ acetic acid solution is $4.95 \times 10^{-5}\text{ S cm}^{-1}$, find the degree of dissociation of the acetic acid solution

- (1) 1.00
- (2) 0.1233
- (3) 1.233
- (4) 0.01233

EC0217

Re-NEET (UG) 2022

45. Two half cell reactions are given below :




The standard EMF of a cell with feasible redox reaction will be :

- (1) +7.09 V
- (2) +0.15 V
- (3) +3.47 V
- (4) -3.47 V

EC0218

46. Standard electrode potential for the cell with cell reaction



is 1.1 V. Calculate the standard gibbs energy change for the cell reaction.

(Given $F = 96487\text{ C mol}^{-1}$)

- (1) -200.27 kJ mol^{-1}
- (2) -212.27 kJ mol^{-1}
- (3) -212.27 J mol^{-1}
- (4) -200.27 J mol^{-1}

EC0219

EXERCISE-II (Previous Year Questions)

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	2	2	3	2	3	1	2	3	2	2	4	4	2	2	4
Que.	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Ans.	3	2	3	4	4	1	4	4	3	1	2	4	3	1	4
Que.	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
Ans.	2	2	3	3	2	2	3	4	3	3	4	B	3	2	3
Que.	46														
Ans.	2														

EXERCISE-III (Analytical Questions)

1. Specific conductance of 0.1 M Nitric acid is $6.3 \times 10^{-2} \text{ ohm}^{-1} \text{ cm}^{-1}$. The molar conductance of the solution is :
- (1) $630 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 - (2) $315 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 - (3) $6.300 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$
 - (4) $63.0 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$

EC0140

2. The highest electrical conductivity of the following aqueous solution is of
- (1) 0.1 M fluoroacetic acid
 - (2) 0.1 M difluoroacetic acid
 - (3) 0.1 M acetic acid
 - (4) 0.1 M chloroacetic acid

EC0141

3. Resistance of a conductivity cell filled with a solution of an electrolyte of concentration 0.1 M is 100Ω . The conductivity of this solution is 1.29 S m^{-1} . Resistance of the same cell when filled with 0.02 M solution of the same electrolyte is 520Ω . The molar conductivity of 0.02 M solution of the electrolyte will be.
- (1) $124 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 - (2) $1240 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 - (3) $1.24 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 - (4) $12.4 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$

EC0142

4. Resistance of 0.2 M solution of an electrolyte is 50Ω . The specific conductance of the solution is 1.3 S m^{-1} . If resistance of the 0.4 M solution of the same electrolyte is 260Ω then its molar conductivity is:-
- (1) $6250 \text{ S m}^2 \text{ mol}^{-1}$
 - (2) $6.25 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 - (3) $625 \times 10^{-4} \text{ S m}^2 \text{ mol}^{-1}$
 - (4) $62.5 \text{ S m}^2 \text{ mol}^{-1}$

EC0144

5. The equivalent conductance of NaCl at concentration C and at infinite dilution are λ_c and λ^∞ , respectively. The correct relationship between λ_c and λ^∞ is given by :
- (where the constant B is positive)
- (1) $\lambda_c = \lambda^\infty - (B) \sqrt{C}$
 - (2) $\lambda_c = \lambda^\infty + (B) \sqrt{C}$
 - (3) $\lambda_c = \lambda^\infty + (B) C$
 - (4) $\lambda_c = \lambda^\infty - (B) C$

EC0145
Master Your Understanding

6. Which of the following statements is TRUE for the electrochemical Daniel cell :
- (1) Electrons flow from copper electrode to zinc electrode.
 - (2) Current flows from zinc electrode to copper electrode.
 - (3) Cations move toward copper electrode.
 - (4) Cations move toward zinc electrode.

EC0147

7. Standard reduction electrode potential of three metal electrodes A, B and C are respectively + 0.5 V, - 3.0 V and -1.2 V. The reducing power of these metals are :
- (1) $C > B > A$
 - (2) $A > C > B$
 - (3) $B > C > A$
 - (4) $A > B > C$


EC0149

8. Consider the following E^0 values

$$E^0_{\text{Fe}^{3+}/\text{Fe}^{2+}} = +0.77 \text{ V}$$

$$E^0_{\text{Sn}^{2+}/\text{Sn}} = -0.14 \text{ V}$$

Under standard conditions the potential for the reaction



- (1) 0.91 V
- (2) 1.40 V
- (3) 1.68 V
- (4) 0.63 V

EC0150

9. For the redox reaction :
- $$\text{Zn(s)} + \text{Cu}^{2+} (0.1\text{M}) \rightarrow \text{Zn}^{2+} (1\text{M}) + \text{Cu(s)}$$
- taking place in a cell, E°_{cell} is 1.10 V. E_{cell} for the cell will be
- $$\left(2.303 \frac{RT}{F} = 0.0591 \right)$$
- (1) 1.07 V
 - (2) 0.82 V
 - (3) 2.14 V
 - (4) 1.80 V

EC0151

10. In a cell that utilises the reaction
- $$\text{Zn(s)} + 2\text{H}^+(\text{aq.}) \rightleftharpoons \text{Zn}^{2+}(\text{aq}) + \text{H}_2(\text{g})$$
- addition of H_2SO_4 to cathode compartment, will
- (1) increase the E_{cell} and shift equilibrium to the right
 - (2) lower the E_{cell} and shift equilibrium to the right
 - (3) lower the E_{cell} and shift equilibrium to the left
 - (4) increase the E_{cell} and shift equilibrium to the left

EC0152

11. The e.m.f. of a Daniell cell at 298 K is E_1 .



(0.01M) (1.0M)

When the concentration of ZnSO_4 is 1.0 M and that of CuSO_4 is 0.01 M, the e.m.f. changed to E_2 . What is the relationship between E_1 and E_2

- (1) $E_1 > E_2$ (2) $E_1 < E_2$
(3) $E_1 = E_2$ (4) $E_2 = 0 \neq E_1$

EC0153

12. The reduction potential of hydrogen half-cell will be negative if :-

- (1) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$
(2) $p(\text{H}_2) = 2 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$
(3) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 2.0 \text{ M}$
(4) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 1.0 \text{ M}$

EC0154

13. For a spontaneous reaction; ΔG , equilibrium constant (K) and E_{Cell}° will be respectively

- (1) -ve, < 1, -ve (2) -ve, > 1, -ve
(3) -ve, > 1, +ve (4) +ve, > 1, -ve

EC0156

14. The metal that cannot be obtained by electrolysis of an aqueous solution of its salts is:

- (1) Cu (2) Cr (3) Ag (4) Ca


EC0160

15. During electrolysis of a solution of AgNO_3 9650 C of charge is passed through the electroplating bath. The mass of silver deposited on the cathode will be :

- (1) 21.6 g (2) 108 g (3) 1.08 g (4) 10.8 g

EC0161

16. Aluminium oxide may be electrolysed at 1000 °C to furnish aluminium metal . The cathode reaction is




The amount of charge required to prepare 5.12 kg of aluminium metal by this method would be :-

- (1) $5.49 \times 10^4 \text{ C}$
(2) $5.49 \times 10^1 \text{ C}$
(3) $5.49 \times 10^7 \text{ C}$
(4) $1.83 \times 10^7 \text{ C}$

EC0162

17. A concentration cell involving a metal M is :




For this cell ;

- (1) $E_{\text{cell}} > 0$; $\Delta G < 0$
(2) $E_{\text{cell}} = 0$; $\Delta G = 0$
(3) $E_{\text{cell}} > 0$; $\Delta G > 0$
(4) $E_{\text{cell}}^\circ > 0$; $\Delta G^\circ < 0$

EC0196

18. $\text{M}^{+n}(\text{aq}) + \text{ne}^- \longrightarrow \text{M}(\text{s})$; $E_{\text{M}^{+n}/\text{M}}^\circ = xV$



if $x > y$ then :

- (I) $\text{M}^{+n}(\text{aq})$ is better oxidising agent compared to $\text{N}^{+n}(\text{aq})$
(II) $\text{N}(\text{s})$ is better reducing agent compared to $\text{M}(\text{s})$
(III) $\text{N}^{+n}(\text{aq})$ is better oxidising agent compared to $\text{M}^{+n}(\text{aq})$
(IV) $\text{M}(\text{s})$ is better reducing agent compared to $\text{N}(\text{s})$

- (1) I and III (2) II, III
(3) I and II (4) III, IV

EC0197

19. Which of the following is incorrect for



- (1) $E_{\text{cell}}^\circ = E_{\text{Cu}/\text{Cu}^{+2}}^\circ + E_{\text{Ag}^{+}/\text{Ag}}^\circ$
(2) $E_{\text{cell}}^\circ = E_{\text{Ag}^{+}/\text{Ag}}^\circ - E_{\text{Cu}^{+2}/\text{Cu}}^\circ$
(3) $E_{\text{cell}}^\circ = E_{\text{Cu}/\text{Cu}^{+2}}^\circ - E_{\text{Ag}^{+}/\text{Ag}}^\circ$
(4) $E_{\text{cell}}^\circ = E_{\text{Ag}/\text{Ag}^{+}}^\circ - E_{\text{Cu}^{+2}/\text{Cu}}^\circ$

EC0198

20. Find the incorrect formula

- (1) $E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{2.303 RT}{nF} \log_{10} Q$
(2) $\Delta G = \Delta G^\circ + RT \ln Q$
(3) $K_{\text{eq}} = 10^{\frac{nE_{\text{cell}}^\circ}{0.0591}}$
(4) At equilibrium $(\Delta G^\circ)_{\text{T,P}} = 0$

EC0199

21. $\Lambda_m^\circ \text{NaCl} = z \text{ S cm}^2 \text{ mol}^{-1}$
 $\Lambda_m^\circ \text{CH}_3\text{COONa} = y \text{ S cm}^2 \text{ mol}^{-1}$
 $\Lambda_m^\circ \text{HCl} = x \text{ S cm}^2 \text{ mol}^{-1}$
 $\Lambda_m^\circ \text{CH}_3\text{COOH} = w \text{ S cm}^2 \text{ mol}^{-1}$
 When its concentration is 0.1 M then pick the correct option :

- (1) $\Lambda_m^\circ \text{CH}_3\text{COOH} = x + y + z$
 (2) $\alpha_{\text{CH}_3\text{COOH}} = \frac{w}{x + y - z}$
 (3) $K_a = 0.1 \left[\frac{w}{x + y - z} \right]$ if $\alpha \ll 1$
 (4) $\Lambda_m^\circ \text{CH}_3\text{COOH} = \frac{\Lambda_{\text{eq}}^\circ \text{CH}_3\text{COOH}}{2}$

EC0200

22. The emf of the cell : $\text{Pt(s)} \mid \text{Br}^- (0.01\text{M}) \mid \text{Br}_2(\text{l}) \parallel \text{H}^+ (0.001 \text{ M}) \mid \text{H}_2(\text{g}) \mid \text{Pt(s)}$ will be

[Given : $E_{\text{Br}_2/\text{Br}^-}^\circ = 1.09 \text{ V}$ and $E_{\text{H}^+/\text{H}_2}^\circ = 0\text{V}$]

- (1) -0.79 V (2) -1.39 V
 (3) -1.30 V (4) -1.90 V

EC0201

23. Given $E_{\text{K}^+/\text{K}}^\circ = -2.93 \text{ V}$, $E_{\text{Ag}^+/\text{Ag}}^\circ = 0.80 \text{ V}$,
 $E_{\text{Hg}^{+2}/\text{Hg}}^\circ = 0.79 \text{ V}$, $E_{\text{Mg}^{+2}/\text{Mg}}^\circ = -2.37 \text{ V}$,
 $E_{\text{Cr}^{+3}/\text{Cr}}^\circ = -0.74 \text{ V}$. Arrange these metals in increasing order of reducing power

- (1) K, Mg, Cr, Hg, Ag (2) Ag, Hg, Cr, Mg, K
 (3) Mg, K, Cr, Ag, Hg (4) Ag, Hg, K, Cr, Mg

EC0202

24. 4.5 gm of aluminium metal is deposited at cathode from Al^{+3} ions by a certain quantity of electric charge. Find volume of $\text{H}_{2(\text{g})}$, produced at S.T.P. from H^+ ions by the same quantity of electric charge?

- (1) 5.6 litre (2) 11.2 litre
 (3) 22.4 litre (4) 44.8 litre

EC0203

25. The molar conductivity of 0.02 M HCOOH is $40.4 \text{ S cm}^2 \text{ mol}^{-1}$, If $\Lambda_m^\circ(\text{H}^+) = 350 \text{ S cm}^2 \text{ mol}^{-1}$ and $\Lambda_m^\circ(\text{HCOO}^-) = 54 \text{ S cm}^2 \text{ mol}^{-1}$ then dissociation constant of HCOOH is –

- (1) 2.22×10^{-3} (2) 2.22×10^{-4}
 (3) 2.22×10^{-5} (4) 1.22×10^{-2}

EC0204

EXERCISE-III (Analytical Questions)

ANSWER KEY

Que.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	1	2	1	2	1	3	3	1	1	1	1	1	3	4	4
Que.	16	17	18	19	20	21	22	23	24	25					
Ans.	3	1	3	4	4	2	2	2	1	2					