

LEADER & ACHIEVER COURSE

PHASE : MLB, MLK, MLM, MLP, MLQ, MLR, MAZE, MAZF, MAZG, MAZH, MAZM, MAZP, MAZQ,

MAZR, MAZU, MAZV, MAAZ, MAAW, SPARK PRO

TARGET : PRE MEDICAL 2025

Test Type : MAJOR

Test Pattern : NEET (UG)

TEST DATE : 25-03-2025

ANSWER KEY


Q.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
A.	3	1	1	2	1	2	1	3	4	3	1	3	4	1	1	1	1	2	4	1	4	3	2	3	1	1	3	1	2	3
Q.	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
A.	1	3	1	3	1	3	4	1	3	4	4	3	4	1	4	1	4	3	4	3	3	4	3	1	3	3	2	4	3	2
Q.	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90
A.	4	2	2	3	1	3	2	3	1	1	1	1	3	1	4	4	1	4	1	1	1	4	1	1	2	1	3	2	3	3
Q.	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
A.	3	4	1	3	1	4	3	4	2	3	4	3	3	4	1	1	4	3	2	2	2	3	3	2	3	2	1	2	2	2
Q.	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150
A.	3	1	2	2	2	4	3	1	3	3	2	4	4	1	3	3	3	4	4	4	3	1	2	4	4	3	4	4	3	4
Q.	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
A.	3	2	2	4	3	4	1	1	3	1	4	3	1	4	2	3	2	1	2	2	1	1	4	4	3	2	1	2	1	4

HINT - SHEET

1. Ans (3)

Simple series parallel combination start from 'AFE' side and combined.

2. Ans (1)




Apply KCL,

$$\frac{X-30}{4} + \frac{X-20}{4} + \frac{X-0}{2} = 0 \Rightarrow 4x = 50$$

$$x = 25/2 \text{ V} \Rightarrow I = \frac{25/2 - 0}{2} \Rightarrow I = \frac{25}{4} \text{ A}$$

3. Ans (1)




Current will flow from A to B.

4. Ans (2)

$$\text{T.P.D.} = \frac{5 \times 4 - 5 \times 1}{1 + 4} = 3\text{V}$$

5. Ans (1)



$$R = 1 + 1 + \frac{4R}{4 + R}$$

$$R = \frac{2(4 + R) + 4R}{4 + R} = \frac{8 + 6R}{6 + R}$$

$$4R + R^2 = 8 + 6R$$


$$R^2 - 4R + 2R - 8 = 0$$

$$R(R-4) + 2(R-4) = 0$$


$$R = 4\Omega$$

$$R = -2\Omega \text{ (Not possible)}$$

6. Ans (2)



Re-Draw



$$R_{AB} = \frac{4R}{3}$$

7. Ans (1)


$$R \propto \ell^2$$

$$\frac{R}{R^1} = \frac{\ell^2}{(2\ell)^2} = \frac{1}{4}$$

$$R^1 = 4R.$$

$$\% \text{ change} = \left(\frac{4R - R}{R} \right) 100\% = 300\%$$

8. Ans (3)



$$R_{eq} = \frac{1100}{5} = 220$$

$$I = \frac{220}{2200} = 1A$$

9. Ans (4)

$$I_g = \frac{2}{2000} = 10^{-3} A$$

$$V = I_g (G + H)$$

$$10 = 10^{-3} (2000 + H)$$

$$H = 10^4 - 2000 = 8000 \Omega$$


10. Ans (3)

$$P_c = \frac{V_A^2}{R} = \frac{V_A^2}{V_A^2} P_R$$


$$= \left(\frac{160}{200} \right)^2 100$$

$$= \frac{16}{25} \times 100 = 64 \text{ w}$$

11. Ans (1)



Re-Draw



$$R_{AB} = \frac{3R}{2}$$

12. Ans (3)

$$I = neAV_d$$

$$0.21 = 8.4 \times 10^{22} \times 10^6 \times 1.6 \times 10^{-19} \times 10^{-6} V_d$$

$$V_d = \frac{0.21}{8.4 \times 1.6 \times 10^3} \text{ m/s}$$

$$= 1.56 \times 10^{-5} \text{ m/s}$$


13. Ans (4)

$$I_1 : I_2 : I_3 = \frac{1}{60} : \frac{1}{20} : \frac{1}{10}$$

$$= 1 : 3 : 6$$

$$I_1 = \left(\frac{1}{10} \right) 20 = 2A$$

14. Ans (1)




$$R_{eq} = R_1 + R_2$$

$$\frac{\rho_{eq}(\ell_1 + \ell_2)}{A} = \frac{\rho_1 \ell_1}{A} + \frac{\rho_2 \ell_2}{A}$$


$$\rho_{eq} = \frac{\rho_1 \ell_1 + \rho_2 \ell_2}{\ell_1 + \ell_2}$$

15. Ans (1)



$$R_{AB} = \frac{3R}{8}$$


16. Ans (1)



$$V_A - 8 + 3 - 4 - 5 + 8 = V_B$$

$$V_A - V_B = 6V$$


17. Ans (1)



18. Ans (2)

$$P_C = \frac{(150)^2}{R/4} = \frac{4(150)^2}{(150)^2/100} = 400 \text{ W}$$

19. Ans (4)



$$V_X = \frac{V}{3}$$

$$V_Y = \frac{2V}{3}$$

$$V_Z = V$$

20. Ans (1)

$$R_{eq} = \frac{R}{5} = \frac{50/5}{5} = 2\Omega$$


21. Ans (4)

$$q_A = 12 \text{ nC}, q_B = -6 \text{ nC}, q_C = 0$$

$$q_A^1 = q_C^1 = \left(\frac{12\text{nC} + 0}{2} \right) = 6 \text{ nC}$$

$$q_C^{11} = q_B^1 = \frac{6\text{nC} - 6\text{nC}}{2} = 0 \text{ C}$$

22. Ans (3)



$$F' = F'' = F = \text{equal in magnitude}$$


$$F_{net} = 0$$

23. Ans (2)

$$q = ne$$

$$n = \frac{2}{1.6 \times 10^{-19}} = 1.25 \times 10^{19}$$

24. Ans (3)



25. Ans (1)

$$\text{Force in AIR } F = \frac{1}{4\pi\epsilon_0} \frac{2\mu \cdot 3\mu}{r^2}$$


$$\text{force in medium } F_m = \frac{1}{4\pi\epsilon_0} \frac{2\mu \cdot 3\mu}{r^2}$$

$$= \frac{\epsilon_0}{\epsilon} \left(\frac{1}{4\pi\epsilon} \frac{2\mu \cdot 3\mu}{r^2} \right)$$

$$= \frac{\epsilon_0}{\epsilon} \left(\frac{1}{4\pi\epsilon_0} \frac{2\mu \cdot 3\mu}{r^2} \right)$$

$$= \frac{\epsilon_0}{\epsilon} F$$

26. Ans (1)



Ball (3) must be neutral

27. Ans (3)

For (A)

$$E_1 = 9 \times 10^9 \times \frac{10^{-8}}{(0.05)^2} = 3.6 \times 10^4 = E_2$$

$$E_A = E_1 + E_2 = 7.2 \times 10^4 \text{ N/C}$$

For (C)

$$E_C = 2E_1 \cos \left(\frac{120}{2} \right)$$

$$= 2 \times 9 \times 10^9 \times \frac{10^{-8}}{(0.1)^2} \times \frac{1}{2} = 9 \times 10^3$$

$$\frac{E_A}{E_C} = \frac{7.2 \times 10}{0.9 \times 10} = \frac{8}{1}$$

28. Ans (1)

Electric potential is a scalar quantity and net potential at any point is algebraic sum of individual potential.

29. Ans (2)

Net flux through closed surface is depend on inside charge only, while electric field on closed surface depend on both inside and outside charge.

30. Ans (3)

By frictional method of charging, charge acquire by bodies are equal in magnitude and opposite in nature.

31. Ans (1)

$$\vec{F}_x = (F_{12} + F_{13} \sin \theta) \hat{i}$$

$$= k \left(\frac{q_1 q_2}{b^2} + \frac{q_1 q_3}{a^2} \sin \theta \right) \hat{i}$$

$$\Rightarrow F_y = F_{13} \cos \theta (-\hat{j})$$

$$= -k \frac{q_1 q_3}{a^2} \cos \theta \hat{j}$$


32. Ans (3)

$$V_A = \frac{1}{4\pi\epsilon_0} \left(\frac{q}{a} + \frac{q}{\sqrt{2}a} + \frac{q}{a} \right) = \frac{q}{4\pi\epsilon_0} \frac{1}{a} \left(2 + \frac{1}{\sqrt{2}} \right)$$

$$V_0 = \frac{1}{4\pi\epsilon_0} \left(\frac{3q}{a/\sqrt{2}} \right) = \frac{1}{4\pi\epsilon_0} \frac{3\sqrt{2}}{a}$$

$$\frac{V_A}{V_0} = \frac{(2\sqrt{2} + 1)}{6}$$

33. Ans (1)



34. Ans (3)

Workdone against the electric field.

$$\begin{aligned} W &= -q \int \vec{E} \cdot d\vec{l} \\ &= -q \int \vec{E} \cdot \hat{i} \cdot (dx\hat{i} + dy\hat{j}) \\ &= -qE(x_2 - x_1) \\ &= -10^{-6} \times 10^3 \times (3 - 1) \\ &= -2 \times 10^{-3} \text{ J} \end{aligned}$$

35. Ans (1)

$$\begin{aligned} \vec{E}_x &= -\frac{dv}{dx} \hat{i} = -(6x + 4)\hat{i} \\ \left(\vec{E}_x\right)_{\text{origin}} &= -4\hat{i} \end{aligned}$$

Force on e^-

$$\begin{aligned} \vec{F} &= q\vec{E} \\ &= (-e)(-4\hat{i}) \\ &= 4e\hat{i} \end{aligned}$$

36. Ans (3)

$$W = q_0(V_B - V_A)$$

$$V_B = V_A = k \frac{q}{a}$$


37. Ans (4)

$$\begin{aligned} E_x &= -\frac{dV}{dx} = (-5) = 5 \\ E_y &= -\frac{dV}{dy} = -3 \\ E_z &= -\frac{dV}{dz} = -\sqrt{15} \\ E &= \sqrt{(5)^2 + (-3)^2 + (-\sqrt{15})^2} \\ &= 7 \end{aligned}$$

39. Ans (3)


$$\begin{aligned} W_{A \rightarrow B} &= q_0(V_B - V_A) \\ &= 4(5 - (-10)) = 60 \text{ J} \end{aligned}$$

41. Ans (4)



$$C_{eq} = \frac{C}{3} + C = \frac{4C}{3}$$

42. Ans (3)



$$V_A - V_D = \frac{V}{2} \Rightarrow V_A = \frac{V}{2}$$

$$V_D - V_B = \frac{V}{2} \Rightarrow V_B = -\frac{V}{2}$$

44. Ans (1)

$$C_{eq} = \frac{110}{3} \mu\text{F}$$

$$C = \frac{Q}{V} \Rightarrow Q = C.V.$$

$$= \frac{110}{3} \mu\text{F} \times 50$$

$$= \frac{5500}{3} \mu\text{C}$$

45. Ans (4)

$$\begin{aligned} U &= \frac{1}{2} \epsilon_0 E^2 = \frac{1}{2} \epsilon_0 \left(\frac{V}{r} \right)^2 \\ &= 2.83 \text{ J/m}^3 \end{aligned}$$

51. Ans (3)

In acidic $\rightarrow$ Colourless

In Basic $\rightarrow$ Pink

52. Ans (4)

Number of g eq of HCl neutralized = $1 \times 1 \times 1 = 1$

Number of g eq of H_2SO_4 neutralized = $1 \times 2 \times 1 = 2$

$$2x = y$$

54. Ans (1)

$$m = \frac{\text{Moles of solute}}{\text{Weight of solvent (g)}} \times 1000$$

$$1 = \frac{0.5}{\text{Weight of solvent (g)}} \times 1000$$

Weight of solvent (g) = 500 g

55. Ans (3)

$X_{\text{H}_2\text{O}} = 0.02$

$\therefore X_{\text{gas}} = 0.98$

$P_{\text{total}} = 1.2 \text{ atm}$

partial pressure of dry-air = $P_T \times \text{mole fraction of dry-air}$

Partial pressure of dry-air = $1.2 \text{ atm} \times 0.98 = 1.176 \text{ atm}$.

56. Ans (3)

$$\frac{P^0 - P_s}{P^0} = i X_{\text{solute}}$$

$$= 2 \times 0.1 = 0.2$$

57. Ans (2)

% w/w = 49

Let weight of solution = 100 g

weight of $\text{H}_2\text{SO}_4 = 49 \text{ g}$

$$\text{vol. of solution} = \frac{\text{mass}}{d} = \frac{100}{1.2} = 83.33 \text{ ml}$$

$$\text{Molarity} = \frac{49}{98} \times \frac{1000}{83.33} = 6 \text{ M}$$

58. Ans (4)

$\pi \propto i c$

59. Ans (3)

At maximum boiling point composition, vapour pressure of solution will be minimum.

60. Ans (2)

For $\text{Mg}(\text{NO}_3)_2$ $i = 1 + 2\alpha = 2.74$

$$2\alpha = 1.74$$

$$\alpha = 0.87$$

$$\alpha = 87\%$$

61. Ans (4)

NCERT page # 53 (XII)

62. Ans (2)

$$\Delta T_f = 2.6 = k_f \times \frac{n}{w(\text{kg})}$$

$$2.6 = 14.4 \times \frac{1.3}{\text{M.w} \times 0.1}$$

$$\text{M.w.} = 72$$

63. Ans (2)

I. Ethyl ether + acetone : +ve deviation from

Raoult's law

II. Chloroform + acetone : -ve deviation from

Raoult's law

III. n-Hexane + h-heptane: ideal solution

64. Ans (3)

in air, $P_{\text{N}_2} = 0.8 \text{ atm}$, $P_{\text{O}_2} = 0.2 \text{ atm}$

$P = K_H X$ (Henry's law)

$$0.8 = 2 \times 10^4 X_{\text{N}_2} \dots\dots(1)$$

$$0.2 = 10^4 X_{\text{O}_2} \dots\dots(2)$$


Ratio of solubility of N_2 & O_2 is 2 : 1

65. Ans (1)

On dilution, H_2O in solution increases so reduction of H_2O occurs at cathode.

66. Ans (3)

At cathode



No. of Equ. of OH^- = No. of Faraday charge

$$\text{OH}^- \Rightarrow 4 \times 10^{-2} \text{ equivalent}$$

$$\Rightarrow 4 \times 10^{-2} \text{ mol}$$

$$\text{Conc. } [\text{OH}^-] = \frac{4 \times 10^{-2}}{\left(\frac{400}{1000}\right)} = 10^{-1} \text{ mol L}^{-1}$$

$$\text{pOH} = -\log(10^{-1}) \Rightarrow \text{pOH} = 1$$

$$\text{pH} = 13$$

67. Ans (2)

Reaction -1 suggest Reduction of B_2 & oxidation of

A^- is spontaneous,

Reaction -2 suggest Reduction of B_2 & oxidation of

C^- is non-spontaneous.

Reaction -3 suggest Reduction of A_2 & oxidation

of D^- is spontaneous.

68. Ans (3)

$$\text{From slow step } r = k_3(\text{Q})^2 (\text{P}) \quad \dots(\text{i})$$

$$\text{from reversible step } k_c = \frac{k_1}{k_2} = \frac{(\text{Q})^2}{(\text{P})} \quad \dots(\text{ii})$$

$$\text{from (i) \& (ii) } r = k_3 \cdot \frac{k_1}{k_2} (\text{P})^2$$

69. Ans (1)

$$2.3 \times 10^{-5} \text{ s}^{-1} \times 90 \times 60 \text{ s} = 2.303 \log \left(\frac{\text{A}_0}{\text{A}_t} \right)$$

$$\log \left(\frac{\text{A}_0}{\text{A}_t} \right) = 0.054 \Rightarrow [\text{A}_t = 0.885 \text{ A}_0]$$

70. Ans (1)

$$\text{Ae}^{-E_a/RT} = \left[\frac{\text{Ae}^{-E_{a1}/RT} \text{Ae}^{-E_{a2}/RT}}{\text{Ae}^{-E_{a3}/RT}} \right]^{2/3}$$

$$= \left[\text{Ae}^{(-E_{a1}-E_{a2}+E_{a3})/RT} \right]^{2/3}$$

$$E_a = \frac{2}{3} [E_{a1} + E_{a2} - E_{a3}]$$

$$= \frac{2}{3} [180 + 80 - 50] = 140 \text{ kJ/mol}$$


71. Ans (1)

[A]	[B]	[C]	rate
0.02	0.02	0.02	2.08×10^{-3}
0.01	0.02	0.02	1.04×10^{-3}
0.02	0.04	0.02	4.16×10^{-3}
0.02	0.02	0.04	8.32×10^{-3}


$$\text{rate} \propto [\text{A}]^1 [\text{B}]^1 [\text{C}]^2$$

$$\text{rate} = K [\text{A}] [\text{B}] [\text{C}]^2$$

72. Ans (1)



$$\text{time} - 3 \times 0.231 = 0.693 \text{ sec}$$



$$\text{time} - 2 \times 0.3465 = 0.693 \text{ sec}$$

$$\frac{\text{C}_A}{\text{C}_B} \rightarrow \frac{[\text{A}]_0/8}{[\text{B}]_0/4} = \frac{1}{2} \times \frac{3}{2} = \frac{3}{4}$$

73. Ans (3)

NCERT Pg # 69-70, para-3.3



74. Ans (1)

$$E^0 = 0 - (E_{M^{+2}/M}^0)$$


$$\Delta G^0 = -nF(-E_{M^{+2}/M}^0)$$

$$96.5 \times 1000 = +2 \times 96500 \times E_{M^{+2}/M}^0$$

75. Ans (4)



on combining



76. Ans (4)

$$\Lambda_M^\infty (Ag_2CrO_4) = 2\Lambda_M^\infty (Ag^+) + \Lambda_M^\infty (CrO_4^{2-})$$

$$= 2 \times 127 + 273$$

$$= 254 + 273 = 527 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$$

$$\Lambda_M^\infty = \frac{k \times 1000}{S}$$

$$s = \frac{2 \times 10^{-2} \times 1000}{527}$$



$$K_{sp} = 4s^3$$

77. Ans (1)

$$Q = n \times n_f \times 96500 = \frac{12.3}{123} \times 6 \times 96500$$

$$= 57900 \text{ C}$$

78. Ans (4)



91. Ans (3)

NCERT-XI, Pg. # 61

92. Ans (4)

NCERT-XI, Pg. # 66, 67

93. Ans (1)

NCERT, Pg. # 67

94. Ans (3)

NCERT XI, Pg. # 67

95. Ans (1)

XI-NCERT Page No. # 58

96. Ans (4)

XI-NCERT Page No. # 57, 58

97. Ans (3)

NCERT XI Pg : 65, 66, 67

98. Ans (4)

NCERT XI Pg : 67

99. Ans (2)

XI-NCERT Page No. # 63, 64, 65

100. Ans (3)

XI-NCERT Page No. # 61, 62, 63, 64, 65

101. Ans (4)

XI-NCERT Page No. # 67, 68

102. Ans (3)

XI-NCERT Page No. # 61

103. Ans (3)

NCERT-XI, Pg. # 65

104. Ans (4)

NCERT-XI, Pg. # 74

105. Ans (1)

NCERT-XI, Pg. # 76-77

106. Ans (1)

NCERT-XI, Pg. # 73, 74

107. Ans (4)

NCERT-XI, Pg # 72

- | | |
|---|---|
| 109. Ans (2)
NCERT-XI, Pg # 75 | 124. Ans (2)
NCERT XII Pg. # 9 |
| 110. Ans (2)
NCERT-XI, Pg # 73 | 126. Ans (4)
NCERT XII, Pg. No. # 14 |
| 111. Ans (2)
NCERT XII Pg # 76 | 127. Ans (3)
NCERT XII Pg. # 21 |
| 112. Ans (3)
NCERT XII Pg # 75, 76 | 128. Ans (1)
NCERT XII, Pg. No. # 13 |
| 113. Ans (3)
NCERT-XI Pg#73 | 129. Ans (3)
NCERT Pg. # 18 |
| 114. Ans (2)
NCERT XI, Pg. # 73-74 | 130. Ans (3)
NCERT, Pg. # 11 |
| 115. Ans (3)
NCERT XI, Page No. 72 | 131. Ans (2)
NCERT XI, Pg. No. # 65 |
| 116. Ans (2)
NCERT-XII Pg. # 22 | 132. Ans (4)
NCERT-XI Pg. # 62 (d) in figure 5.13 |
| 117. Ans (1)
NCERT-XII Pg. # 7 | 133. Ans (4)
NCERT-XI, Pg # 64 |
| 118. Ans (2)
NCERTXII Pg. # 11 | 134. Ans (1)
NCERT (XIth) Pg. # 61, 62 |
| 119. Ans (2)
NCERT XII Page No. 16 | 135. Ans (3)
NCERT XI, Pg. # 59 |
| 120. Ans (2)
NCERT XII Pg. # 13 | 136. Ans (3)
NCERT Pg. # 46 |
| 121. Ans (3)
NCERT-XII Pg. # 6 | 137. Ans (3)
NCERT-XII Pg. # 52(E) & 57(H) |
| 122. Ans (1)
NCERT XII Pg. # 16 | 138. Ans (4)
NCERT(XII) Pg# 43 Para:3.1 |
| 123. Ans (2)
NCERT XII Pg. # 7, 9, 10 | 139. Ans (4)
NCERT(XII) Pg# 50 Para:3.4 |

- | | |
|---|--|
| <p>140. Ans (4)
NCERT(XII) Pg#54/59(H) Para: 3.6</p> <p>141. Ans (3)
NCERT(XII) Pg# 54 (E), 58 (H)</p> <p>142. Ans (1)
NCERT (XIIth) Pg. # 52 (E) 57(H) , Fig. (3.11)</p> <p>143. Ans (2)
NCERT-XII Page#48</p> <p>144. Ans (4)
Module Page#24</p> <p>145. Ans (4)
NCERT, Pg # 49,50

In general oxytocin do not have role in regulation of menstrual cycle.</p> <p>146. Ans (3)
NCERT, Pg # 51,52</p> <p>147. Ans (4)
NCERT Pg. # 50,51</p> <p>148. Ans (4)
NCERT XII Pg # 55, para 1, Pg#51(E)</p> <p>149. Ans (3)
Application based - Allen module.</p> <p>150. Ans (4)
NCERT-XII, Pg. # 60</p> <p>151. Ans (3)
NCERT XII Page # 27</p> <p>152. Ans (2)
NCERT Pg. # 36</p> <p>153. Ans (2)
NCERT - XII, Pg-47 [Ist Para]</p> | <p>154. Ans (4)
NCERT - XII, Pg-49 [Fig-3.8]</p> <p>155. Ans (3)
NCERT (XII) Pg # 44/77(H) Fig. : 3.2</p> <p>156. Ans (4)
NCERT-XII # Page No. 43</p> <p>157. Ans (1)
NCERT Pg. # 30-31</p> <p>158. Ans (1)
NCERT Pg. # 30</p> <p>159. Ans (3)
NCERT-XII, Page # Eng. 47, Hindi 51</p> <p>160. Ans (1)
NCERT Page # Eng. 48, Hindi 52</p> <p>161. Ans (4)
NCERT Page # Eng. 48, Hindi 53</p> <p>162. Ans (3)
NCERT Page # Eng. 48, Hindi 53</p> <p>163. Ans (1)
NCERT Page - 47</p> <p>164. Ans (4)
Second meiotic division occur when sperm comes in contact with ovum in perivitelline space during fertilisation.</p> <p>165. Ans (2)
NCERT-XII, Pg. # 44</p> <p>166. Ans (3)
NCERT Pg. # 46</p> <p>167. Ans (2)
NCERT Pg. # 48</p> |
|---|--|

168. Ans (1)

NCERT, Pg. # 45

Saheli is non-steroidal preparation.

169. Ans (2)

NCERT Pg # 42

170. Ans (2)

NCERT Pg. # 44

171. Ans (1)

NCERT, Pg. # 45 (E)

172. Ans (1)

NCERT, Pg. # 48 (E)

173. Ans (4)

NCERT Pg. # 48

174. Ans (4)

NCERT Pg. # 48

175. Ans (3)

NCERT XII Pg. # 44

176. Ans (2)

NCERT XII, Pg. # 46 & 48

177. Ans (1)

NCERT-XII, Pg. # 44

178. Ans (2)

Module-8, Pg. # 104

179. Ans (1)

NCERT Pg # 57

180. Ans (4)

NCERT Pg No # 60