1394. Minimum Path Cost in a Grid

Difficulty: Medium


https://leetcode.com/problems/minimum-path-cost-in-a-grid

You are given a **0-indexed** $m \times n$ integer matrix grid consisting of **distinct** integers from 0 to $m \times n - 1$. You can move in this matrix from a cell to any other cell in the **next** row. That is, if you are in cell (x, y) such that x < m - 1, you can move to any of the cells (x + 1, 0), (x + 1, 1), ..., (x + 1, n - 1). **Note** that it is not possible to move from cells in the last row.

Each possible move has a cost given by a **0-indexed** 2D array moveCost of size $(m * n) \times n$, where moveCost[i][j] is the cost of moving from a cell with value i to a cell in column j of the next row. The cost of moving from cells in the last row of grid can be ignored.

The cost of a path in grid is the **sum** of all values of cells visited plus the **sum** of costs of all the moves made. Return *the* **minimum** cost of a path that starts from any cell in the **first** row and ends at any cell in the **last** row.

Example 1:



Input: grid = [[5,3],[4,0],[2,1]], moveCost = [[9,8],[1,5],[10,12],[18,6],[2,4],[14,3]]
Output: 17

Explanation: The path with the minimum possible cost is the path 5 -> 0 -> 1.

- The sum of the values of cells visited is 5 + 0 + 1 = 6.
- The cost of moving from 5 to 0 is 3.
- The cost of moving from 0 to 1 is 8.

So the total cost of the path is 6 + 3 + 8 = 17.

So the total cost of this path is 5 + 1 = 6.

Example 2:

```
Input: grid = [[5,1,2],[4,0,3]], moveCost = [[12,10,15],[20,23,8],[21,7,1],[8,1,13],[9,10,25],[5,3,2]]
Output: 6
Explanation: The path with the minimum possible cost is the path 2 -> 3.
- The sum of the values of cells visited is 2 + 3 = 5.
- The cost of moving from 2 to 3 is 1.
```

Constraints:

```
• m == grid.length
```

- n == grid[i].length
- 2 <= m, n <= 50
- qrid consists of distinct integers from 0 to m * n 1.
- moveCost.length == m * n
- moveCost[i].length == n
- 1 <= moveCost[i][j] <= 100