

1583. Paint House III

Difficulty : Hard

<https://leetcode.com/problems/paint-house-iii>

There is a row of m houses in a small city, each house must be painted with one of the n colors (labeled from 1 to n), some houses that have been painted last summer should not be painted again.

A neighborhood is a maximal group of continuous houses that are painted with the same color.

- For example: `houses = [1,2,2,3,3,2,1,1]` contains 5 neighborhoods `[{1}, {2,2}, {3,3}, {2}, {1,1}]`.

Given an array `houses`, an $m \times n$ matrix `cost` and an integer `target` where:

- `houses[i]`: is the color of the house i , and `0` if the house is not painted yet.
- `cost[i][j]`: is the cost of paint the house i with the color $j + 1$.

Return *the minimum cost of painting all the remaining houses in such a way that there are exactly `target` neighborhoods*. If it is not possible, return `-1`.

Example 1:

Input: `houses = [0,0,0,0,0]`, `cost = [[1,10],[10,1],[10,1],[1,10],[5,1]]`, `m = 5`, `n = 2`, `target = 3`

Output: 9

Explanation: Paint houses of this way `[1,2,2,1,1]`

This array contains `target = 3` neighborhoods, `[{1}, {2,2}, {1,1}]`.

Cost of paint all houses $(1 + 1 + 1 + 1 + 5) = 9$.

Example 2:

Input: `houses = [0,2,1,2,0]`, `cost = [[1,10],[10,1],[10,1],[1,10],[5,1]]`, `m = 5`, `n = 2`, `target = 3`

Output: 11

Explanation: Some houses are already painted, Paint the houses of this way `[2,2,1,2,2]`

This array contains `target = 3` neighborhoods, `[{2,2}, {1}, {2,2}]`.

Cost of paint the first and last house $(10 + 1) = 11$.

Example 3:

Input: `houses = [3,1,2,3]`, `cost = [[1,1,1],[1,1,1],[1,1,1],[1,1,1]]`, `m = 4`, `n = 3`, `target = 3`

Output: -1

Explanation: Houses are already painted with a total of 4 neighborhoods `[{3},{1},{2},{3}]` different of `target = 3`.

Constraints:

- `m == houses.length == cost.length`
- `n == cost[i].length`
- `1 <= m <= 100`
- `1 <= n <= 20`
- `1 <= target <= m`
- `0 <= houses[i] <= n`
- `1 <= cost[i][j] <= 104`