Драчов Ярослав Факультет общей и прикладной физики МФТИ

22 ноября 2020 г.

$$\begin{cases} \frac{\partial \mathcal{L}}{\partial \varphi} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\varphi}} = 0, \\ \frac{\partial \mathcal{L}}{\partial z} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{z}} = 0, \\ \varphi(0) = \varphi_0, \\ \dot{\varphi}(0) = \omega_0, \\ z(0) = z_0, \\ \dot{z}(0) = v_0 \end{cases}$$

$$A\cos \omega t + B\sin \omega t.$$

$$x(t) = A\cos \omega t.$$

$$\ddot{x} = -\omega^2 x.$$

$$a = \ddot{x} = -\omega^2 x.$$

$$v = \frac{dx}{dt} = \dot{x}, \qquad a = \frac{dv}{dt} = \frac{d^2x}{dt^2} = \ddot{x}.$$

$$\begin{cases} \ddot{x} = -\omega^2 t, \\ x(0) = A, \\ \dot{x}(0) = 0. \end{cases}$$

$$x(t) = A\cos \omega t.$$

$$\varphi(0) = 0.$$

$$\varphi = \operatorname{arctg} \frac{x}{y}.$$