

# Tutorial #1: Introduction to Algebra

## 1 Exercise 1:

**Proof:**

$$\sum_{i=1}^n i^2 = \frac{(n)(n+1)(2n+1)}{6} \quad (1)$$

**Base case:** When  $n = 1$ , the left hand side (LHS) of (1) is  $1^2 = 1$ , and the right hand side (RHS) of (1) is  $\frac{1 \times (1+1) \times (2 \times 1 + 1)}{6} = 1$ . Thus, (1) is true for  $n = 1$

**Induction step:**

Let  $k \in \mathbb{N}$  be given and suppose (1) is true for  $n = k$ . Then:

$$\begin{aligned} \sum_{i=1}^{k+1} i^2 &= \frac{(k+1)(k+2)(2k+3)}{6} \\ \sum_{i=1}^k i^2 + (k+1)^2 &= \frac{(k+1)(k+2)(2k+3)}{6} \\ \frac{(k)(k+1)(2k+1)}{6} + (k+1)^2 &= \frac{(k+1)(k+2)(2k+3)}{6} \\ \frac{(k+1)(2k^2 + k + 6(k+1))}{6} &= \frac{(k+1)(k+2)(2k+3)}{6} \\ \frac{(k+1)(2k^2 + 7k + 6)}{6} &= \frac{(k+1)(k+2)(2k+3)}{6} \end{aligned}$$

Thus, (1) holds to  $n = k+1$  and the proof of induction step is complete.

**Conclusion:** By the principle of induction, (1) is true for all  $n \in \mathbb{N}$

## 2 Exercise 2:

**Proof:**

$$n! > 2^n \text{ for } n \geq 4 \quad (2)$$

**Base case:** When  $n = 4$ ,  $LHS = 4! = 24$ ,  $RHS = 2^4 = 16$ . Thus, (2) holds for  $n = 4$ .

**Induction step:** Let  $k \in \mathbb{N}$  be given and suppose (2) is true for  $n = k$ . Then:

$$\begin{aligned} (k+1)! &> 2^{k+1} \\ (k+1)! &= k! \times (k+1) > 2k! > 2^{k+1} \end{aligned}$$

Thus, (2) holds to  $n = k+1$  and the proof of induction step is complete.

**Conclusion:** By the principle of induction, (2) is true for all  $n \geq 4$

## 3 Exercise 3:

**Proof:**  $10^{n+1} + 10^n + 1$  is divisible by 3  $\forall n \in \mathbb{N}$ .

As  $10^{n+1} + 10^n + 1$  is represented as 11000.....001. Thus, the total of digit values is 3, which is divisible for 3. Thus,  $10^{n+1} + 10^n + 1$  is divisible by 3  $\forall n \in \mathbb{N}$ .

## 4 Exercise 4:

**Proof:**

$$\sum_{i=0}^n 2^i = 2^{n+1} - 1 \quad (3)$$

**Base case:** When  $n = 1$ ,  $LHS = 2^0 + 2^1 = 3$ ,  $RHS = 2^{1+1} - 1 = 3$ . Thus, (3) holds for  $n = 1$ .

**Induction step:** Let  $k \in \mathbb{N}$  be given and suppose (3) is true for  $n = k$ .

Then:

$$\begin{aligned}\sum_{i=0}^{k+1} 2^i &= 2^{k+2} - 1 \\ \sum_{i=0}^k 2^i + 2^{k+1} &= 2^{k+2} - 1 \\ 2^{k+1} - 1 + 2^{k+1} &= 2^{k+2} - 1 \\ 2^{k+1} + 2^{k+1} &= 2^{k+2}\end{aligned}$$

Thus, (3) holds for  $n = k+1$  and the proof of induction step is complete.

**Conclusion:** By the principle of induction, (3) is true  $\forall n \in \mathbb{N}$

## 5 Exercise 5:

a)

$$\begin{aligned}3x &\equiv 2 \pmod{7} \\ 3x \times 5 &\equiv 2 \times 5 \pmod{7} \\ x &\equiv 3 \pmod{7}\end{aligned}$$

b)

$$\begin{aligned}5x + 1 &\equiv 13 \pmod{23} \\ 5x &\equiv 12 \pmod{23} \\ 5x \times 14 &\equiv 12 \times 14 \pmod{23} \\ x &\equiv 7 \pmod{23}\end{aligned}$$

c

$$2x \equiv 1 \pmod{6}$$

Since there are not exist such  $k$  that  $2k = 1$ , (c) has not solution.

## 6 Exercise 6:

$$\begin{cases} 3x + 7y \equiv 4 \pmod{11} \\ 8x + 6y \equiv 1 \pmod{11} \end{cases}$$

Cayley table for  $\mathbb{Z}_{11}$

$\mathbb{Z}_{11}$	1	2	3	4	5	6	7	8	9	10
1	1	2	3	4	5	6	7	8	9	10
2	2	4	6	8	10	1	3	5	7	9
3	3	6	9	1	4	7	10	2	5	8
4	4	8	1	5	9	2	6	10	3	7
5	5	10	4	9	3	8	2	7	1	6
6	6	1	7	2	8	3	9	4	10	5
7	7	3	10	6	2	9	5	1	8	4
8	8	5	2	10	7	4	1	9	6	3
9	9	7	5	3	1	10	8	6	4	2
10	10	9	8	7	6	5	4	3	2	1

$$\begin{cases} 3x + 7y \equiv 4 \pmod{11} \\ 8x + 6y \equiv 1 \pmod{11} \end{cases} \iff \begin{cases} 8x + 4y \equiv 7 \pmod{11} \\ 8x + 6y \equiv 1 \pmod{11} \end{cases}$$

$$\iff \begin{cases} 8x + 4y \equiv 7 \pmod{11} \\ 2y \equiv 5 \pmod{11} \end{cases} \iff \begin{cases} 8x + 4 \times 8 \equiv 7 \pmod{11} \\ y \equiv 8 \pmod{11} \end{cases}$$

$$\iff \begin{cases} 8x \equiv 8 \pmod{11} \\ y \equiv 8 \pmod{11} \end{cases} \iff \begin{cases} x \equiv 1 \pmod{11} \\ y \equiv 8 \pmod{11} \end{cases}$$

## 7 Exercise 7:

$$\begin{cases} 3x + 5y - 7z \equiv 8 \pmod{83} \\ 8x - 9y + 13z \equiv 13 \pmod{83} \\ 7x + 4y + 5z \equiv 15 \pmod{83} \end{cases} \iff \begin{cases} x + 57y + 53z \equiv 58 \pmod{83} (\times 28) \\ 8x - 9y + 13z \equiv 13 \pmod{83} \\ 7x + 4y + 5z \equiv 15 \pmod{83} \end{cases}$$

$$\iff \begin{cases} x + 57y + 53z \equiv 58 \pmod{83} \\ -56y + 29z \equiv -29 \pmod{83} \\ -6y + 19z \equiv -1 \pmod{83} \end{cases} \iff \begin{cases} x + 57y + 53z \equiv 58 \pmod{83} \\ 27y + 29z \equiv -29 \pmod{83} \\ -6y + 19z \equiv -1 \pmod{83} \end{cases}$$

$$\iff \begin{cases} x + 57y + 53z \equiv 58 \pmod{83} \\ y + 81z \equiv 2 \pmod{83} (\times 40) \\ -6y + 19z \equiv -1 \pmod{83} \end{cases} \iff \begin{cases} x + 37y - 2z \equiv 26 \pmod{83} \\ y + 81z \equiv 2 \pmod{83} \\ 7z \equiv 15 \pmod{83} \end{cases}$$

$$\iff \begin{cases} x + 57y + 53z \equiv 58 \pmod{83} \\ y + 81z \equiv 2 \pmod{83} \\ z \equiv 14 \pmod{83} \end{cases} \iff \begin{cases} x \equiv 23 \pmod{83} \\ y \equiv 30 \pmod{83} \\ z \equiv 14 \pmod{83} \end{cases}$$

Any mistakes might be considered later ;)

## 8 Exercise 8:

Cayley table for  $\mathbb{Z}_7$

$\mathbb{Z}_7$	1	2	3	4	5	6
1	1	2	3	4	5	6
2	2	4	6	1	3	5
3	3	6	2	5	1	4
4	4	1	5	2	6	3
5	5	3	1	6	4	2
6	6	5	4	3	2	1

$$A = \begin{bmatrix} 5 & 2 \\ 6 & 3 \end{bmatrix} \Rightarrow [A|I] = \begin{bmatrix} 5 & 2 & 1 & 0 \\ 6 & 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 6 & 3 & 0 \\ 6 & 3 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 6 & 3 & 0 \\ 0 & 2 & 3 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 6 & 3 & 0 \\ 0 & 1 & 5 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1 & 4 \\ 0 & 1 & 5 & 4 \end{bmatrix} = [I|A^{-1}]$$

Thus, the inverse of  $A$  is  $A^{-1} = \begin{bmatrix} 1 & 4 \\ 5 & 4 \end{bmatrix}$