

## Modular Arithmetic Intro I

Note 6

**Modular arithmetic:** working number “mod  $m$ ”: restrict to only  $\{0, 1, \dots, m-1\}$ ; other numbers are *equivalent* to some number in this set.

Think of a clock; “13-o’clock” is the same as “1-o’clock”;  $2m$  is the same as  $m$ , which is the same as 0. We can go the other way too; 0 is the same as  $-m$ , etc.

Saying  $a \equiv b \pmod{m}$  means:

- $a, b$  have same remainder when divided by  $m$
- $a = b + km$  for some integer  $k$
- $m \mid (a - b)$

**Operations:** Suppose  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ .

Addition/subtraction:  $a \pm c \equiv b \pm d \pmod{m}$

Multiplication:  $ac \equiv bd \pmod{m}$

Exponentiation:  $a^k \equiv b^k \pmod{m}$ . You *cannot* apply the mod to the exponent.

There is no division; there are multiplicative inverses though:  $x^{-1} \equiv a \pmod{m}$  means  $ax \equiv 1 \pmod{m}$ .

## 1 Party Tricks

Note 6

You are at a party celebrating your completion of the CS 70 midterm. Show off your modular arithmetic skills and impress your friends by quickly figuring out the last digit(s) of each of the following numbers:

- Find the last digit of  $11^{3142}$ .
- Find the last digit of  $9^{9999}$ .
- Find the last digit of  $3^{641}$ .

**Solution:**

- First, we notice that  $11 \equiv 1 \pmod{10}$ . So  $11^{3142} \equiv 1^{3142} \equiv 1 \pmod{10}$ , so the last digit is a 1.

(b) 9 is its own multiplicative inverse mod 10, so  $9^2 \equiv 1 \pmod{10}$ . Then

$$9^{9999} = 9^{2(4999)} \cdot 9 \equiv 1^{4999} \cdot 9 \equiv 9 \pmod{10},$$

so the last digit is a 9.

Another solution: We know  $9 \equiv -1 \pmod{10}$ , so

$$9^{9999} \equiv (-1)^{9999} \equiv -1 \equiv 9 \pmod{10}.$$

You could have also used this to say

$$9^{9999} \equiv (-1)^{9998} \cdot 9 \equiv 9 \pmod{10}.$$

(c) Notice that  $3^4 = 9^2$  so using that  $9^2 = 81 \equiv 1 \pmod{10}$ , we have  $3^4 \equiv 1 \pmod{10}$ . We also have that  $641 = 160 \cdot 4 + 1$ , so

$$3^{641} \equiv 3^{4(160)} \cdot 3 \equiv 1^{160} \cdot 3 \equiv 3 \pmod{10},$$

making the last digit a 3.

## 2 Modular Potpourri

### Note 6

Prove or disprove the following statements:

- (a) There exists some  $x \in \mathbb{Z}$  such that  $x \equiv 3 \pmod{16}$  and  $x \equiv 4 \pmod{6}$ .
- (b)  $2x \equiv 4 \pmod{12} \iff x \equiv 2 \pmod{12}$ .
- (c)  $2x \equiv 4 \pmod{12} \iff x \equiv 2 \pmod{6}$ .

### Solution:

- (a) Impossible.

Suppose there exists an  $x$  satisfying both equations.

From  $x \equiv 3 \pmod{16}$ , we have  $x = 3 + 16k$  for some integer  $k$ . This implies  $x \equiv 3 \pmod{2}$ .

From  $x \equiv 4 \pmod{6}$ , we have  $x = 4 + 6l$  for some integer  $l$ . This implies  $x \equiv 0 \pmod{2}$ .

Now we have  $x \equiv 3 \pmod{2}$  and  $x \equiv 0 \pmod{2}$ . Contradiction.

- (b) False, consider  $x \equiv 8 \pmod{12}$ .

The reason we can't eliminate the 2 in the first equation to get the second equation is because 2 does not have a multiplicative inverse modulo 12, as 2 and 12 are not coprime.

- (c) True. We can write  $2x \equiv 4 \pmod{12}$  as  $2x = 4 + 12k$  for some  $k \in \mathbb{Z}$ . Dividing by 2, we have  $x = 2 + 6k$  for the same  $k \in \mathbb{Z}$ . This is equivalent to saying  $x \equiv 2 \pmod{6}$ .

### 3 Modular Inverses

**Note 6** Recall the definition of inverses from lecture: let  $a, m \in \mathbb{Z}$  and  $m > 0$ ; if  $x \in \mathbb{Z}$  satisfies  $ax \equiv 1 \pmod{m}$ , then we say  $x$  is an **inverse of  $a$  modulo  $m$** .

Now, we will investigate the existence and uniqueness of inverses.

- (a) Is 3 an inverse of 5 modulo 14?
- (b) Is 3 an inverse of 5 modulo 10?
- (c) For all  $n \in \mathbb{N}$ , is  $3 + 14n$  an inverse of 5 modulo 14?
- (d) Does 4 have an inverse modulo 8?
- (e) Suppose  $x, x' \in \mathbb{Z}$  are both inverses of  $a$  modulo  $m$ . Is it possible that  $x \not\equiv x' \pmod{m}$ ?

**Solution:**

- (a) Yes, because  $3 \cdot 5 = 15 \equiv 1 \pmod{14}$ .
- (b) No, because  $3 \cdot 5 = 15 \equiv 5 \pmod{10}$ .
- (c) Yes, because  $(3 + 14n) \cdot 5 = 15 + 14 \cdot 5n \equiv 15 \equiv 1 \pmod{14}$ .
- (d) No. For contradiction, assume  $x \in \mathbb{Z}$  is an inverse of 4 modulo 8. Then  $4x \equiv 1 \pmod{8}$ . Then  $8 \mid 4x - 1$ , which is impossible, since  $4x - 1$  is odd (and thus cannot be divisible by 8 either).
- (e) No. We have  $xa \equiv x'a \equiv 1 \pmod{m}$ . So

$$xa - x'a = a(x - x') \equiv 0 \pmod{m}.$$

Multiply both sides by  $x$ , we get

$$xa(x - x') \equiv 0 \cdot x \pmod{m}$$

$$\implies x - x' \equiv 0 \pmod{m}.$$

$$\implies x \equiv x' \pmod{m}$$