



## Lecture 4: Interval Estimation & Goodness of Estimation

STA261 – Probability & Statistics II

Ofir Harari

Department of Statistical Sciences

University of Toronto



# Outline

## Interval Estimation

- Confidence Intervals

- Asymptotic Confidence Intervals

## Goodness of Estimation

- Bias and the Mean Squared Error

- Efficiency and the Cramér-Rao Lower Bound



# point estimation does not reveal uncertainty

## Confidence Intervals

- The last couple of lectures dealt with *point estimation*: finding an estimator  $\hat{\theta}$  with good properties (e.g. consistency) that will hopefully land “in the ballpark” of  $\theta$ .
- But we will inevitably err –

$$\mathbb{P}(\hat{\theta} = \theta) = 0 \text{ (for continuous data)}$$

– and then what...? **because MLE estimator has normal distribution (continuous)**

- We have learned about the notion of standard error (SE) of an estimator –
  - Could report the point estimate along with its SE – a good start
  - Is that what the “margin of error:  $\pm 4$  percentage points” in the newspapers is all about ?
  - Somewhat misleading if the sampling distribution of the estimator is asymmetrical



## Confidence Intervals (cont.)

- The idea of confidence intervals is to provide a range of plausible values for  $\theta$ , rather than a single number.

### Definition

Let  $X_1, \dots, X_n \sim f_\theta$ . A  $100(1 - \alpha)\%$  *confidence interval* for  $\theta$  is a pair of statistics  $L = L(X_1, \dots, X_n)$  and  $U = U(X_1, \dots, X_n)$  such that

$$\mathbb{P}(L \leq \theta \leq U) = 1 - \alpha.$$

We call  $100(1 - \alpha)\%$  the *confidence level*.

note  $\theta$  is fixed,  $L$  and  $U$  are random



## Example: Normal mean with known variance

### Example

1. Let  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$ , where  $\sigma^2$  is assumed to be known. Find a  $100(1 - \alpha)\%$  confidence interval for  $\mu$ .
2. Assuming  $\sigma = 5$ , find a 95% confidence interval for  $\mu$ , if  $n = 16$  and  $\bar{X} = 175$ .

### Solution:

1. Recall that  $\bar{X} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$ , or, equivalently:  $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim \mathcal{N}(0, 1)$ .

Think of a pair of numbers,  $a$  and  $b$ , that satisfy –


$$\mathbb{P}\left(a \leq \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq b\right) = 1 - \alpha$$

– infinitely many options, but a natural choice would be  $a = z_{\alpha/2}$  and  $b = z_{1-\alpha/2}$  – the quantiles of the standard Normal distribution.

the symmetric range over normal curve



## Normal mean with known variance (cont.)



$$1 - \alpha = \mathbb{P} \left( -z_{1-\alpha/2} \leq \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq z_{1-\alpha/2} \right)$$

$$= \mathbb{P} \left( \bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\alpha/2} \leq \mu \leq \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\alpha/2} \right)$$



## Normal mean with known variance (cont.)

We have shown that  $\mathbb{P}\left(\bar{X} - \frac{\sigma}{\sqrt{n}}z_{1-\alpha/2} \leq \mu \leq \bar{X} + \frac{\sigma}{\sqrt{n}}z_{1-\alpha/2}\right) = 1 - \alpha$ ,

hence

$$\left[\bar{X} - \frac{\sigma}{\sqrt{n}}z_{1-\alpha/2}, \bar{X} + \frac{\sigma}{\sqrt{n}}z_{1-\alpha/2}\right]$$

is a  $100(1 - \alpha)\%$  confidence interval for  $\mu$ .

2. Here  $\alpha = 0.05 \implies 1 - \frac{\alpha}{2} = 0.975$ . Substitute

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}}z_{1-\alpha/2} = 175 \pm \frac{5}{\sqrt{16}}z_{0.975} = 175 \pm 1.25 \times 1.96$$

$$\implies \begin{cases} U = 177.45, \\ L = 172.55, \end{cases}$$

thus  $[172.55, 177.45]$  is a 95% confidence interval for  $\mu$  in this case.

idea is find a pivot that approximates parameter  
in this case the pivot is the standardization of sample mean



OO

the true population mean is always the center  
of sampling distribution

OOOOOOOO  
OOOOOOOOOOOOOO

## Understanding confidence intervals

- So,  $[172.55, 177.45]$  is a 95% confidence interval for  $\mu$
- Surely that means “ $\mu$  has a 95% chance of lying between 172.55 and 177.45”...?
  - An outrageous statement!  $\mu$  is a fixed scalar (albeit an unknown one)
  - What is the chance of 5 lying between 4 and 6? Between 3 and 4?
- In the construction of confidence intervals, it is the interval itself that is random
- A 95% Confidence level suggests that if we had infinitely many random samples and calculated the confidence limits for each, 95% of the resultant intervals would include the true parameter value
- Can only hope that the one we have is a good one...






## R simulation

```

> N_Samples <- 100 #No. of random samples
>
> x <- matrix(rnorm(16*N_Samples, mean=175, sd=5), ncol=16) #100 samples of size 16
> xBar <- apply(x, 1, mean) #vector of sample means
> U <- xBar + qnorm(.975)*sigma/4 #upper interval limits
> L <- xBar - qnorm(.975)*sigma/4 #lower interval limits
> uncovered <- which((L>175)|(U<175)) #locating "bad" intervals
>
> plot(c(1:N_Samples), rep(175, N_Samples), type='l', lty=2, col=4, lwd=2)
> segments(1:N_Samples, L, 1:N_Samples, U, lwd=2)
> segments(uncovered, L[uncovered], uncovered, U[uncovered], lwd=4, col=2)

```

95% confidence intervals for  $\mu$  ( $n=16$ ,  $X \sim N(175, 1)$ )





## The pivotal method

### Definition

A *pivotal quantity* (or simply “a pivot”) is a function  $g(X_1, \dots, X_n; \theta)$  of the data and parameter of interest, whose distribution does not depend on any unknown parameter.

- In the last example,  $\bar{X} - \mu \sim \mathcal{N}\left(0, \frac{\sigma^2}{n}\right)$  served as a pivot note sigma and n are all given in the question
- The pivotal method for confidence interval goes as follows:
  1. Find a pivot  $g(X_1, \dots, X_n; \theta)$  and identify its distribution
  2. Find  $a$  and  $b$  such that  $\mathbb{P}(a \leq g(X_1, \dots, X_n; \theta) \leq b) = 1 - \alpha$
  3. Find  $L$  and  $U$  such that  $\mathbb{P}(L \leq \theta \leq U) = 1 - \alpha$



## Example: Normal mean with unknown variance

### Example

Repeat the last example, this time with  $\sigma^2$  unknown, and assuming  $S^2 = 25$ .

- This time  $\bar{X} - \mu$  is no longer a pivot – because  $\sigma^2$  is unknown.
- However, in the first lecture we verified that  $\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t_{n-1}$ , and is therefore a pivot.  
note no population param in pivot
- Now if we look for  $a$  and  $b$  to satisfy


$$\mathbb{P} \left( a \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq b \right) = 1 - \alpha,$$

we can choose  $a = \underline{t_{n-1, \alpha/2}}$  and  $b = t_{n-1, 1-\alpha/2}$  – the quantiles of the  $t_{n-1}$  distribution!

note n-1 d.f.



## Normal mean with unknown variance (cont.)



$$\begin{aligned}
 1 - \alpha &= \mathbb{P} \left( -t_{n-1, 1-\alpha/2} \leq \frac{\bar{X} - \mu}{S/\sqrt{n}} \leq t_{n-1, -\alpha/2} \right) \\
 &= \mathbb{P} \left( \bar{X} - \frac{S}{\sqrt{n}} t_{n-1, 1-\alpha/2} \leq \mu \leq \bar{X} + \frac{S}{\sqrt{n}} t_{n-1, -\alpha/2} \right)
 \end{aligned}$$



## Normal mean with unknown variance (cont.)

- We just showed that

$$\left[ \bar{X} - \frac{S}{\sqrt{n}} t_{n-1, 1-\alpha/2}, \bar{X} + \frac{S}{\sqrt{n}} t_{n-1, 1-\alpha/2} \right]$$

is a  $100(1 - \alpha)\%$  confidence interval for  $\mu$ .

- For our data

$$\bar{X} \pm \frac{S}{\sqrt{n}} t_{n-1, 1-\alpha/2} = 175 \pm \frac{5}{\sqrt{16}} t_{15, 0.975} = 175 \pm 1.25 \times 2.131$$

$$\Rightarrow \begin{cases} U = 177.66, \\ L = 172.34, \end{cases}$$

- Interval of length 5.32 compared to 4.9 when  $\sigma^2$  was assumed to be known  
 CI gets larger compared to if  $\sigma^2$  is known.



## Example: CI for Normal variance

### Example

Find a  $100(1 - \alpha)\%$  confidence interval for  $\sigma^2$ , based on  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$ .


### Solution:

- Recall that  $\frac{(n-1)S^2}{\sigma^2} = \sum_{i=1}^n \left( \frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi_{n-1}^2$  (a pivot).
- We need to find  $a$  and  $b$  such that  $\mathbb{P}\left(a \leq \frac{(n-1)S^2}{\sigma^2} \leq b\right) = 1 - \alpha$   
**problem chi squared not symmetric**
- Ideally, choose them such that the length of the eventual CI is minimized
- A hard optimization problem – not always worth the trouble
- Simply choose  $a = \chi_{n-1, \alpha/2}^2$  and  $b = \chi_{n-1, 1-\alpha/2}^2$ , then a  $(1 - \alpha)100\%$  CI for  $\sigma^2$  will be

$$\left[ \frac{(n-1)S^2}{\chi_{n-1, 1-\alpha/2}^2}, \frac{(n-1)S^2}{\chi_{n-1, \alpha/2}^2} \right]$$



## The $\chi^2$ quantiles



$$\begin{aligned} 1 - \alpha &= \mathbb{P} \left( \chi_{n-1, \alpha/2}^2 \leq \frac{(n-1)S^2}{\sigma^2} \leq \chi_{n-1, 1-\alpha/2}^2 \right) \\ &= \mathbb{P} \left( \frac{(n-1)S^2}{\chi_{n-1, 1-\alpha/2}^2} \leq \sigma^2 \leq \frac{(n-1)S^2}{\chi_{n-1, \alpha/2}^2} \right). \end{aligned}$$



## Asymptotic confidence intervals

- When pivots are hard to find, one can invoke large sample theory, namely:

$$\hat{\theta}_{\text{MLE}} \sim AN(\theta, \mathcal{I}^{-1}(\hat{\theta}_{\text{MLE}})) \quad \text{plugging in}$$

- Can be taken advantage of to construct  $100(1 - \alpha)\%$  *asymptotic confidence interval* of the form

this is CI for normal 's mean

$$\left[ \hat{\theta}_{\text{MLE}} - \frac{z_{1-\alpha/2}}{\sqrt{\mathcal{I}(\hat{\theta}_{\text{MLE}})}}, \hat{\theta}_{\text{MLE}} + \frac{z_{1-\alpha/2}}{\sqrt{\mathcal{I}(\hat{\theta}_{\text{MLE}})}} \right].$$

- For example, for  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \text{Exp}(\lambda)$  we calculated  $\hat{\lambda}_{\text{MLE}} = 1/\bar{X}$  and  $\mathcal{I}(\lambda) = n/\lambda^2$ . A  $100(1 - \alpha)\%$  confidence interval for  $\theta$  would then be  
 substitute mle estimator for true param by plugin principle

$$\left[ \frac{1}{\bar{X}} - \frac{z_{1-\alpha/2}}{\bar{X}\sqrt{n}}, \frac{1}{\bar{X}} + \frac{z_{1-\alpha/2}}{\bar{X}\sqrt{n}} \right].$$





## Comparing different estimators

- So far we have covered two methods of parameter estimation: the Method of Moments and the Maximum Likelihood principle
- Various other methods exist: Bayesian estimation, Least-Squares estimation etc.
- How do we choose between the different types of estimators then?
- Consider the following *loss function*:

$$\mathcal{L}(\hat{\theta}, \theta) = (\theta - \hat{\theta})^2 \quad (\text{the squared error loss})$$

- Inflicts harsh penalties on large deviations from the true parameter value
- Forgiving when it comes to small deviations
- Overall a good candidate for a measure of estimation accuracy – except that... it's a random variable!



## The Mean Squared Error

### Definition

The *Mean Squared Error* of an estimator  $\hat{\theta}$  of a parameter  $\theta$  is

$$\text{MSE}(\hat{\theta}, \theta) = \mathbb{E} \left\{ (\hat{\theta} - \theta)^2 \right\}.$$

- By and large, we use the MSE to assess goodness-of-estimation out of mathematical convenience

**MSE assesses quality of an estimator**

- It could be argued that a more appropriate measure would be the *Mean Absolute Error*  $\mathbb{E} \left\{ |\theta - \hat{\theta}| \right\}$ , but the latter is not differentiable at the origin
- It does not have the following lovely property either –



## The Bias-Variance decomposition

### Proposition

Let  $\hat{\theta}$  be an estimator of a parameter  $\theta$ , and denote

$$b(\hat{\theta}, \theta) = \mathbb{E}[\hat{\theta}] - \theta \quad (\text{the bias of } \hat{\theta}).$$

Then

$$\text{MSE}(\hat{\theta}, \theta) = b^2(\hat{\theta}, \theta) + \text{Var}[\hat{\theta}].$$

**Proof:**

note that  $\theta$  is the RV here,  $\theta$  is just a constant

$$\text{MSE}(\hat{\theta}, \theta) = \mathbb{E} \left\{ (\hat{\theta} - \theta)^2 \right\} = \mathbb{E} \left\{ (\hat{\theta} - \mathbb{E}[\hat{\theta}] + \mathbb{E}[\hat{\theta}] - \theta)^2 \right\}$$

recognize this is bias<sup>2</sup>

$$= \mathbb{E} \left\{ (\hat{\theta} - \mathbb{E}[\hat{\theta}])^2 \right\} + \mathbb{E} \left\{ (\mathbb{E}[\hat{\theta}] - \theta)^2 \right\} + 2\mathbb{E} \left\{ (\hat{\theta} - \mathbb{E}[\hat{\theta}]) (\mathbb{E}[\hat{\theta}] - \theta) \right\}$$

recognize this is the variance of estimator testing

$$= b^2(\hat{\theta}, \theta) + \text{Var}[\hat{\theta}] + 2b(\hat{\theta}, \theta)\mathbb{E} \left\{ (\hat{\theta} - \mathbb{E}[\hat{\theta}]) \right\} = b^2(\hat{\theta}, \theta) + \text{Var}[\hat{\theta}].$$

Note estimator is a constant, so cancel out



## Making sense of the Bias-Variance decomp.

- Think of an Olympic shooter, trying to earn her bread at a competition




sportskeeda.com



## The Bias-Variance decomposition (cont.)

- A shaky hand will not win her any medals




- This is the variance!



## The Bias-Variance decomposition (cont.)

- But if her rifle is out of whack, not even the steadiest of hands will save her




- This is the bias!



## The Bias-Variance decomposition (cont.)

- High accuracy requires both a steady hand and zeroed sights



- This is the MSE!



## Example: Bernoulli trials

### Example

Suppose that we observe a series of Bernoulli trials  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \text{Binom}(1, p)$ . Compare the following estimators of  $p$  (in terms of their MSE):

1.  $\hat{p}_1 = \bar{X}$  (MME and MLE)
2.  $\hat{p}_2 = \frac{\sum_{i=1}^n X_i + 1}{n + 2}$  (Bayesian estimator)
3.  $\hat{p}_3 = X_1$

### Solution:

1. As always with the sample mean,  $\mathbb{E}[\hat{p}_1] = \mathbb{E}[\bar{X}] = \mathbb{E}[X] = p$ . The MSE thus reduces to the variance (why?):

$$\text{MSE}(\hat{p}_1, p) = \text{Var}[\hat{p}_1] = \text{Var}[\bar{X}] = \frac{\text{Var}[X]}{n} = \frac{p(1-p)}{n}.$$





## Bernoulli trials (cont.)

### Solution (cont.):

2. First, let us calculate

$$\mathbb{E}[\hat{p}_2] = \mathbb{E}\left[\frac{\sum_{i=1}^n X_i + 1}{n + 2}\right] = \frac{\sum_{i=1}^n \mathbb{E}[X_i] + 1}{n + 2} = \frac{np + 1}{n + 2},$$

and so the bias is  $b(\hat{p}_2, p) = \frac{np + 1}{n + 2} - p = \frac{1 - 2p}{n + 2}$ . As for the variance,

$$\text{Var}[\hat{p}_2] = \text{Var}\left[\frac{\sum_{i=1}^n X_i + 1}{n + 2}\right] = \frac{\sum_{i=1}^n \text{Var}[X_i]}{(n + 2)^2} = \frac{np(1 - p)}{(n + 2)^2},$$


and finally

$$\text{MSE}(\hat{p}_2, p) = b^2(\hat{p}_2, p) + \text{Var}[\hat{p}_2] = \frac{(1 - 2p)^2 + np(1 - p)}{(n + 2)^2}.$$

3. Trivially,  $\mathbb{E}[\hat{p}_3] = p$ , therefore  $\text{MSE}(\hat{p}_3, p) = \text{Var}[\hat{p}_3] = p(1 - p)$ .



## Bernoulli trials (cont.)





## Example: variance of a Normal population

### Example

For  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$ , Compare the following estimators of  $\sigma^2$ :

1.  $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$  (the sample variance)
2.  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$  (MME and MLE)

**Solution:** 1. easy to calculate because we find a pivot for  $S^2$

1. Recall that  $\frac{(n-1)S^2}{\sigma^2} = \sum_{i=1}^n \left( \frac{X_i - \bar{X}}{\sigma} \right)^2 \sim \chi_{n-1}^2$ , therefore

$$\mathbb{E}[S^2] = \frac{\sigma^2}{n-1} \mathbb{E}[\chi_{n-1}^2] = \frac{\sigma^2}{n-1} \cdot (n-1) = \sigma^2, \quad \text{note } S^2 \text{ is unbiased}$$

hence

since chi squared with n d.f. is gamma(n/2, 1/2) with mean n and variance 2n

$$\text{MSE}(S^2, \sigma^2) = \text{Var}[S^2] = \frac{\sigma^4}{(n-1)^2} \text{Var}[\chi_{n-1}^2] = \frac{\sigma^4 \cdot 2(n-1)}{(n-1)^2} = \frac{2\sigma^4}{n-1}.$$



## Variance of a Normal population (cont.)

**Solution (cont.):**

use the fact that this estimator is a transformation of  $S^2$

2. Clearly  $\hat{\sigma}^2 = \frac{(n-1)S^2}{n}$ , thus

so mme and mle are biased

therefore

$$\mathbb{E}[\hat{\sigma}^2] = \frac{n-1}{n} \mathbb{E}[S^2] = \frac{(n-1)\sigma^2}{n},$$

the asymptotic normality still holds  $n \rightarrow \infty$

$$b(\hat{\sigma}^2, \sigma^2) = \frac{(n-1)\sigma^2}{n} - \sigma^2 = -\frac{\sigma^2}{n}.$$

In addition,

$$\text{Var}[\hat{\sigma}^2] = \frac{(n-1)^2}{n^2} \text{Var}[S^2] = \frac{(n-1)^2}{n^2} \cdot \frac{2\sigma^4}{n-1} = \frac{2(n-1)\sigma^4}{n^2},$$

and finally

$$\text{MSE}(\hat{\sigma}^2, \sigma^2) = b^2(\hat{\sigma}^2, \sigma^2) + \text{Var}[\hat{\sigma}^2]$$

$$= \frac{(2n-1)\sigma^4}{n^2} < \frac{2\sigma^4}{n-1} = \text{MSE}(S^2, \sigma^2) \quad \text{for any } n \geq 2.$$

so mme and mle estimator is more accurate: bias not necessarily bad



## Unbiased estimators

### Definition

We say that  $\hat{\theta}$  is an *unbiased* estimator of  $\theta$  if  $\mathbb{E}[\hat{\theta}] = \theta$  (i.e.  $b(\hat{\theta}, \theta) = 0$ ).

- $\bar{X}$  is always an unbiased estimator of  $\mu = \mathbb{E}[X]$  by LLN
- $S^2$  is always an unbiased estimator of  $\sigma^2 = \text{Var}[X]$  (Practice Problem Set 1)
- Can always correct bias by scaling or shifting – not always beneficial in terms of the MSE
- Unbiased estimators are not necessarily superior to biased ones – yet we love them. Mostly because

1. For an unbiased  $\hat{\theta}$ ,

$$\underline{\text{MSE}(\hat{\theta}, \theta) = \text{Var}[\hat{\theta}]}$$

– compact!

2. We have some seriously nice theory for unbiased estimators



## The Cramér–Rao lower bound



Harald Cramér, 1893–1985

Source: [insurancehalloffame.org](http://insurancehalloffame.org)



Calyampudi R. Rao, 1920–

Source: [isical.ac.in](http://isical.ac.in)



## The Cramér–Rao lower bound (cont.)

### Theorem

Let  $X_1, \dots, X_n \sim f_\theta$ , and let  $\hat{\theta}$  be an unbiased estimator of  $\theta$ . Under some regularity conditions

$$\text{Var}[\hat{\theta}] \geq \mathcal{I}^{-1}(\theta),$$

where  $\mathcal{I}(\theta)$  is the Fisher Information.

**Proof:**

variance for mle are as good as it gets  
for unbiased estimator asymptotically

Denoting  $\underline{x} = (x_1, \dots, x_n)$ , we have

score  $\ell'(\theta) = \frac{\partial \log f(\underline{x}|\theta)}{\partial \theta} = \frac{\frac{\partial f(\underline{x}|\theta)}{\partial \theta}}{f(\underline{x}|\theta)} \Rightarrow \frac{\partial f(\underline{x}|\theta)}{\partial \theta} = \ell'(\theta)f(\underline{x}|\theta) = \underline{u}(\theta)f(\underline{x}|\theta),$

where  $u(\theta)$  is the Score statistic. Now, since  $\hat{\theta}$  is unbiased, we know that

$$\underline{\theta} = \mathbb{E}[\hat{\theta}] = \int \hat{\theta}(\underline{x})f(\underline{x}|\theta)d\underline{x},$$

uses the fact that theta is unbiased here



## The Cramér–Rao lower bound (cont.)

**Proof (cont.):**

theta hat is not a function of theta, so skip..

Having established that

$$\theta = \mathbb{E}[\hat{\theta}] = \int \hat{\theta}(\underline{x}) f(\underline{x}|\theta) d\underline{x},$$

we can differentiate to obtain

$$1 = \frac{\partial \theta}{\partial \theta} = \frac{\partial}{\partial \theta} \int \hat{\theta}(\underline{x}) f(\underline{x}|\theta) d\underline{x} = \int \hat{\theta}(\underline{x}) \frac{\partial f(\underline{x}|\theta)}{\partial \theta} d\underline{x}$$

$\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$

by previously

Text

$$= \int \hat{\theta}(\underline{x}) u(\theta) f(\underline{x}|\theta) d\underline{x} = \mathbb{E}[\hat{\theta} \cdot u(\theta)] = \text{Cov}(\hat{\theta}, u(\theta)) \quad (\text{why?})$$

$$\leq \sqrt{\text{Var}[\hat{\theta}]} \cdot \sqrt{\text{Var}[u(\theta)]} = \sqrt{\text{Var}[\hat{\theta}]} \cdot \sqrt{\mathcal{I}(\theta)},$$

+  $E[\theta]E[u(\theta)]$ ,  
which is 0 because  
 $E[u(\theta)] = 0$

since we proved last week that  $\text{Var}[\hat{\theta}] = \mathcal{I}(\theta)$ , which completes the proof.

this is true by the fact that  $\text{Corr}(X, Y)$   
=  $\text{Cov}(X, Y) / \sqrt{\text{Var}\{X\}\text{Var}\{Y\}} \leq 1$   
i.e. correlation is between -1 and 1

Note  $\text{Var}[u(\theta)] = \mathcal{I}(\theta)$





## Example: the Poisson distribution

- For  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \text{Pois}(\lambda)$  we have already calculated the log-likelihood

$$\ell(\lambda) = n\bar{X} \log \lambda - n\lambda + \text{const}$$

and concluded that the MLE of  $\lambda$  was  $\hat{\lambda}_{\text{MLE}} = \bar{X}$ . In particular, it is unbiased.

- Further calculations yield

$$\ell'(\lambda) = \frac{n\bar{X}}{\lambda} - n \quad \text{and} \quad \ell''(\lambda) = -\frac{n\bar{X}}{\lambda^2}$$

- Note that  $n\bar{X} = \sum_{i=1}^n X_i \sim \text{Pois}(n\lambda)$ , thus  $\mathbb{E}[n\bar{X}] = \text{Var}[n\bar{X}] = n\lambda$ .

**poisson processes**

- The Fisher Information is therefore

$$\mathcal{I}(\lambda) = -\mathbb{E}[\ell''(\lambda)] = \mathbb{E}\left[\frac{n\bar{X}}{\lambda^2}\right] = \frac{n\lambda}{\lambda^2} = \frac{n}{\lambda}.$$



## Example: Poisson distribution (cont.)

- We have calculated  $\mathcal{I}(\lambda) = \frac{n}{\lambda}$
- The CR bound guarantees that for any unbiased estimator  $\hat{\lambda}$  of  $\lambda$

$$\text{Var}[\hat{\lambda}] \geq \mathcal{I}^{-1}(\lambda) = \frac{\lambda}{n}.$$

unbiased

- However, for  $\hat{\lambda}_{\text{MLE}} = \bar{X}$  we have

$$\text{Var}[\hat{\lambda}_{\text{MLE}}] = \text{Var}[\bar{X}] = \frac{\text{Var}[X]}{n} = \frac{\lambda}{n}.$$

- The MLE achieves the CR bound in this case!
- We know for sure then that no unbiased estimator of  $\lambda$  outperforms  $\bar{X}$ .

achieves CR bound: allows to prove  
optimality of unbiased estimators



## Example: Normal distribution

- For  $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$  we have already calculated the log-likelihood

$$\ell(\mu, \sigma^2) = -\frac{n}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^n (X_i - \mu)^2 + \text{const.}$$

- $\frac{\partial \ell}{\partial \sigma^2} = -\frac{n}{2\sigma^2} + \frac{1}{2(\sigma^2)^2} \sum_{i=1}^n (X_i - \mu)^2.$

- $\frac{\partial^2 \ell}{\partial (\sigma^2)^2} = \frac{n}{2\sigma^4} - \frac{1}{\sigma^6} \sum_{i=1}^n (X_i - \mu)^2 = \frac{n}{2\sigma^4} - \frac{1}{\sigma^4} \sum_{i=1}^n \left( \frac{X_i - \mu}{\sigma} \right)^2$

- Recall that  $\sum_{i=1}^n \left( \frac{X_i - \mu}{\sigma} \right)^2 \sim \chi_n^2$ , then **easier to find expected value...**

$$\mathcal{I}(\sigma^2) = -\mathbb{E} \left\{ \frac{\partial^2 \ell}{\partial (\sigma^2)^2} \right\} = -\frac{n}{2\sigma^4} + \frac{1}{\sigma^4} \mathbb{E} \left\{ \sum_{i=1}^n \left( \frac{X_i - \mu}{\sigma} \right)^2 \right\}$$

$$= -\frac{n}{2\sigma^4} + \frac{n}{\sigma^4} = \frac{n}{2\sigma^4}.$$

**sigma<sup>2</sup> is the unit of differentiation**



## Example: Normal distribution (cont.)

- We just calculated:  $\mathcal{I}(\sigma^2) = \frac{n}{2\sigma^4}$
- The CR bound for any unbiased estimator  $\hat{\sigma}^2$  of  $\sigma^2$  is thus

$$\text{Var}[\hat{\sigma}^2] \geq \mathcal{I}^{-1}(\sigma^2) = \frac{2\sigma^4}{n}$$

- The sample variance  $S^2$  is unbiased, and we calculated

$$\text{Var}[S^2] = \frac{2\sigma^4}{n-1} \implies \text{does not achieve the CR bound.}$$

- However,  $\lim_{n \rightarrow \infty} \frac{\text{Var}[S^2]}{\mathcal{I}^{-1}(\sigma^2)} = 1.$

note,  $S^2$  does not achieve CR bound  
but its negligible. We say  $S^2$  is asymptotically efficient



## Efficiency

### Definition

1. We say that an unbiased estimator  $\hat{\theta}$  of a parameter  $\theta$  is *finite sample efficient* (or simply “efficient”) if

$$\text{Var}[\hat{\theta}] = \mathcal{I}^{-1}(\theta).$$

(i.e. it achieves the CR lower bound).

2. We say that  $\hat{\theta}$  is *asymptotically efficient* if

$$\lim_{n \rightarrow \infty} \frac{\text{Var}[\hat{\theta}]}{\mathcal{I}^{-1}(\theta)} = 1.$$

3. The *Relative Efficiency* of an unbiased estimator  $\hat{\theta}_1$  of  $\theta$  with respect to another unbiased estimator  $\hat{\theta}_2$  is

$$\text{eff}(\hat{\theta}_1, \hat{\theta}_2) = \frac{\text{Var}[\hat{\theta}_2]}{\text{Var}[\hat{\theta}_1]}.$$



## Efficiency (cont.)

- In the Poisson example,  $\hat{\lambda}_{\text{MLE}} = \bar{X}$  achieved the CR lower bound, hence it is efficient.

- In the Normal example

$$\lim_{n \rightarrow \infty} \frac{\text{Var}[S^2]}{\mathcal{I}^{-1}(\sigma^2)} = 1,$$

thus  $S^2$  is asymptotically efficient.

note  $S^2$  is not an MLE, but still asymptotically efficient

- When we learned about large sample properties of Maximum Likelihood Estimators, we proved that (under some conditions)

$$\hat{\theta}_{\text{MLE}} \sim AN(\theta, \mathcal{I}^{-1}(\theta)),$$

therefore MLEs are asymptotically unbiased and asymptotically efficient.

doesn't imply that finite sample of MLE is efficient still have to check



## Muon decay example

- $X$  was the cosine of the angle at which electrons are released, with pdf

$$f(x|\alpha) = \frac{1 + \alpha x}{2}, \quad -1 \leq x \leq 1, \quad -1 \leq \alpha \leq 1.$$

- We calculated  $\mathbb{E}[X] = \frac{\alpha}{3}$ . Similarly,

$$\mathbb{E}[X^2] = \int_{-1}^1 x^2 \frac{1 + \alpha x}{2} dx = \frac{1}{3}$$

question from HW

$$\Rightarrow \text{Var}[X] = \mathbb{E}[X^2] - \{\mathbb{E}[X]\}^2 = \frac{1}{3} - \frac{\alpha^2}{9} = \frac{3 - \alpha^2}{9}.$$

- The Method of Moments estimator was found to be  $\hat{\alpha}_{\text{MME}} = 3\bar{X}$ , with

and

$$\mathbb{E}[\hat{\alpha}_{\text{MME}}] = 3\mathbb{E}[\bar{X}] = 3\mathbb{E}[X] = \alpha \Rightarrow \text{unbiased},$$

$$\text{Var}[\hat{\alpha}_{\text{MME}}] = 9\text{Var}[\bar{X}] = \frac{9\text{Var}[X]}{n} = \frac{3 - \alpha^2}{n}.$$

method of moments estimator



## Muon decay example (cont.)

remember we used newton raphson previously

- The Maximum Likelihood estimator,  $\hat{\alpha}_{MLE}$ , is not given in a closed form: cannot calculate its exact sampling distribution.
- We do know that for large samples,  $\hat{\alpha}_{MLE} \sim \mathcal{N}(\alpha, \mathcal{I}^{-1}(\alpha))$  (approximately).
- Calculate **by asymptotic normality**

$$\begin{aligned} \mathcal{I}(\alpha) &= n\mathcal{I}^*(\alpha) = -n\mathbb{E}\left[\frac{\partial^2 \log f(x|\alpha)}{\partial \alpha^2}\right] = -n \int \frac{\partial^2 \log f(x|\alpha)}{\partial \alpha^2} f(x|\alpha) dx \\ &= n \int_{-1}^1 \frac{x^2}{(1+\alpha x)^2} \frac{1+\alpha x}{2} dx = \begin{cases} \frac{n \left( \log \frac{1+\alpha}{1-\alpha} - 2\alpha \right)}{2\alpha^3} & , \quad \alpha \neq 0, \\ \frac{n}{3} & , \quad \alpha = 0. \end{cases} \end{aligned}$$

can also calculate fisher info with  $\mathcal{I}^*$






## Muon decay example (cont.)

- The *asymptotic* relative efficiency is thus

$$\text{eff}(\hat{\alpha}_{\text{MME}}, \hat{\alpha}_{\text{MLE}}) = \frac{\text{Var}[\hat{\alpha}_{\text{MLE}}]}{\text{Var}[\hat{\alpha}_{\text{MME}}]} = \frac{2\alpha^3}{3 - \alpha^2} \left( \log \frac{1 + \alpha}{1 - \alpha} - 2\alpha \right)^{-1} \quad (\alpha \neq 0).$$



$\text{Var}[\alpha_{\text{MME}}]$  increasing toward boundary

- Note how much efficiency the MME loses (relative to the MLE) close to the boundary of the parameter space!