

1 Compactness

Definition. Let S be a non-empty set of real numbers.

1. A real number x is called an upper bound for S if $x \geq s$ for all $s \in S$.
2. A real number x is the **least upper bound** (or supremum) for S if x is an upper bound for S and $x \leq y$ for every upper bound y of S .
3. $p = \sup A$ if and only if for every $\epsilon > 0$ there is an $x \in A$ with $x > p - \epsilon$, and $x \leq p$

Theorem. By completeness of real numbers, every non-empty set $A \in \mathbb{R}$ which is bounded above has a least upper bound, i.e. $\sup(A)$ exists.

Proposition. A closed and bounded set $A \in \mathbb{R}$, $\inf(A) \in A \wedge \sup(A) \in A$. Implies that supremum/infimum are equivalent to maximum/minimum

Proof.

Assume the set didn't contain its supremum (which exists since the set is bounded). Since the set is closed, an entire neighbourhood of the supremum lies outside the set. This neighbourhood contains upper bounds of the set that are lower than the supremum, which is a contradiction. \square

Note. While the maximum of a set must be contained in the set, the supremum does not necessarily have to be contained in the set

Definition. A set $S \subseteq \mathbb{R}^n$ is **compact** if it is both closed and bounded.

Remark. A finite union of compact sets is compact because a finite union of closed and bounded sets is closed and bounded; however an infinite union of compact is not necessarily compact.

For example, $S_n = [\frac{1}{n}, 1]$ and $U = \bigcup_i S_i$. $x = 0 \subseteq U^c$, whose neighborhood meets U and U^c . Since x a boundary point of U but not in U , then U not closed.

Theorem. The following are equivalent,

1. S is compact
2. **Bolzano-Weierstrass** Every sequence in S has a convergent subsequence, that is, if $(X_n)_{n=1}^\infty \subseteq S$ (note (X_n) may not be convergent), then there exists a subsequence X_{n_k} and a point $x \in S$ such that $X_{n_k} \rightarrow x$
3. **Heine-Borel** Every open cover of S admits a finite subcover; that is, if $\{\cup_i\}_{i \in \mathbb{I}}$ is a collection of open sets such that $S \subseteq \{\cup_i\}_{i \in \mathbb{I}} U_i$, then there exists a finite subset $J \subseteq \mathbb{I}$ such that $S \subseteq \{\cup_i\}_{i \in J} U_i$

Theorem. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a continuous function. If $K \subseteq \mathbb{R}^n$, then $f(K)$ is also compact.

Theorem. Extreme Value Theorem If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuous function and $K \subseteq \mathbb{R}^n$ is a compact set, then there exists $x_{\min}, x_{\max} \in K$ such that for every $x \in K$, $f(x_{\min}) \leq f(x) \leq f(x_{\max})$; that is, f achieves both its extreme values on K .

Remark. x_{\min}, x_{\max} need not be minimum and maximum in K .

Connectedness

Definition 1.1. * A set $S \subseteq \mathbb{R}^n$ is said to be **disconnected** if there exist non-empty $S_1, S_2 \subseteq S$ such that

1. $S_1 \neq \emptyset \wedge S_2 \neq \emptyset$
2. $S = S_1 \cup S_2$
3. $\overline{S_1} \cap S_2 = \emptyset$ and $S_1 \cap \overline{S_2} = \emptyset$

Here (S_1, S_2) is a disconnection of S . If S admits no disconnection, we say that S is **connected**.

Theorem. A set $S \subseteq \mathbb{R}$ is connected if and only if S is an interval

Definition. If $S \subseteq \mathbb{R}^n$ then a **path** in S is any continuous map $\gamma : \mathbb{R} \rightarrow \mathbb{R}^n : [0, 1] \rightarrow S$. We say that S is **path-connected** if for every two points $a, b \in S$ there exists a path $\gamma : [0, 1] \rightarrow S$ such that $\gamma(0) = a$ and $\gamma(1) = b$

Remark. path-connected is a weaker statement than connectedness. Any path-connected set is necessarily connected; the converse not always true. Geometrically, a set is path-connected if between any points in the set, we can draw a line between those two points that never leaves the set

Theorem. The continuous image of a (path) connected set is (path) connected. More precisely, If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous and $S \subseteq \mathbb{R}^n$ is (path) connected, then $f(S)$ is (path) connected.

Theorem. Intermediate Value Theorem Let $V \subseteq \mathbb{R}^n$ be a (path) connected set and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function. Let $a, b \in V$ and assume that $f(a) < f(b)$. Then for every $c \in \mathbb{R}$, such that $f(a) < c < f(b)$ there exists an $x \in V$ such that $f(x) = c$

Theorem. disconnection from a set to subset

Let (S_1, S_2) be disconnection for S . Let $U \subseteq S$ and $U_1 = S_1 \cap U$, $U_2 = S_2 \cap U$. If U meets both S_1 and S_2 , i.e., $U_1 \neq \emptyset$ and $U_2 \neq \emptyset$, then (U_1, U_2) is a disconnection for U

Corollary. If a set S is disconnected (S_1, S_2) and a subset $U \subseteq S$ is connected. Then U must be contained in either S_1 or S_2 , i.e. $U \subseteq S_1$ or $U \subseteq S_2$

Corollary. *Let A be a connected set, suppose B satisfies $A \subseteq B \subseteq \overline{B}$, then B is connected.*

Theorem. *Any path connected set is also connected*

Proof.

Proof by contradiction, assume S is not connected. Then there exists a disconnection (S_1, S_2) . Consider $a \in S_1, b \in S_2$. Since S is path-connected, there exists $\gamma : [0, 1] \rightarrow S$, such that $\gamma[0] = a, \gamma[1] = b$. Let $U = \gamma([0, 1]) \subseteq S$, then U is path-connected since γ is continuous and $[0, 1]$ an interval in \mathbb{R} is path-connected. Now introduce contradiction by showing U is not connected. Let $U_1 = U \cap S_1, U_2 = U \cap S_2$.

$$U_1 \cup U_2 = (U \cap S_1) \cup (U \cap S_2) = U \cap (S_1 \cup S_2) = U$$

and

$$\overline{U_1} \cap U_2 \subseteq (\overline{U} \cap \overline{S_1}) \cap (U \cap S_2) = \overline{S_1} \cap S_2 = \emptyset$$

Similarly $U_1 \cap \overline{U_2} = \emptyset$. Then shown (U_1, U_2) is a disconnection of U , which is a contradiction. \square

Theorem. *If $S \subseteq \mathbb{R}^n$ is connected and open, then S is path-connected.*

Remark. For example, the topologist's sine curve is connected but not path-connected.

$$\{(x, \sin(\frac{1}{x})) : x \in \mathbb{R} \setminus \{0\}\} \cup (0, 0)$$

	closed sets	compact sets
finite union	yes	yes
infinite union	no i.e. $S = [\frac{1}{n}, 1]$	no
finite intersection	yes	yes
infinite intersection	yes	yes

Definition. A set $S \subseteq \mathbb{R}^n$ is **convex** if for all $a, b \in S$, there exists a path

$$\lambda : [0, 1] \rightarrow l$$

where $l \subseteq S$ is a line and $\lambda(0) = a$ and $\lambda(1) = b$

Remark. Convex set is path-connected and thus connected.