



Lecture 6: The Elements of Hypothesis Testing

STA261 – Probability & Statistics II

Ofir Harari

Department of Statistical Sciences

University of Toronto



Outline

Introduction

The Statistical Hypothesis Testing Framework

Basic Definitions

Simple Hypotheses

Significance Level and Power

Likelihood Ratio Tests and the Neyman–Pearson Lemma



Introduction

- In order to receive the FDA's approval, new drugs must go through Clinical Research, or *Clinical Trials* (specifically: *Treatment Trials*).
- Clinical trials are conducted in phases:
 - *Phase I and II trials* (20–300 patients): evaluating safety, identifying side effects and determining effectiveness.
 - *Phase III trials* (1,000–3,000 patients): confirming effectiveness. We shall focus on this phase.
- Typically there will be (at least) two groups: treatment and control.
- Each group consists of patients (volunteers) with the disease/condition, who have met the selection criteria, blindly and randomly allocated to avoid research bias. i.e. woman more prone to disease..
- In "placebo-controlled" trials, patients in the treatment group receive the drug under investigation, while patients in the control group receive a placebo.
double blind: both patient and doctor does not know which group peps are in



Introduction

- How do we determine the effectiveness of a treatment?

- Denote:

$$\left\{ \begin{array}{l} n_{\text{tr}} - \text{Number of patients in the treatment group} \\ n_{\text{pl}} - \text{Number of patients in the placebo group} \\ p_{\text{tr}} - \text{the probability of being cured among patients in the treatment group} \\ p_{\text{pl}} - \text{the probability of being cured among patients in the placebo group} \\ X_{\text{tr}} - \text{Number of patients in the treatment group who were cured} \\ X_{\text{pl}} - \text{Number of patients in the placebo group who were cured} \end{array} \right.$$

Then $X_{\text{tr}} \sim \text{Binom}(n_{\text{tr}}, p_{\text{tr}})$ and $X_{\text{pl}} \sim \text{Binom}(n_{\text{pl}}, p_{\text{pl}})$

- To prove the effectiveness of the drug under investigation, the pharmaceutical company must **use the data** to show that

$$p_{\Delta} := p_{\text{tr}} - p_{\text{pl}} > 0$$

beyond a reasonable doubt.



Hypotheses and types of errors

- The data: $X_1, \dots, X_n \sim f_\theta$
- The parameter: θ (in our example $p_\Delta := p_{\text{tr}} - p_{\text{pl}}$)
publication bias: discard result that does not support alternative hypothesis
- The Null Hypothesis \mathcal{H}_0 – usually the conservative view (“no effect”)
 - In our example: $\mathcal{H}_0 : p_\Delta \leq 0$ (i.e. the new drug is no better than placebo)
- The Alternative Hypothesis \mathcal{H}_1 – usually represents change of reality
 - In our example: $\mathcal{H}_1 : p_\Delta > 0$ (i.e. the new drug IS better than placebo)

rejection rule

- The decision: to reject \mathcal{H}_0 or not to reject \mathcal{H}_0 there is no accept \mathcal{H}_1
- Type I Error: incorrectly rejecting \mathcal{H}_0 (i.e. a *false discovery*)
 - In our example: falsely declaring the medication effective
- Type II Error: incorrectly retaining \mathcal{H}_0
 - In our example: failing to approve an authentically effective medication

★ Which type of error would you consider to be more serious?

type 1: false discovery is more serious than sparing the world an authentic discovery



Formulating the hypotheses

- The parameter space: Θ – all possible values of θ
 - In our example: $\Theta = [-1, 1]$ (all possible values of $p_{\Delta} := p_{\text{tr}} - p_{\text{pl}}$)

- The competing hypotheses are typically of the form

$$\begin{cases} \mathcal{H}_0 : \theta \in \Theta_0 \\ \mathcal{H}_1 : \theta \in \Theta_1 \end{cases}$$

- In our example:

theta_0 and theta_1 definitely dont overlap

$$\begin{cases} \mathcal{H}_0 : p_{\Delta} \in [-1, 0] \\ \mathcal{H}_1 : p_{\Delta} \in (0, 1] \end{cases}$$

- When $\Theta = \{\theta_0, \theta_1\}$, the hypotheses

$$\begin{cases} \mathcal{H}_0 : \theta = \theta_0 \\ \mathcal{H}_1 : \theta = \theta_1 \end{cases}$$

are called *simple hypotheses*.

- A statistical test: a data driven, probabilistic decision rule with regard to \mathcal{H}_0 (reject/not reject). input: sample ==> output: hypothesis



Significance Level and Power

Definition

Suppose that we test the simple hypotheses

$$\begin{cases} \mathcal{H}_0 : \theta = \theta_0, \\ \mathcal{H}_1 : \theta = \theta_1. \end{cases}$$

1. The significance level of the test is the probability of a type I error,

$$\alpha = \mathbb{P} \left(\begin{array}{c} \text{rejecting} \\ \mathcal{H}_0 \end{array} \middle| \theta = \theta_0 \right).$$

not conditional probability, just with given param

2. The power of a statistical test is the probability of NOT making a type II error, $\pi = 1 - \beta$, where

$$\beta = \mathbb{P} \left(\begin{array}{c} \text{not rejecting} \\ \mathcal{H}_0 \end{array} \middle| \theta = \theta_1 \right).$$

rejecting H_0 when we should reject it

a good test \rightarrow small beta and high power



Example

Example

You just purchased a matchbox. The company states on its website that 10% of their matches are defective, but you have that growing feeling that the true proportion is 50%. The company's customer relations representative proposes that you sample two matches at random, and if at least one turns out to be defective – your claim will be accepted.

- Denote: X – the number of defective matches in a random sample of size 2;
- $X \sim \text{Binom}(2, p)$ specifies probability distribution completely \Rightarrow hence simple hypotheses
- The (simple) hypotheses in this case are –

$$\begin{cases} \mathcal{H}_0 : p = 0.1 & \text{null: 10\% defective} \\ \mathcal{H}_1 : p = 0.5 & \text{alternative: 50\% defective} \end{cases}$$

- The test: reject \mathcal{H}_0 if $X \geq 1$



Example (cont.)

- $$\begin{cases} \mathcal{H}_0 : p = 0.1 \\ \mathcal{H}_1 : p = 0.5 \end{cases}, \quad \text{reject } \mathcal{H}_0 \text{ if } X \geq 1$$
- The significance level of the test is – **rejecting \mathcal{H}_0 when \mathcal{H}_0 is true**

$$\alpha = \mathbb{P} \left(\begin{array}{c} \text{rejecting} \\ \mathcal{H}_0 \end{array} \middle| p = 0.1 \right) = \mathbb{P}(X \geq 1 | p = 0.1)$$

$$= 1 - \mathbb{P}(X = 0 | p = 0.1) = 1 - 0.9^2 = 0.18$$

just binom

- The probability of a type II error is – **not rejecting \mathcal{H}_0 when \mathcal{H}_1 is true**

$$\beta = \mathbb{P} \left(\begin{array}{c} \text{not rejecting} \\ \mathcal{H}_0 \end{array} \middle| p = 0.5 \right) = \mathbb{P}(X = 0 | p = 0.5)$$

$$= 0.5^2 = 0.25,$$

note here alpha and beta can be calculated before observations, with given decision rule, test statistic, and its underlying distribution

and the power of the test is $\pi = 1 - \beta = 0.75$.



The α - β tradeoff

Decision \ Truth	\mathcal{H}_0 is correct	\mathcal{H}_0 is incorrect
Not rejecting \mathcal{H}_0	Correct decision ($1 - \alpha$)	Type I error (Significance level α)
Rejecting \mathcal{H}_0	Type II error (β)	Correct decision (Power $\pi = 1 - \beta$)

beta: false negative

alpha: false positive

- Ideally, we would like both α and β to be as small as possible
- Unfortunately, they have conflicting agendas...
- Suppose that we wish to test

$$\begin{cases} \mathcal{H}_0 : \mu = \mu_0 \\ \mathcal{H}_1 : \mu = \mu_1 \end{cases},$$


where $\mu_1 > \mu_0$, based on a sample $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$, where σ^2 is known.

- We reject \mathcal{H}_0 if $\underline{\bar{X}} \geq c$ for some threshold c , and recall that $\bar{X} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$.
test statistic \rightarrow mean



The α - β tradeoff (cont.)

- As α decreases...



falsely retaining H_0

falsely rejecting H_0



The α - β tradeoff (cont.)

- The general convention is that type I errors are more dangerous than type II errors, ergo, controlling α is prioritized over controlling β .
- The acceptable practice is thus to focus on tests with significance level α (where α is a small number, typically 0.05 or less), and among those to search for the test with the smallest β .
 - also known as the most powerful test.
- But can we search for such a test in a principled way?
- Yes! And (perhaps not so) surprisingly, it involves the likelihood function.



Test Statistics

- Any statistical test is based on a test statistic $T(\underline{X})$: a sample statistic whose distribution under \mathcal{H}_0 is known.
- In a previous example wished to test

$$\begin{cases} \mathcal{H}_0 : \mu = \mu_0 \\ \mathcal{H}_1 : \mu = \mu_1 \end{cases},$$

where $\mu_1 > \mu_0$, based on a sample $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$, where σ^2 was known.

- The proposed test was: reject \mathcal{H}_0 if $\bar{X} \geq c$, for some c .
- Here $\bar{X} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$, hence $\bar{X} \stackrel{\mathcal{H}_0}{\sim} \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right)$.
- In this example \bar{X} is our test statistic.

known by H_0



Rejection Regions

$$\begin{cases} \mathcal{H}_0 : \mu = \mu_0 \\ \mathcal{H}_1 : \mu = \mu_1 (> \mu_0) \end{cases}, \quad \text{reject } \mathcal{H}_0 \text{ if } \bar{X} \geq c$$

- Remember that we restrict ourselves to tests with significance level α
- Can we find a value of c for which the probability of a type I error will be α ?
- Recall that under \mathcal{H}_0 (i.e. assuming $\mu = \mu_0$), $\bar{X} \sim \mathcal{N}\left(\mu_0, \frac{\sigma^2}{n}\right)$, thus


$$\alpha = \mathbb{P}\left(\begin{array}{c} \text{rejecting} \\ \mathcal{H}_0 \end{array} \middle| \mu = \mu_0\right) = \mathbb{P}(\bar{X} \geq c | \mu = \mu_0) = 1 - \mathbb{P}(\bar{X} \leq c | \mu = \mu_0)$$

$$= 1 - \mathbb{P}\left(\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \leq \frac{c - \mu_0}{\sigma/\sqrt{n}} \middle| \mu = \mu_0\right) = 1 - \Phi\left(\frac{c - \mu_0}{\sigma/\sqrt{n}}\right)$$

$$\Rightarrow \Phi\left(\frac{c - \mu_0}{\sigma/\sqrt{n}}\right) = 1 - \alpha \Rightarrow \frac{c - \mu_0}{\sigma/\sqrt{n}} = z_{1-\alpha} \Rightarrow c = \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha}$$



Rejection Regions (cont.)





Rejection Regions (cont.)

- We calculated $c = \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha}$, hence the test –

$$\text{“reject } \mathcal{H}_0 \text{ if } \bar{x} \geq \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha} \text{”}$$

has significance level α .

- In general, any statistical test that is based on test statistic $T(\underline{X})$ will be of the form

$$\text{“reject } \mathcal{H}_0 \text{ if } T(\underline{x}) \in \mathcal{C}, \text{” for some region } \mathcal{C} \subset \mathbb{R}^n.$$

We call \mathcal{C} the Rejection Region.

- In our example the rejection region was

$$\mathcal{C} = \left\{ \underline{x} \in \mathbb{R}^n : \bar{x} \geq \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha} \right\}.$$

reject H_0 if \bar{x} in rejection region \rightarrow gives confidence level of α



Rejection Regions (cont.)

- For example, suppose that $n = 16$, $\sigma^2 = 25$, and we wish to test

$$\begin{cases} \mathcal{H}_0 : \mu = 175 & \text{vs.} \\ \mathcal{H}_0 : \mu = 180 \end{cases}$$

at a 5% level.

- Here the rejection region is

$$\begin{aligned} \mathcal{C} &= \left\{ \underline{x} : \bar{x} \geq \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha} \right\} = \left\{ \underline{x} : \bar{x} \geq 175 + \frac{5}{4} \underbrace{z_{0.95}}_{1.645} \right\} \\ &= \{ \underline{x} : \bar{x} \geq 177.06 \}. \end{aligned}$$

- Congratulations! You just designed your first statistical test. But is it the optimal (i.e. most powerful) test?



The Likelihood Ratio statistic

- Consider again the problem of testing the simple hypotheses

$$\begin{cases} \mathcal{H}_0 : \theta = \theta_0 \\ \mathcal{H}_1 : \theta = \theta_1 \end{cases}$$

based on $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} f_\theta$.

- The Likelihood Ratio is

$$\lambda(\underline{x}) := \frac{\mathcal{L}(\theta_1)}{\mathcal{L}(\theta_0)} = \frac{f(x_1, \dots, x_n | \theta_1)}{f(x_1, \dots, x_n | \theta_0)}.$$

- Loosely speaking, $\lambda(\underline{x})$ measures how likely \mathcal{H}_1 is to be true compared to \mathcal{H}_0 , with large values supporting the case for rejecting \mathcal{H}_0 .
- It is thus reasonable to consider tests with rejection regions of the form

$$\mathcal{C} = \{\underline{x} \in \mathbb{R}^n : \underline{\lambda(\underline{x})} \geq c\}.$$



Likelihood Ratio Tests

Definition

A statistical test based on the rejection region

$$\mathcal{C} = \left\{ \underline{x} \in \mathbb{R}^n : \lambda(\underline{x}) = \frac{\mathcal{L}(\theta_1)}{\mathcal{L}(\theta_0)} \geq c \right\},$$

for c satisfying $\mathbb{P}(\lambda(\underline{x}) \geq c | \theta = \theta_0) = \alpha$, is called a *likelihood ratio test* (LRT) at (significance) level α .

- Back to our example:

$$\begin{cases} \mathcal{H}_0 : \mu = \mu_0 \\ \mathcal{H}_1 : \mu = \underline{\mu_1} (> \mu_0) \end{cases}, \quad X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mu, \sigma^2)$$

- Here

$$\lambda(\underline{x}) = \frac{\mathcal{L}(\mu_1)}{\mathcal{L}(\mu_0)} = \frac{\frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu_1)^2 \right\}}{\frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu_0)^2 \right\}}$$



Likelihood Ratio Tests (cont.)

$$\lambda(\underline{x}) = \frac{\frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left\{ -\frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu_1)^2 \right\}}{\frac{1}{(2\pi\sigma^2)^{n/2}} \exp \left\{ -\frac{1}{\sigma^2} \sum_{i=1}^n (x_i - \mu_0)^2 \right\}} = \frac{\exp \left\{ -\frac{1}{\sigma^2} \sum_{i=1}^n x_i^2 + \frac{n\mu_1}{\sigma^2} \bar{x} - \frac{n\mu_1^2}{2\sigma^2} \right\}}{\exp \left\{ -\frac{1}{\sigma^2} \sum_{i=1}^n x_i^2 + \frac{n\mu_0}{\sigma^2} \bar{x} - \frac{n\mu_0^2}{2\sigma^2} \right\}}$$

$$= \exp \left\{ \frac{n(\mu_1 - \mu_0)}{\sigma^2} \bar{x} - \frac{n(\mu_1^2 - \mu_0^2)}{\sigma^2} \right\},$$

2

therefore

$$\lambda(\underline{x}) \geq c \iff \frac{n(\mu_1 - \mu_0)}{\sigma^2} \bar{x} - \frac{n(\mu_1^2 - \mu_0^2)}{\sigma^2} \geq c_1$$

a constant

$$\iff \frac{n(\mu_1 - \mu_0)}{\sigma^2} \bar{x} \geq c_2 \iff \bar{x} \geq c_3, \text{ (since } \mu_1 > \mu_0 \text{)}$$

2

hence

positive constant

$$\mathcal{C} = \{\underline{x} \in \mathbb{R}^n : \lambda(\underline{x}) \geq c\} = \{\underline{x} \in \mathbb{R}^n : \bar{x} \geq c_3\}.$$

this is the test we did earlier on normal distribution samples



Likelihood Ratio Tests (cont.)

$$\mathcal{C} = \{\underline{x} \in \mathbb{R}^n : \lambda(\underline{x}) \geq c\} = \{\underline{x} \in \mathbb{R}^n : \bar{x} \geq c_3\}$$

- It now remains to find c_3 such that

$$\alpha = \mathbb{P} \left(\begin{array}{c} \text{Type I} \\ \text{error} \end{array} \right) = \mathbb{P}(\underline{X} \in \mathcal{C} | \mu = \mu_0) = \mathbb{P}(\bar{X} \geq c_3 | \mu = \mu_0)$$

in rejection region

- Wait a minute – we’ve already calculated it! It was

$$\underline{c_3 = \mu_0 + \frac{\sigma}{\sqrt{n}} z_{1-\alpha}}$$

- It turns out that the test we proposed was a likelihood ratio test all along.
- But is it the *most powerful* (MP) test at level α ?



The Neyman–Pearson Lemma

Definition

Suppose that we observe $X_1, \dots, X_n \stackrel{\text{i.i.d.}}{\sim} f_\theta$, and consider the simple hypotheses $\mathcal{H}_0 : \theta = \theta_0$ vs. $\mathcal{H}_1 : \theta = \theta_1$. We say that a test is a most powerful (MP) test at level α if

1. the significance level of the test is α , and
2. no other test at level α has greater power.

Lemma (Neyman–Pearson Lemma)


The likelihood ratio test, based on the rejection region

$$\mathcal{C} = \left\{ \underline{x} \in \mathbb{R}^n : \lambda(\underline{x}) = \frac{\mathcal{L}(\theta_1)}{\mathcal{L}(\theta_0)} \geq c \right\},$$

with c satisfying $\mathbb{P}(\lambda(\underline{X}) \geq c) = \alpha$ is an MP test at level α .



The Neyman–Pearson Lemma (cont.)



Egon Pearson, 1895-1980

Source: swlearning.com



The Neyman–Pearson Lemma (cont.)

Proof for a continuous f_θ :

Denote the rejection region of the LRT by \mathcal{C} . Note that

$$\alpha = \mathbb{P}(\underline{X} \in \mathcal{C} | \theta = \theta_0) = \int_{\mathcal{C}} f(\underline{x} | \theta_0) d\underline{x} = \int_{\mathcal{C}} \mathcal{L}(\theta_0) d\underline{x}.$$

rejecting null given null correct

Consider now another test at level α , with rejection region \mathcal{D} . The same

calculations would yield $\alpha = \int_{\mathcal{D}} \mathcal{L}(\theta_0) d\underline{x}$, hence **prove that \mathcal{C} yields higher power when compared to any other region \mathcal{D} (i.e. MP)**

$$\int_{\mathcal{C}} \mathcal{L}(\theta_0) d\underline{x} = \int_{\mathcal{D}} \mathcal{L}(\theta_0) d\underline{x}. \quad \text{since both equal to alpha}$$

Write $\mathcal{C} = (\mathcal{C} \cap \mathcal{D}) \cup (\mathcal{C} \cap \overline{\mathcal{D}})$ (a disjoint union), we have

$$\int_{\mathcal{C}} \mathcal{L}(\theta_0) d\underline{x} = \int_{\mathcal{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\underline{x} + \int_{\mathcal{C} \cap \overline{\mathcal{D}}} \mathcal{L}(\theta_0) d\underline{x}.$$

by one property of integration

Likewise, $\int_{\mathcal{D}} \mathcal{L}(\theta_0) d\underline{x} = \int_{\mathcal{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\underline{x} + \int_{\overline{\mathcal{C}} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\underline{x}.$



The Neyman–Pearson Lemma (cont.)

Proof (cont.):

We have shown so far

$$\int_{C \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x} + \int_{C \cap \overline{\mathcal{D}}} \mathcal{L}(\theta_0) d\mathbf{x} = \int_{C \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x} + \int_{\overline{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x},$$

or simply

$$\int_{C \cap \overline{\mathcal{D}}} \mathcal{L}(\theta_0) d\mathbf{x} = \int_{\overline{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x}.$$

It is now time to recall that for any $\mathbf{x} \in C$ we have $\lambda(\mathbf{x}) = \frac{\mathcal{L}(\theta_1)}{\mathcal{L}(\theta_0)} \geq c$, and for

every $\mathbf{x} \in \overline{C}$ we have $\frac{\mathcal{L}(\theta_1)}{\mathcal{L}(\theta_0)} < c$, hence

by definition of region C

$$c \int_{C \cap \overline{\mathcal{D}}} \mathcal{L}(\theta_0) d\mathbf{x} \leq \int_{C \cap \overline{\mathcal{D}}} \mathcal{L}(\theta_1) d\mathbf{x} \quad \text{and} \quad c \int_{\overline{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x} \geq \int_{\overline{C} \cap \mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x}.$$



Example: Binomial distribution



The Neyman–Pearson Lemma (cont.)

Proof (cont.):

Thus far we have

by equality of alpha for region C and D

by definition of region C

$$\int_{\bar{C} \cap \mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x} \leq c \int_{\bar{C} \cap \mathcal{D}} \mathcal{L}(\theta_0) d\mathbf{x} = c \int_{C \cap \bar{\mathcal{D}}} \mathcal{L}(\theta_0) d\mathbf{x} \leq \int_{C \cap \bar{\mathcal{D}}} \mathcal{L}(\theta_1) d\mathbf{x}.$$

Let π be the power of the LRT, and let π' be the power of the other test. We have
 power = reject null when H_1 is true

$$\begin{aligned} \pi &= \mathbb{P}(\underline{X} \in \mathcal{C} | \theta = \theta_1) = \int_{\mathcal{C}} f(\underline{x} | \theta_1) d\mathbf{x} = \int_{\mathcal{C}} \mathcal{L}(\theta_1) d\mathbf{x} \\ &= \int_{C \cap \mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x} + \int_{C \cap \bar{\mathcal{D}}} \mathcal{L}(\theta_1) d\mathbf{x} \geq \int_{C \cap \mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x} + \int_{\bar{C} \cap \mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x} \\ &= \int_{\mathcal{D}} \mathcal{L}(\theta_1) d\mathbf{x} = \int_{\mathcal{D}} f(\underline{x} | \theta_1) d\mathbf{x} = \mathbb{P}(\underline{X} \in \mathcal{D} | \theta = \theta_1) = \pi'. \end{aligned}$$

prove that power of LRT > power of any other test



Example: Bernoulli data

Example

Suppose that we wish to test $\mathcal{H}_0 : p = p_0$ vs. $\mathcal{H}_1 : p = p_1$ (for $p_1 > p_0$), based on sequence X_1, \dots, X_n of Bernoulli trials with an unknown probability of success p . Find an MP test at level α . **should be does not exceed alpha.**

Solution: **just use LRT**

Here

note do not use pdf of binomial distribution, just product of bernoulli pdf

$$\begin{aligned}
 \lambda(\underline{x}) &= \frac{\mathcal{L}(p_1)}{\mathcal{L}(p_0)} = \frac{p_1^{\sum x_i} (1-p_1)^{n-\sum x_i}}{p_0^{\sum x_i} (1-p_0)^{n-\sum x_i}} = \left(\frac{p_1}{p_0}\right)^{\sum x_i} \left(\frac{1-p_1}{1-p_0}\right)^{n-\sum x_i} \\
 &= \underbrace{\left(\frac{p_1}{p_0}\right)}_{>1} \cdot \underbrace{\left(\frac{1-p_1}{1-p_0}\right)}_{>1}^{\sum x_i} \underbrace{\left(\frac{1-p_1}{1-p_0}\right)^n}_{<1} = a^{\sum x_i} b,
 \end{aligned}$$

where $a > 1$ and $b > 0$.



Example: Bernoulli data (cont.)

Solution (cont.):

$$\lambda(\underline{x}) = a^{\sum x_i} b, \text{ where } a > 1 \text{ and } b > 0.$$

monotonic increasing function

Now,

$$\lambda(\underline{x}) \geq c \iff a^{\sum x_i} b \geq c \iff a^{\sum x_i} \geq c_1 \iff \sum x_i \geq c_2,$$

hence, the rejection region of the LRT is

$$\mathcal{C} = \{\underline{x} : \lambda(\underline{x}) \geq c\} = \left\{ \underline{x} : \sum x_i \geq c_2 \right\}.$$

To find c_2 , recall that $\sum_{i=1}^n X_i \sim \text{Binom}(n, p)$, hence

note here how alpha is the upper bound on possible size of test

$$\alpha \geq \mathbb{P}(\underline{X} \in \mathcal{C} | p = p_0) = \mathbb{P}\left(\sum_{i=1}^n X_i \geq c_2 \mid p = p_0\right)$$

rejecting null when null is true

note this is a problem for discrete distribution, where can't really find size of test to be exact alpha

$$= 1 - \mathbb{P}\left(\sum_{i=1}^n X_i < c_2 \mid p = p_0\right) = 1 - \sum_{k=0}^{\lfloor c_2 \rfloor - 1} \binom{n}{k} p_0^k (1 - p_0)^{n-k}.$$

sum < c SAME AS sum ≤ c - 1

cdf of binomial



Example: Bernoulli data (cont.)

Solution (cont.): take smallest c_2 such that the output $\leq \alpha$

So c_2 is the smallest integer satisfying

$$\sum_{k=0}^{c_2-1} \binom{n}{k} p_0^k (1-p_0)^{n-k} \geq 1 - \alpha.$$

For example, to test $\mathcal{H}_0 : p = 0.2$ vs. $\mathcal{H}_1 : p = 0.3$ at a 5% level with $n = 50$, we need to look for the smallest c_2 satisfying

$$\sum_{k=0}^{c_2-1} \binom{50}{k} 0.2^k 0.8^{50-k} \geq 0.95.$$

```
> probs <- pbinom(c(0:50), size=50, prob=.2)
```

```
> (c2 <- min(which(probs > .95)))
```

```
[1] 16
```

assume that c_2 is integer, because sum of int is int, no point in setting a threshold that is not int

and the rejection region in this case is $\mathcal{C} = \left\{ \underline{x} : \sum_{i=1}^{50} x_i \geq 16 \right\}$.

so reject null if sum of x is ≥ 16 .

gives significant level of $\alpha = 0.05$ with smallest beta



Example: Bernoulli data (cont.)

Solution (cont.):

a conservative test; not exactly 5% because pdf is discrete

- ★ CAUTION: the actual significance level of this test is actually not 5%, but

$$\text{alpha} = \mathbb{P}(\underline{X} \in \mathcal{C} | p = 0.2) = \mathbb{P}\left(\sum_{i=1}^{50} X_i \geq 16 \middle| p = 0.2\right) = 0.031$$

(but if we chose the cutoff to be 15 instead, it would be 0.061).

- ★ The Neyman-Pearson Lemma states that no test with greater power exists at the 0.031 level, but there could be a test with a significance level that is closer to 5% and a greater power. **because pearson lemma states MP test at exactly alpha**
- For a large n , $\text{Binom}(n, p) \overset{\text{CLT}}{\approx} \mathcal{N}(np, np(1-p))$. We can then calculate

$$\alpha = \mathbb{P}(\underline{X} \in \mathcal{C} | p = p_0) = \mathbb{P}\left(\sum_{i=1}^n X_i \geq c_2 \middle| p = p_0\right) \approx 1 - \Phi\left(\frac{c_2 - np_0}{\sqrt{np_0(1-p_0)}}\right)$$

$$\implies c_2 = np_0 + z_{1-\alpha} \sqrt{np_0(1-p_0)}$$



Example: Bernoulli data (cont.)

Solution (cont.):

- The rejection region based on the large sample approximation is

$$C' = \left\{ \underline{x} : \sum_{i=1}^n x_i \geq np_0 + z_{1-\alpha} \sqrt{np_0(1-p_0)} \right\}$$

- For $n = 50$, $\alpha = 0.05$ and $p_0 = 0.2$ we get

$$\begin{aligned} C' &= \left\{ \underline{x} : \sum_{i=1}^{50} x_i \geq 50 \times 0.2 + 1.645 \sqrt{50 \times 0.2 \times 0.8} \right\} \\ &= \left\{ \underline{x} : \sum_{i=1}^{50} x_i \geq 14.65 \right\} = \left\{ \underline{x} : \sum_{i=1}^{50} x_i \geq 15 \right\}. \end{aligned}$$

in this case, normal approximation is continuous hence able to find c such that $P(\text{Type I error}) = \alpha$. But still since binomial is discrete, c gives to a natural number, whose actual significance level is not 5%, in fact in this case it is 6.1%