

MAT237 notes

Mark Wang

April 15, 2017

Contents

[11pt]article
 marginnote caption subcaption mathtools amsmath amssymb amsthm pgfplots relsize
 bbm [noline, lined, boxed, titled, noend]algorithm2e
 multicol
 listings color tikz
 tikz-qtrees forest enumitem graphicx
 [a4paper, total=6in, 8in]geometry hyperref
 Theorem Theorem[section] Corollary Corollary[theorem] Proposition Proposition Lemma
 Lemma
 Definition Definition[section]
 Example
 Remark Note Case
 *arg min arg min * arg maxarg max

1 Set and Notation

Definition 1.1. Let S be a set and choose 2 sets $A, B \subseteq S$. The **union** of A and B is

$$A \cup B = \{x \in S \mid x \in A \vee x \in B\}$$

$$\bigcup_{i \in I} A_i = \{x \in S \mid \exists i \in I, x \in A_i\}$$

Definition 1.2. Let S be a set and choose 2 sets $A, B \subseteq S$. The **intersection** of A and B is

$$A \cap B = \{x \in S \mid x \in A \wedge x \in B\}$$

$$\bigcap_{i \in I} A_i = \{x \in S \mid \forall i \in I, x \in A_i\}$$

Definition 1.3. If $A \subseteq S$, then the **complement** of A with respect to S is all elements which are not in A , that is

$$A^c = \{x \in S : x \notin A\}$$

Definition 1.4. Given 2 sets A, B , a **function** $f : A \rightarrow B$ is a map which assigns to every point in A a unique point of B , that is

$$f : a \mapsto f(a), \text{ where } a \in A, f(a) \in B$$

Definition 1.5. Let $f : A \rightarrow B$ be a function.

1. If $U \subseteq A$, then we define the **image** of U to be

$$f(U) = \{y \in B : \exists x \in U, f(x) = y\} = \{f(x) : x \in U\}$$

2. If $V \subseteq B$ we define the **pre-image** of V to be

$$f^{-1}(V) = \{x \in A : f(x) \in V\}$$

Remark. Note U, V are sets, not variable.

Definition 1.6. Let $f : A \rightarrow B$ be a function. We say that

1. f is **injective** if whenever $f(x) = f(y)$ then $x = y$
2. f is **surjective** if for every $y \in B$ there exists $x \in A$ such that $f(x) = y$
3. f is **bijective** if f is both injective and surjective

Remark. Testing injectivity by using the horizontal line test in \mathbb{R}^2 : An injective function is one whose graph that never intersect any horizontal line twice. Test surjectivity by ensuring that every horizontal line in the domain is crossed at least once by the graph.