

Structure of pHs RLC

Antoine Falaize¹ and Thomas Helie²

¹Project-team S3*, , STMS, IRCAM-CNRS-UPMC (UMR 9912), , 1 Place
Igor-Stravinsky, 75004 Paris, France

²Project-team S3[†], , STMS, IRCAM-CNRS-UPMC (UMR 9912), , 1 Place
Igor-Stravinsky, 75004 Paris, France

November 12, 2016

1 System netlist

line	label	dictionary	component	nodes	parameters
------	-------	------------	-----------	-------	------------

$\dim(\mathbf{x}) = n_{\mathbf{x}} = 2;$
 $\dim(\mathbf{w}) = n_{\mathbf{w}} = 1;$
 $\dim(\mathbf{y}) = n_{\mathbf{y}} = 1;$
 $\dim(\mathbf{p}) = n_{\mathbf{p}} = 0;$

2 System variables

State variable $\mathbf{x} = \begin{pmatrix} x_L \\ x_C \end{pmatrix};$

Dissipation variable $\mathbf{w} = (w_R);$

Input $\mathbf{u} = (u_{\text{out}});$

Output $\mathbf{y} = (y_{\text{out}});$

*<http://s3.ircam.fr>

[†]<http://s3.ircam.fr>



Figure 1: Graph of system RLC.

3 Constitutive relations

$$\text{Hamiltonian } H(\mathbf{x}) = \frac{x_L^2}{2 \cdot L} + \frac{x_C^2}{2 \cdot C};$$

$$\text{Hamiltonian gradient } \nabla H(\mathbf{x}) = \begin{pmatrix} \frac{x_L}{L} \\ \frac{x_C}{C} \end{pmatrix};$$

Dissipation function $\mathbf{z}(\mathbf{w}) = (\mathbf{R} \cdot w_{\mathbf{R}})$;
Jacobian of dissipation function $\mathcal{J}_{\mathbf{z}}(\mathbf{w}) = (\mathbf{R})$;

4 System parameters

4.1 Constant

parameter	value (SI)
C :	2e-09
R :	1000.0
L :	0.05

$$\mathbf{M} = \begin{pmatrix} 0 & -1 & -1 & -1 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix};$$

$$\mathbf{J} = \begin{pmatrix} 0 & -1.0 & -1.0 & -1.0 \\ 1.0 & 0 & 0 & 0 \\ 1.0 & 0 & 0 & 0 \\ 1.0 & 0 & 0 & 0 \end{pmatrix};$$

$$\mathbf{R} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix};$$