

MAS242 해석학 II

Notes

한승우

November 9, 2023

CONTENTS

CHAPTER	DIFFERENTIATION	PAGE 2
	1.1 Higher order partial derivatives	2
	1.2 Extreme Values of differentiable Functions	3
CHAPTER	INVERSE FUNCTION THEOREM	PAGE 5
	2.1 Jacobian	5
	2.2 The Inverse Function Theorem	5
	2.3 Implicit Function Theorem	7
	2.4 Applications of IMFT: Lagrange's Method	9
CHAPTER	SERIES OF VECTORS	PAGE 11
	3.1 Preliminaries	11
	3.2 Finite Dimensional Banach Spaces	12
	3.3 Conditional Convergence	13
	3.4 The Cauchy Product	14
	3.5 Series on Infinite Dimensional Banach Spaces	15
CHAPTER	ANALYSIS FOR SERIES FUNCTIONS	PAGE 17
	4.1 Calculus of Series Functions	17

Chapter 1

Differentiation

1.1 Higher order partial derivatives

Definition 1.1.1

Given $f : U \rightarrow \mathbb{R}$ where U is an open set in \mathbb{R}^m , define $\partial_{ij} \triangleq \partial_i(\partial_j f)(x)$ for each $i, j \in [m]$ to be *2nd order partial derivatives*. Any higher order partial derivatives can be defined inductively.

Definition 1.1.2: C^k -regularity

$f : U \rightarrow \mathbb{R}$ is C^k -regular if all partial derivatives up to order k and they are continuous.

Theorem 1.1.1

$f : U(\subseteq \mathbb{R}^2) \rightarrow \mathbb{R}$ is C^2 at a point $c \in U$, i.e., $\exists \delta > 0$, f is C^2 in $B_\delta(c)$. Then, $\partial_{12}f(c) = \partial_{21}f(c)$.

Proof. Let $|h| < \delta$. Define $A(h) \triangleq f(c_1 + h_1, c_2 + h_2) - f(c_1 + h_1, c_2) - f(c_1, c_2 + h_2) + f(c_1, c_2)$. Define $u(x_1) \triangleq f(x_1, c_2 + h_2) - f(x_1, c_2)$ and $v(x_2) \triangleq f(c_1 + h_1, x_2) - f(c_1, x_2)$. Note that u and v are differentiable.

Then, $A(h) = u(c_1 + h_1) - u(c_1)$ and $A(h) = v(c_2 + h) - v(c_2)$. By MVT, $\exists c_1^* \in (c_1, c_1 + h_1)$ and $c_2^* \in (c_2, c_2 + h_2)$ s.t. $A(h) = u'(c_1^*)h_1 = h_1(\partial_1 f(c_1^*, c_2 + h) - \partial_1 f(c_1^*, c_2)) = h_1 h_2 \partial_{21}f(c_1^*, c_2^*)$

Similarly, $\exists c_1^{**}, c_2^{**}$ such that $A(h) = h_1 h_2 \partial_{12}f(c_1^{**}, c_2^{**})$. $\partial_{21}f(c_1^*, c_2^*) = \partial_{12}f(c_1^{**}, c_2^{**})$. Hence, as $|h| \rightarrow 0$, due to the continuity, $\partial_{21}(c) = \partial_{12}(c)$. \square

Corollary 1.1.1

Suppose $f : U(\subseteq \mathbb{R}^m) \rightarrow \mathbb{R}$ is C^k at $c \in U$. Then $\partial_{j_1 j_2 \dots j_k} f(c) = \partial_{j'_1 j'_2 \dots j'_k} f(c)$ where $j'_1 \dots$ are a permutation of $j_1 \dots$.

1.2 Extreme Values of differentiable Functions

Definition 1.2.1: Hessian

Let $f : U(\subseteq \mathbb{R}^m) \rightarrow \mathbb{R}$ be C_2 in U . Suppose $p \in U$ is a critical point of f , i.e., $\nabla f(p) = 0$. Define

$$\mathcal{H}f(x) \triangleq \begin{pmatrix} \partial_{11}f(x) & \partial_{21}f(x) & \cdots & \partial_{m1}f(x) \\ \partial_{12}f(x) & \partial_{22}f(x) & \cdots & \partial_{m2}f(x) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{1m}f(x) & \partial_{2m}f(x) & \cdots & \partial_{mm}f(x) \end{pmatrix}.$$

(Sometimes $\mathcal{H}f(x) = D^2f(x)$.)

Define $D(x) = \det \mathcal{H}f(x)$. (Note that $\mathcal{H}f(x)$ is symmetric when f is C^2 by the theorem above.)

Theorem 1.2.1 2nd-order derivative test for two variable functions.

When $m = 2$ and f is C^2 , a critical point p is

- a local maximum if $D(p) > 0$ and $\partial_{11}f(p) > 0$ (or $\partial_{22}f(p) > 0$).
- a local minimum if $D(p) > 0$ and $\partial_{11}f(p) < 0$ (or $\partial_{22}f(p) < 0$).
- a saddle point if $D(p) < 0$.

The test fails when $D(p) = 0$.

Proof. Given a unit vector $\mathbf{u} = (u_1, u_2) \in \mathbb{R}^2$, $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} = u_1\partial_1f + u_2\partial_2f$, and thus

$$D_{\mathbf{u}}^2f = (u_1\partial_1 + u_2\partial_2)(u_1\partial_1f + u_2\partial_2f) = u_1^2\partial_{11}f + u_1u_2(2\partial_{12}f) + u_2^2\partial_{22}f.$$

WLOG, $u_1 \neq 0$. Set $z = u_2/u_1$. Then,

$$D_{\mathbf{u}}^2f(p) = u_1^2(\partial_{11}f(p) + 2\partial_{12}f(p)z + \partial_{22}f(p)z^2).$$

Note that, if $D(p) > 0$, $D_{\mathbf{u}}^2f(p)$ has no real root.

- If $D(p) > 0$ and $\partial_{11}f(p) < 0$, Then, $D^2\mathbf{u} < 0$ for all unit vector \mathbf{u} .
- If $D(p) > 0$ and $\partial_{11}f(p) > 0$, Then, $D^2\mathbf{u} > 0$ for all unit vector \mathbf{u} .
- If $D(p) < 0$, $D_{\mathbf{u}}^2f(p)$ has different signs depending on \mathbf{u} .

For general m ?

$$D_{\mathbf{u}}(D_{\mathbf{u}}f) = D_{\mathbf{u}} \sum_{j=1}^m \partial_j f u_j = \sum_{j=1}^m ((\nabla \partial_j f) \cdot \mathbf{u}) u_j = \sum_{j=1}^m \sum_{k=1}^m u_k u_j \partial_{kj} f.$$

Hence,

$$D_{\mathbf{u}}^2f(p) = \mathbf{u}^T \cdot D^2f(p) \cdot \mathbf{u}$$

Since $D^2f(p)$ is symmetric, its eigenvalues $\lambda_1, \dots, \lambda_m$ exists and they are real numbers. Also, there exists an $m \times m$ orthogonal matrix \mathcal{O} such that $D^2f(p) = \mathcal{O}\Lambda(p)\mathcal{O}^T$ where $\Lambda(p)$ is the diagonal matrix with entries are the eigenvalues.

Then, we can write $D_{\mathbf{u}}^2f(p) = \mathbf{u}\mathcal{O}\Lambda(p)\mathcal{O}^T\mathbf{u}^T = (\mathbf{u}\mathcal{O})\Lambda(p) = (\mathbf{u}\mathcal{O})^T$. Since \mathcal{O} is orthogonal, $\mathbf{u}\mathcal{O}$ is another arbitrary unit vector. \square

Theorem 1.2.2 Generalized 2nd order partial derivatives test

When f is C^2 , a critical point p is

- a local maximum if all eigenvalues of $D^2f(p)$ are negative.

- a local minimum if all eigenvalues of $D^2f(p)$ are positive.
 - a saddle point if there are both negative eigenvalues and positive eigenvalues.
- The test fails when there are zero eigenvalues.

Chapter 2

Inverse Function Theorem

2.1 Jacobian

Definition 2.1.1: Jacobian

Let $\mathbf{f}: U(\subseteq \mathbb{R}^m) \rightarrow \mathbb{R}^n$ be differentiable. The function $J_{\mathbf{f}}: U \rightarrow \mathbb{R}$ defined by

$$J_{\mathbf{f}}(\mathbf{x}) = \det \begin{bmatrix} \partial_1 f_1(\mathbf{x}) & \cdots & \partial_n f_1(\mathbf{x}) \\ \vdots & \ddots & \vdots \\ \partial_1 f_n(\mathbf{x}) & \cdots & \partial_n f_n(\mathbf{x}) \end{bmatrix}$$

is called the *Jacobian* of \mathbf{f} at \mathbf{x} .

Lemma 2.1.1

If $f: V(\subseteq \mathbb{R}^n) \rightarrow \mathbb{R}$ and $\mathbf{g}: U \rightarrow V$ are differentiable, then

$$J_{f \circ \mathbf{g}}(\mathbf{x}) = J_f(\mathbf{g}(\mathbf{x})) \cdot J_{\mathbf{g}}(\mathbf{x}).$$

Note:-

The linear mapping $df(c)$ is invertible if and only if $J_{\mathbf{f}}(c)$ is nonzero.

2.2 The Inverse Function Theorem

Lemma 2.2.1 Contraction Mapping Principle

Let (X, d) be a complete metric space. Let $\varphi: X \rightarrow X$. Suppose that there exists $M \in [0, 1)$ such that $d(\varphi(x_1), \varphi(x_2)) \leq M d(x_1, x_2)$. (We call it a *contraction mapping*.) Then, there uniquely exists $x_* \in X$ such that $\varphi(x_*) = x_*$.

Proof. Fix any $x_0 \in X$. Since $\{x_j\}_{j \in \mathbb{Z}_+}$, where $x_j = \varphi(x_{j-1})$ for each $j \in \mathbb{Z}_+$, is continuous. It converges to some x_* . As φ is continuous, we have $\varphi(x_*) = x_*$. The uniqueness follows trivially. \square

Note:-

- For each $v \in \mathbb{R}^n \setminus \{0\}$, $|Av| = |v| \cdot |A \frac{v}{|v|}| \leq \|A\|_L \cdot |v|$. The result is trivial when $v = 0$.
- For each $u \in \mathbb{R}^n$ with $|u| = 1$, $|ABu| \leq \|A\|_L |Bu| \leq \|A\|_L \|B\|_L$. Hence, $\|AB\|_L = \|A\|_L \|B\|_L$.
- Given invertible $A \in L(\mathbb{R}^n, \mathbb{R}^n)$, $A^{-1}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is linear. Moreover, $\|A\|_L > 0$.

Lemma 2.2.2

Given two linear mappings $A, B: \mathbb{R}^n \rightarrow \mathbb{R}^n$ with invertibility of A ,

$$\|A - B\|_L \cdot \|A^{-1}\|_L < 1 \implies B \text{ is invertible.}$$

Proof. Let $\|A^{-1}\|_L = 1/\alpha$ and $\|B - A\|_L = \beta$ so that $\beta < \alpha$. Then, for every $\mathbf{x} \in \mathbb{R}^n$,

$$\begin{aligned} \alpha|\mathbf{x}| &= \alpha|A^{-1}A\mathbf{x}| \leq \alpha\|A^{-1}\| \cdot |A\mathbf{x}| \\ &= |A\mathbf{x}| \leq |(A - B)\mathbf{x}| + |B\mathbf{x}| \leq \beta|\mathbf{x}| + |B\mathbf{x}|; \end{aligned}$$

hence $(\alpha - \beta)|\mathbf{x}| \leq |B\mathbf{x}|$ where $\mathbf{x} \in \mathbb{R}^n$ is arbitrary. As $\alpha > \beta$, it holds that $B\mathbf{x} = 0 \implies \mathbf{x} = 0$. \square

Corollary 2.2.1

The set $\Omega \subseteq L(\mathbb{R}^n, \mathbb{R}^n)$ of invertible linear transformations is open.

Lemma 2.2.3

The mapping from Ω onto Ω defined by $A \mapsto A^{-1}$ is continuous.

Proof. Let A and B be invertible linear transformations from \mathbb{R}^n to \mathbb{R}^n . Let $\|A^{-1}\| = 1/\alpha$ and $\|B - A\|_L = \beta$. We have $(\alpha - \beta)|\mathbf{x}| \leq |B\mathbf{x}|$ by the same reasoning as in the proof of Lemma 2.2.2. Hence, the following holds.

$$\forall \mathbf{y} \in \mathbb{R}^n, (\alpha - \beta)|B^{-1}\mathbf{y}| \leq |BB^{-1}\mathbf{y}| = |\mathbf{y}|$$

This shows that $\|B^{-1}\|_L \leq (\alpha - \beta)^{-1}$.

Hence, we have

$$\|B^{-1} - A^{-1}\|_L \leq \|B^{-1}\|_L \|A - B\|_L \|A^{-1}\|_L \leq \frac{\beta}{\alpha(\alpha - \beta)}.$$

This implies that $\|B^{-1} - A^{-1}\|_L \rightarrow 0$ as $B \rightarrow A$. \square

Theorem 2.2.1 Inverse Function Theorem

Let $\mathbf{f}: E(\subseteq \mathbb{R}^n) \rightarrow \mathbb{R}^n$ be C^1 in E and $\mathbf{c} \in E$. Suppose that $J_{\mathbf{f}}(\mathbf{c}) \neq 0$. Then, the following hold.

- (i) There exists a neighborhood U of \mathbf{a} such that $\mathbf{f}|_U$ is bijective and $V \triangleq \mathbf{f}(U)$ is open.
- (ii) The inverse map of $\mathbf{f}|_U$ is C^1 in V .

Proof. Let $A \triangleq d\mathbf{f}(\mathbf{c})$. Define $\lambda \in \mathbb{R}_+$ by $2\lambda\|A^{-1}\|_L = 1$. Since $d\mathbf{f}$ is continuous, there exists a neighborhood U of \mathbf{c} such that $\|d\mathbf{f}(\mathbf{x}) - A\|_L < \lambda$ for each $\mathbf{x} \in U$.

Given a point $\mathbf{y} \in \mathbb{R}^n$, we define $\varphi(\cdot; \mathbf{y})$ by

$$\begin{aligned} \varphi(\cdot; \mathbf{y}) : B_{\delta}(\mathbf{c}) &\longrightarrow \mathbb{R}^n \\ \mathbf{x} &\longmapsto \mathbf{x} + A^{-1}(\mathbf{y} - \mathbf{f}(\mathbf{x})) \end{aligned}$$

Note that \mathbf{x} is a fixed point of $\varphi(\cdot; \mathbf{y})$ if and only if $A^{-1}(\mathbf{y} - \mathbf{f}(\mathbf{x})) = 0$, i.e., $\mathbf{y} = \mathbf{f}(\mathbf{x})$. Note also that φ is differentiable and $d\varphi(\mathbf{x}; \mathbf{y}) = \text{Id} - A^{-1} d\mathbf{f}(\mathbf{x}) = A^{-1}(A - d\mathbf{f}(\mathbf{x}))$ for each $\mathbf{x} \in U$.

Hence, for all $\mathbf{x} \in U$,

$$\|d\varphi(\mathbf{x}; \mathbf{y})\|_L = \|A^{-1}(A - d\mathbf{f}(\mathbf{x}))\|_L \leq \|A^{-1}\|_L \cdot \|A - d\mathbf{f}(\mathbf{x})\|_L < 1/(2\lambda) \cdot \lambda = 1/2.$$

Thus, MVT gives

$$|\varphi(\mathbf{x}_1; \mathbf{y}) - \varphi(\mathbf{x}_2; \mathbf{y})| \leq \frac{1}{2} |\mathbf{x}_1 - \mathbf{x}_2|$$

whenever $\mathbf{x}_1, \mathbf{x}_2 \in U$. Note that this implies there is at most one fixed point of $\varphi(\cdot; \mathbf{y})$ in U , i.e., $\mathbf{f}|_U$ is bijective.

Now, we shall show that $V = \mathbf{f}(U)$ is open. Take any $\mathbf{y}_0 \in V$. There (uniquely) exists $\mathbf{x}_0 \in U$ such that $\mathbf{y}_0 = \mathbf{f}(\mathbf{x}_0)$. Fix any $r \in \mathbb{R}_+$ such that $\bar{B} \subseteq U$ where $B = B_r(\mathbf{x}_0)$. Take any $\mathbf{y} \in B_{\lambda r}(\mathbf{y}_0)$. Then,

$$|\varphi(\mathbf{x}_0; \mathbf{y}) - \mathbf{x}_0| = |A^{-1}(\mathbf{y} - \mathbf{y}_0)| < \|A^{-1}\|_L \lambda r = \frac{r}{2}.$$

Moreover, for any $\mathbf{x} \in \bar{B}$,

$$|\varphi(\mathbf{x}; \mathbf{y}) - \mathbf{x}_0| \leq |\varphi(\mathbf{x}; \mathbf{y}) - \varphi(\mathbf{x}_0; \mathbf{y})| + |\varphi(\mathbf{x}_0; \mathbf{y}) - \mathbf{x}_0| \leq \frac{1}{2} |\mathbf{x} - \mathbf{x}_0| + \frac{r}{2} < r.$$

This directly implies that $\varphi(\bar{B}; \mathbf{y}) \subseteq B \subseteq \bar{B}$. Hence, $\varphi(\cdot, \mathbf{y})$ is a contraction mapping on a complete metric space \bar{B} . By Lemma 2.2.1, there exists a fixed point $\mathbf{x} \in \bar{B}$, which satisfies $\mathbf{y} = \mathbf{f}(\mathbf{x})$. Thus, $\mathbf{y} \in \mathbf{f}(\bar{B}) \subseteq \mathbf{f}(U) = V$. Hence, $B \subseteq V$, V is open. This proves (i).

Now, let $\mathbf{g}: V \rightarrow U$ be the local inverse map of $\mathbf{f}|_U$. Take any $\mathbf{y} \in V$ and $\mathbf{y} + \mathbf{k} \in V$. There are unique $\mathbf{x} \in U$ and $\mathbf{x} + \mathbf{h} \in U$ such that $\mathbf{y} = \mathbf{f}(\mathbf{x})$ and $\mathbf{y} + \mathbf{k} = \mathbf{f}(\mathbf{x} + \mathbf{h})$. Then, we have

$$\varphi(\mathbf{x} + \mathbf{h}; \mathbf{y}) - \varphi(\mathbf{x}; \mathbf{y}) = \mathbf{h} + A^{-1}(\mathbf{f}(\mathbf{x}) - \mathbf{f}(\mathbf{x} + \mathbf{h})) = \mathbf{h} - A^{-1}\mathbf{k},$$

which implies $|\mathbf{h} - A^{-1}\mathbf{k}| \leq |\mathbf{h}|/2$. Hence, $|A^{-1}\mathbf{k}| \geq |\mathbf{h}|/2$ is obtained by the triangle inequality; $|\mathbf{h}| \leq 2\|A^{-1}\|_L |\mathbf{k}| = \lambda^{-1} |\mathbf{k}|$.

Then, since $\|\mathbf{d}\mathbf{f}(\mathbf{x}) - A\|_L \|A^{-1}\|_L < \lambda \cdot 1/(2\lambda) = 1/2$, Lemma 2.2.2 implies that $\mathbf{d}\mathbf{f}(\mathbf{x})$ is invertible. Let $T \triangleq \mathbf{d}\mathbf{f}(\mathbf{x})$. Then, we have

$$\mathbf{g}(\mathbf{y} + \mathbf{k}) - \mathbf{g}(\mathbf{y}) - T^{-1}\mathbf{k} = \mathbf{h} - T^{-1}\mathbf{k} = -T^{-1}(\mathbf{f}(\mathbf{x} + \mathbf{h}) - \mathbf{f}(\mathbf{x}) - T\mathbf{h}),$$

and thus

$$\frac{|\mathbf{g}(\mathbf{y} + \mathbf{k}) - \mathbf{g}(\mathbf{y}) - T^{-1}\mathbf{k}|}{|\mathbf{k}|} \leq \frac{\|T^{-1}\|_L}{\lambda} \cdot \frac{|\mathbf{f}(\mathbf{x} + \mathbf{h}) - \mathbf{f}(\mathbf{x}) - T\mathbf{h}|}{|\mathbf{h}|}.$$

The equation implies that \mathbf{g} is differentiable on V , and that $\mathbf{d}\mathbf{g}(\mathbf{y}) = T^{-1} = \mathbf{d}\mathbf{f}(\mathbf{g}(\mathbf{y}))^{-1}$. Since $\mathbf{d}\mathbf{g}$ is a composition of continuous functions, $\mathbf{d}\mathbf{g}$ itself is continuous. \square

Corollary 2.2.2

Let $\mathbf{f}: E(\subseteq \mathbb{R}^n) \rightarrow \mathbb{R}^n$ be C^1 in E and $J_{\mathbf{f}}(\mathbf{x}) \neq 0$ for all $\mathbf{x} \in E$. Then, for every open set $W \subseteq E$, $\mathbf{f}(W)$ is open.

Proof. This directly follows from (i) of Theorem 2.2.1. \square

2.3 Implicit Function Theorem

Definition 2.3.1

- If $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $\mathbf{y} = (y_1, \dots, y_m) \in \mathbb{R}^m$, let us write (\mathbf{x}, \mathbf{y}) for the point $(x_1, \dots, x_n, y_1, \dots, y_m) \in \mathbb{R}^{n+m}$.
- Every $A \in L(\mathbb{R}^{n+m}, \mathbb{R}^n)$ can be split into $A_x \in L(\mathbb{R}^n, \mathbb{R}^n)$ and $A_y \in L(\mathbb{R}^m, \mathbb{R}^n)$ where $A(\mathbf{h}, \mathbf{k}) = A_x \mathbf{h} + A_y \mathbf{k}$ for each $\mathbf{h} \in \mathbb{R}^n$ and $\mathbf{k} \in \mathbb{R}^m$.

Lemma 2.3.1

If $A \in L(\mathbb{R}^{n+m}, \mathbb{R}^n)$ and if A_x is invertible, then

$$\forall \mathbf{k} \in \mathbb{R}^m, \exists! \mathbf{h} \in \mathbb{R}^n, A(\mathbf{h}, \mathbf{k}) = \mathbf{0}.$$

Proof. $A(\mathbf{h}, \mathbf{k}) = A_x \mathbf{h} + A_y \mathbf{k} = \mathbf{0}$ if and only if $\mathbf{h} = -(A_x)^{-1} A_y \mathbf{k}$. □

Theorem 2.3.1 Implicit Function Theorem

Let $\mathbf{f}: E \rightarrow \mathbb{R}^n$ be a C^1 mapping where E is an open set in \mathbb{R}^{n+m} . Let $(\mathbf{a}, \mathbf{b}) \in E$ satisfy $\mathbf{f}(\mathbf{a}, \mathbf{b}) = \mathbf{0}$. Let $A = d\mathbf{f}(\mathbf{a}, \mathbf{b})$ and suppose A_x is invertible. Then, there exist open sets $U \subseteq \mathbb{R}^{n+m}$ and $W \subseteq \mathbb{R}^m$ that satisfy the following.

- (i) $(\mathbf{a}, \mathbf{b}) \in U$ and $\mathbf{b} \in W$.
- (ii) $\forall \mathbf{y} \in W, \exists! \mathbf{x} \in \mathbb{R}^n, (\mathbf{x}, \mathbf{y}) \in U \wedge \mathbf{f}(\mathbf{x}, \mathbf{y}) = \mathbf{0}$.
- (iii) If the unique \mathbf{x} in (ii) is denoted by $\mathbf{g}(\mathbf{y})$, then $\mathbf{g}: W \rightarrow \mathbb{R}^n$ is C^1 on W .
- (iv) Moreover, $d\mathbf{g}(\mathbf{b}) = -(A_x)^{-1} A_y$.

Proof. Define $\mathbf{F}: E \rightarrow \mathbb{R}^{n+m}$ by $\mathbf{F}(\mathbf{x}, \mathbf{y}) \triangleq (\mathbf{f}(\mathbf{x}, \mathbf{y}), \mathbf{y})$. Then, \mathbf{F} is C^1 . Since $\mathbf{f}(\mathbf{a}, \mathbf{b}) = \mathbf{0}$, if $\mathbf{r}(\mathbf{h}, \mathbf{k}) \triangleq \mathbf{f}(\mathbf{a} + \mathbf{h}, \mathbf{b} + \mathbf{k}) - A(\mathbf{h}, \mathbf{k})$, we have $\lim_{\mathbf{h}, \mathbf{k} \rightarrow \mathbf{0}} |\mathbf{r}(\mathbf{h}, \mathbf{k})| / |(\mathbf{h}, \mathbf{k})| = 0$. Hence, from

$$\mathbf{F}(\mathbf{a} + \mathbf{h}, \mathbf{b} + \mathbf{k}) - \mathbf{F}(\mathbf{a}, \mathbf{b}) = (\mathbf{f}(\mathbf{a} + \mathbf{h}, \mathbf{b} + \mathbf{k}), \mathbf{k}) = (A(\mathbf{h}, \mathbf{k}), \mathbf{k}) + (\mathbf{r}(\mathbf{h}, \mathbf{k}), \mathbf{0}),$$

it is obtained that $d\mathbf{F}(\mathbf{a}, \mathbf{b})(\mathbf{h}', \mathbf{k}') = (A(\mathbf{h}', \mathbf{k}'), \mathbf{k}')$ for each $(\mathbf{h}', \mathbf{k}') \in \mathbb{R}^{n+m}$. If $d\mathbf{F}(\mathbf{a}, \mathbf{b})(\mathbf{h}', \mathbf{k}') = \mathbf{0}$, then $\mathbf{k}' = \mathbf{0}$ and $A(\mathbf{h}', \mathbf{0}) = \mathbf{0}$; thus $\mathbf{h}' = \mathbf{0}$ as A_x is invertible. Hence, $d\mathbf{F}(\mathbf{a}, \mathbf{b})$ is invertible; Theorem 2.2.1 can be applied to \mathbf{F} at (\mathbf{a}, \mathbf{b}) .

By Theorem 2.2.1, there exists a neighborhood $U \subseteq E$ of (\mathbf{a}, \mathbf{b}) such that $\mathbf{F}|_U$ is bijective, $\mathbf{F}(U)$ is open, and its inverse is C^1 . Let $W \triangleq \{\mathbf{y} \in \mathbb{R}^m \mid (\mathbf{0}, \mathbf{y}) \in \mathbf{F}(U)\}$. W is open as $\mathbf{F}(U)$ is open. Noting that $\mathbf{b} \in W$, we finish the proof for (i).

Take any $\mathbf{y} \in W$. Then, there exists $(\mathbf{x}, \mathbf{y}) \in U$ such that $\mathbf{F}(\mathbf{x}, \mathbf{y}) = (\mathbf{0}, \mathbf{y})$; thus $\mathbf{f}(\mathbf{x}, \mathbf{y}) = \mathbf{0}$. If \mathbf{x}, \mathbf{x}' are two such point corresponding to \mathbf{y} , then

$$\mathbf{F}(\mathbf{x}', \mathbf{y}) = (\mathbf{f}(\mathbf{x}', \mathbf{y}), \mathbf{y}) = (\mathbf{0}, \mathbf{y}) = (\mathbf{f}(\mathbf{x}, \mathbf{y}), \mathbf{y}) = \mathbf{F}(\mathbf{x}, \mathbf{y}).$$

However, as \mathbf{F} being injective, $\mathbf{x} = \mathbf{x}'$. This proves (ii).

Let $V \triangleq \mathbf{F}(U)$. Let $\mathbf{G}: V \rightarrow U$ be the inverse of \mathbf{F} , which is C^1 by Theorem 2.2.1. Hence, for each $\mathbf{y} \in W$, from $\mathbf{F}(\mathbf{g}(\mathbf{y}), \mathbf{y}) = (\mathbf{0}, \mathbf{y})$, we have $(\mathbf{g}(\mathbf{y}), \mathbf{y}) = \mathbf{G}(\mathbf{0}, \mathbf{y})$. This directly shows that \mathbf{g} is C^1 as well. This proves (iii).

Let $\Phi: W \rightarrow U$ be defined by $\Phi(\mathbf{y}) = \mathbf{G}(\mathbf{0}, \mathbf{y}) = (\mathbf{g}(\mathbf{y}), \mathbf{y})$, which is C^1 , indeed. Then, $d\Phi(\mathbf{y}) = (d\mathbf{g}(\mathbf{y}), I_m)$. Differentiating both sides of the equality $\mathbf{f}(\Phi(\mathbf{y})) = \mathbf{0}$, we get

$$d\mathbf{f}(\Phi(\mathbf{y})) d\Phi(\mathbf{y}) = \mathbf{0}.$$

Putting $\mathbf{y} := \mathbf{b}$, as $\Phi(\mathbf{b}) = (\mathbf{a}, \mathbf{b})$, we get $A d\Phi(\mathbf{b}) = \mathbf{0}$, or

$$A_x d\mathbf{g}(\mathbf{b}) + A_y = \mathbf{0},$$

i.e., $d\mathbf{g}(\mathbf{b}) = -(A_x)^{-1} A_y$. □

Definition 2.3.2: C^1 -norm

Suppose $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ is C^1 . Then,

$$\begin{aligned}\|\varphi\|_{C^0(\bar{\Omega})} &\triangleq \sup_{\mathbf{x} \in \bar{\Omega}} |\varphi(\mathbf{x})| \\ \|\varphi\|_{C^1(\bar{\Omega})} &\triangleq \|\varphi\|_{C^0(\bar{\Omega})} + \sum_{j=1}^n \|\partial_j \varphi\|_{C^0(\bar{\Omega})}.\end{aligned}$$

This is only for Example 2.3.1.

Example 2.3.1 (Level Sets)

Define $\Omega \triangleq \{(x_1, x_2) \in \mathbb{R}^2 \mid |x_2| \leq 1\}$. Given two constants, $a, b \in \mathbb{R}$ with $a < b$, define $\bar{\varphi}(x_1, x_2) = ax_1$ and $\bar{\psi}(x_1, x_2) = bx_1$. Then, $\Gamma_0 = \{\mathbf{x} \in \Omega \mid \bar{\varphi}(\mathbf{x}) - \bar{\psi}(\mathbf{x}) = 0\} = \{\mathbf{x} \in \Omega \mid x_1 = 0\}$.

Suppose that $\varphi, \psi : \Omega \rightarrow \mathbb{R}$ satisfy

$$\|\varphi - \bar{\varphi}\|_{C^1(\bar{\Omega})} + \|\psi - \bar{\psi}\|_{C^1(\bar{\Omega})} \leq \frac{1}{4}|a - b|.$$

Then, what would be the expression for $\Gamma = \{\mathbf{x} \in \Omega \mid \varphi(\mathbf{x}) - \psi(\mathbf{x}) = 0\}$?

Observe that $(\varphi - \psi) = (\varphi - \bar{\varphi}) + (\bar{\varphi} - \bar{\psi}) + (\bar{\psi} - \psi)$ and thus $|(\varphi - \psi)(x_1, x_2) - (a - b)x_1| \leq |a - b|/4$. This implies $\lim_{x_1 \rightarrow \pm\infty} (\varphi - \psi)(x_1, x_2) = \mp\infty$. Hence, for every $x_2 \in [-1, 1]$, there exists $x_1^* \in \mathbb{R}$ such that $(\varphi - \psi)(x_1^*, x_2) = 0$.

Moreover, $\partial_1(\varphi - \psi) = \partial_1(\varphi - \bar{\varphi}) + (a - b) + \partial_1(\bar{\psi} - \psi)$, and thus $|\partial_1(\varphi - \psi)| \geq \frac{3}{4}|a - b| > 0$. Hence, the x_1^* in the previous paragraph is unique. This means that $\Gamma = \{(f(x_2), x_2) \mid x_2 \in \mathbb{R}\}$ for some f .

$(\varphi - \psi)(f(x_2), x_2) - (\bar{\varphi} - \bar{\psi})(f(x_2), x_2) = -(\bar{\varphi} - \bar{\psi})(f(x_2), x_2) = (b - a)f(x_2)$. Hence,

$$f(x_2) = \frac{(\varphi - \bar{\varphi})(f(x_2), x_2) - (\psi - \bar{\psi})(f(x_2), x_2)}{b - a}.$$

This is the implicit representation of f . Moreover, $|f(x_2)| = \frac{|b - a|/4}{|b - a|} = 1/4$.

2.4 Applications of IMFT: Lagrange's Method

Theorem 2.4.1 Optimization Under Multiple Constraints

Let $f, g_1, g_2, \dots, g_k : E \rightarrow \mathbb{R}$ be C^1 where E is an open set in \mathbb{R}^n and $n > k$. Let $Z \triangleq \bigcap_{j=1}^k \{\mathbf{z} \in \mathbb{R}^n \mid g_j(\mathbf{z}) = 0\}$. Suppose $\mathbf{z}_0 \in Z$ is a local maximum point with respect to f on Z . Suppose also that

$$\Delta \triangleq \det \begin{bmatrix} \partial_1 g_1(\mathbf{z}_0) & \cdots & \partial_1 g_k(\mathbf{z}_0) \\ \vdots & \ddots & \vdots \\ \partial_k g_1(\mathbf{z}_0) & \cdots & \partial_k g_k(\mathbf{z}_0) \end{bmatrix} \neq 0.$$

Then, there exists $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ such that $\nabla f(\mathbf{z}_0) = \sum_{m=1}^k \lambda_m \nabla g_m(\mathbf{z}_0)$.

Proof. Since $\Delta \neq 0$, there exists a unique solution $(\lambda_1, \dots, \lambda_k) \in \mathbb{R}^k$ for the linear system

$$\begin{bmatrix} \partial_1 g_1(\mathbf{z}_0) & \cdots & \partial_1 g_k(\mathbf{z}_0) \\ \vdots & \ddots & \vdots \\ \partial_k g_1(\mathbf{z}_0) & \cdots & \partial_k g_k(\mathbf{z}_0) \end{bmatrix} \begin{bmatrix} \lambda_1 \\ \vdots \\ \lambda_k \end{bmatrix} = \begin{bmatrix} \partial_1 f(\mathbf{z}_0) \\ \vdots \\ \partial_k f(\mathbf{z}_0) \end{bmatrix}.$$

For each point $\mathbf{z} = (z_1, \dots, z_n) \in \mathbb{R}^n$, let $\mathbf{x} = (z_1, \dots, z_k)$ and $\mathbf{y} = (z_{k+1}, \dots, z_n)$. Let $\mathbf{z}_0 = (\mathbf{x}_0, \mathbf{y}_0)$. Let $\mathbf{g}: E \rightarrow \mathbb{R}^k$ be defined by $\mathbf{g}(\mathbf{z}) = (g_1(\mathbf{z}), \dots, g_k(\mathbf{z}))$.

Since \mathbf{g} is C^1 , $\mathbf{g}(\mathbf{z}_0) = \mathbf{0}$, and $(d\mathbf{g}(\mathbf{z}_0))_{\mathbf{x}}$ is invertible, by Theorem 2.3.1, there exists an open neighborhood $W \subseteq \mathbb{R}^{n-k}$ of \mathbf{y}_0 and a C^1 function $\mathbf{s}: W \rightarrow \mathbb{R}^k$ such that $\mathbf{g}(\mathbf{s}(\mathbf{y}), \mathbf{y}) = \mathbf{0}$ for each $\mathbf{y} \in W$. Note that $\mathbf{s}(\mathbf{y}_0) = \mathbf{x}_0$.

Define $F: W \rightarrow \mathbb{R}$ by $\mathbf{y} \mapsto f(\mathbf{s}(\mathbf{y}), \mathbf{y})$. As \mathbf{z}_0 is a local maximum point, so is \mathbf{y}_0 . Hence, $\nabla F(\mathbf{y}_0) = \mathbf{0}$. For each $j \in [k]$, define $G_j: W \rightarrow \mathbb{R}$ by $\mathbf{y} \mapsto g_j(\mathbf{s}(\mathbf{y}), \mathbf{y})$. As $(\mathbf{s}(\mathbf{y}), \mathbf{y}) \in Z$, we have $G_j = 0$ for each $j \in [k]$. Thus, $\nabla G_j(\mathbf{y}) = \mathbf{0}$.

Let $\mathbf{s} = (s_1, s_2, \dots, s_k)$ where each $s_j: W \rightarrow \mathbb{R}$. Since

$$\begin{aligned} \nabla F(\mathbf{y}) &= df(\mathbf{s}(\mathbf{y}), \mathbf{y}) d(\mathbf{s}(\mathbf{y}), \mathbf{y}) \\ &= \begin{bmatrix} \partial_1 f(\mathbf{s}(\mathbf{y}), \mathbf{y}) & \cdots & \partial_n f(\mathbf{s}(\mathbf{y}), \mathbf{y}) \end{bmatrix} \begin{bmatrix} \partial_1 s_1(\mathbf{y}) & \partial_2 s_1(\mathbf{y}) & \cdots & \partial_{n-k} s_1(\mathbf{y}) \\ \partial_1 s_2(\mathbf{y}) & \partial_2 s_2(\mathbf{y}) & \cdots & \partial_{n-k} s_2(\mathbf{y}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_1 s_k(\mathbf{y}) & \partial_2 s_k(\mathbf{y}) & \cdots & \partial_{n-k} s_k(\mathbf{y}) \\ 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}, \end{aligned}$$

$\nabla F(\mathbf{y}_0) = \mathbf{0}$ implies

$$\partial_{k+j} f(\mathbf{z}_0) + \sum_{i=1}^k \partial_i f(\mathbf{z}_0) \partial_j s_i(\mathbf{y}_0) = 0$$

for each $j \in [n-k]$. Similarly, $\nabla G_m(\mathbf{y}_0) = \mathbf{0}$ for each $m \in [k]$ implies that

$$-\lambda_m \left[\partial_{k+j} g_m(\mathbf{z}_0) + \sum_{i=1}^k \partial_i g_m(\mathbf{z}_0) \partial_j s_i(\mathbf{y}_0) \right] = 0$$

for each $j \in [n-k]$ and $m \in [k]$.

Adding the $k+1$ equations together for each $j \in [n-k]$,

$$0 = \left[\partial_{k+j} f(\mathbf{z}_0) - \sum_{m=1}^k \lambda_m \partial_{k+j} g_m(\mathbf{z}_0) \right] + \sum_{i=1}^k \left[\partial_i f(\mathbf{z}_0) - \sum_{m=1}^k \lambda_m \partial_i g_m(\mathbf{z}_0) \right] \partial_j s_i(\mathbf{y}_0).$$

By the definition of $\lambda_1, \dots, \lambda_k$, we are left with only

$$\partial_j f(\mathbf{z}_0) = \sum_{m=1}^k \lambda_m \partial_j g_m(\mathbf{z}_0)$$

for each $j \in \{k+1, \dots, n\}$. For $j \in [k]$, the same equation holds by the definition of $\lambda_1, \dots, \lambda_k$. Hence, we have $\nabla f(\mathbf{z}_0) = \sum_{m=1}^k \lambda_m \nabla g_m(\mathbf{z}_0)$. \square

Chapter 3

Series of Vectors

3.1 Preliminaries

Definition 3.1.1: Normed Vector Space

Let V be a (real/complex) vector space equipped with a norm $\|\cdot\|$, i.e., the space $(V, \|\cdot\|)$ satisfies the following properties.

- (i) $0 \in V$
- (ii) $\|\mathbf{x}\| \geq 0$ for all $x \in V$ and $\|\mathbf{x}\| = 0$ iff $\mathbf{x} = \mathbf{0}$. (*positive definiteness*)
- (iii) $\|\beta\mathbf{x}\| = |\beta| \cdot \|\mathbf{x}\|$ for all $\mathbf{x} \in V$ and $\beta \in \mathbb{R}$. (*absolute homogeneity*)
- (iv) $\|\mathbf{x}_1 + \mathbf{x}_2\| \leq \|\mathbf{x}_1\| + \|\mathbf{x}_2\|$ for all $\mathbf{x}_1, \mathbf{x}_2 \in V$. (*triangle inequality*)

Note:-

Note that $(V, \|\cdot\|)$ is naturally a metric space with the metric function $d(\mathbf{x}_1, \mathbf{x}_2) = \|\mathbf{x}_1 - \mathbf{x}_2\|$.

Definition 3.1.2: Banach Space

A normed vector space $(V, \|\cdot\|)$ is called a *Banach space* if, for every Cauchy sequence $\{\mathbf{x}_j\}_{j \in \mathbb{N}}$, there exists a unique $\mathbf{x}_* \in V$ such that $\lim_{n \rightarrow \infty} \|\mathbf{x}_n - \mathbf{x}_*\| = 0$.

Example 3.1.1

Let A be a compact subset of \mathbb{R}^n . $(V, \|\cdot\|)$ where $V = \{f : A \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ and $\|f\| = \sup_{x \in A} |f(x)|$ forms a Banach space.

Note:-

A Banach space is a normed vector space whose naturally induced metric space is complete.

Definition 3.1.3: Series

Let $(V, \|\cdot\|)$ be a normed vector space. Given a sequence $\{x_j\}_{j \in \mathbb{N}} \subseteq V$, define $S_k \triangleq \sum_{j=1}^k x_j$ for each $k \in \mathbb{N}$. Then, each S_k is called a *partial sum* of $\{x_j\}$. If $\{S_k\}_{k \in \mathbb{N}}$ converges to S_* with respect to $\|\cdot\|$, then we write

$$S_* = \sum_{j=1}^{\infty} x_j.$$

If the limit S_* exists, we symbolically say that “ $\sum_{j=1}^{\infty} x_j$ converges.”

Lemma 3.1.1

Let $(V, \|\cdot\|)$ be a normed vector space. Let $\{x_j\}_{j \in \mathbb{N}} \subseteq V$ be a sequence. If a series $\sum_{j=1}^{\infty} x_j$ converges, then $\lim_{k \rightarrow \infty} \|x_k\| = 0$.

Proof. $\{S_k\}_{k \in \mathbb{N}}$ is a Cauchy sequence. Hence, $\lim_{k \rightarrow \infty} \|x_k\| = \lim_{k \rightarrow \infty} \|S_{k+1} - S_k\| = 0$. \square

Lemma 3.1.2

Let $(V, \|\cdot\|)$ be a Banach space. Let $\{x_j\}_{j \in \mathbb{N}} \subseteq V$ be a sequence. A series $\sum_{j=1}^{\infty} x_j$ converges if and only if $\{S_k\}_{k \in \mathbb{N}}$ is Cauchy.

Proof. The definition of Banach spaces. \square

3.2 Finite Dimensional Banach Spaces

Theorem 3.2.1 Comparison Test

Given two real sequence $\{a_j\}$ and $\{b_j\}$, suppose $0 \leq a_j \leq b_j$ for all $j \geq k_0$ where $k_0 \in \mathbb{N}$ is a fixed constant. Then, if $\sum_{j=1}^{\infty} b_j$ converges, then $\sum_{j=1}^{\infty} a_j$ converges.

Proof. Let $S_k = \sum_{j=k_0}^k a_j$ and $T_k = \sum_{j=k_0}^k b_j$. Then, $0 \leq S_n - S_m = \sum_{j=m+1}^n a_j \leq \sum_{j=m+1}^n b_j = T_n - T_m$ whenever $n \geq m \geq k_0$. As $\{T_k\}_{k \in \mathbb{N}}$ is Cauchy, $\{S_k\}_{k \in \mathbb{N}}$ is Cauchy as well. As $(\mathbb{R}, \|\cdot\|)$ is a Banach space, $\sum a_j$ converges. \square

Theorem 3.2.2 Absolute Convergence Test

Let $(V, \|\cdot\|)$ be a Banach space. Let $\{x_j\}_{j \in \mathbb{N}} \subseteq V$ be a sequence. If $\sum_{j=1}^{\infty} \|x_j\|$ converges (in \mathbb{R}), then $\sum_{j=1}^{\infty} x_j$ converges.

Proof. Let $S_k = \sum_{j=1}^k x_j \in V$ and $T_k = \sum_{j=1}^k \|x_j\| \in \mathbb{R}$. Then, $\|S_n - S_m\| = \|\sum_{j=m+1}^n x_j\| \leq \sum_{j=m+1}^n \|x_j\| = T_n - T_m$ whenever $n \geq m$. As $\{T_k\}$ is Cauchy, $\{S_k\}$ is Cauchy as well. Hence, $\sum x_j$ converges. \square

Theorem 3.2.3 Summation by Parts

Let $\{a_j\}$ and $\{b_j\}$ be two real sequences. If $\sum a_j$ converges and $\{b_j\}$ is monotonic and convergent, then $\sum_{j=1}^{\infty} a_j b_j$ converges.

Proof. Let $S_k = \sum_{j=1}^k a_j b_j \in V$ and $A_k = \sum_{j=1}^k a_j \in \mathbb{R}$. ($A_0 = 0$.) Then, $S_k = \sum_{j=1}^k (A_j - A_{j-1}) b_j = \sum_{j=1}^k A_j b_j - \sum_{j=0}^k A_0 b_{j+1} + A_k b_{k+1} = A_k b_{k+1} - \sum_{j=1}^k A_j (b_{j+1} - b_j)$.
Let $T_k = \sum_{j=1}^k |A_j (b_{j+1} - b_j)|$. Then, whenever $n < m$, we have

$$0 \leq T_m - T_n \leq M \sum_{j=n+1}^m |b_{j+1} - b_j| = M |b_{m+1} - b_{n+1}| \rightarrow 0,$$

$\{T_k\}$ is Cauchy, and thus converges; $\{S_k\}$ converges as well. □

3.3 Conditional Convergence

Definition 3.3.1: Conditional Convergence

Given a real sequence $\{a_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}$, if $\sum a_j$ converges, and if $\sum |a_j|$ does not converge, then we say that $\sum a_j$ *converges conditionally*.

Theorem 3.3.1 Alternating Series Test

Let $\{a_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}$ be a real sequence. If $a_j \geq 0$ for all $j \in \mathbb{N}$, and if $\lim_{j \rightarrow \infty} a_j = 0$, then $\sum (-1)^j a_j$ converges.

Proof. MAS101. □

Example 3.3.1

$\sum (-1)^j / j$ conditionally converges.

Note:-

Given, a real sequence $\{a_j\}$, we shall use the following definition for now.

For $j \in \mathbb{N}$, define

$$a_j^+ \triangleq \frac{|a_j| + a_j}{2} = \begin{cases} a_j & \text{if } a_j \geq 0 \\ 0 & \text{if } a_j < 0 \end{cases} \quad \text{and} \quad a_j^- \triangleq \frac{|a_j| - a_j}{2} = \begin{cases} 0 & \text{if } a_j \geq 0 \\ -a_j & \text{if } a_j < 0 \end{cases}.$$

Then, $a_j^+, a_j^- \geq 0$, $|a_j| = a_j^+ + a_j^-$, and $a_j = a_j^+ - a_j^-$.

Lemma 3.3.1

Let $\{a_j\}_{j \in \mathbb{N}}$ be a real sequence.

- (i) If $\sum a_j$ converges absolutely, then both $\sum a_j^+$ and $\sum a_j^-$ converge. Moreover, $\sum a_j = \sum a_j^+ - \sum a_j^-$.
- (ii) If $\sum a_j$ converges conditionally, then both $\sum a_j^+$ and $\sum a_j^-$ diverge.

Proof.

- (i) By the definition of a_j^+ and a_j^- .
- (ii) If one of $\sum a_j^+$ or $\sum a_j^-$ converges, since $a_j = a_j^+ - a_j^-$, the other converges as well. If they both converge, as $|a_j| = a_j^+ + a_j^-$, $\sum a_j$ converges absolutely. □

Definition 3.3.2: Rearrangement of Series

Let $\phi: \mathbb{N} \rightarrow \mathbb{N}$ be bijective. Given a sequence $\{a_j\}_{j \in \mathbb{N}}$, the series $\sum a_{\phi(j)}$ is called a *rearrangement* of $\sum a_j$.

Theorem 3.3.2 Riemann's Rearrangement Theorem

Let $\{a_j\}_{j \in \mathbb{N}}$ be a conditionally convergent real sequence. Then, for any given $-\infty \leq \alpha \leq \beta \leq \infty$ ($\pm\infty$ is allowed for α and β), there exists a rearrangement $\phi: \mathbb{N} \rightarrow \mathbb{N}$ such that $\liminf_{k \rightarrow \infty} \sum_{j=1}^k a_{\phi(j)} = \alpha$ and $\limsup_{k \rightarrow \infty} \sum_{j=1}^k a_{\phi(j)} = \beta$.

Proof. Let $\{P_j\}_{j \in \mathbb{N}}$ and $\{Q_j\}_{j \in \mathbb{N}}$ be nonnegative terms and absolute value of negative terms of $\{a_j\}_{j \in \mathbb{N}}$. Then, since they differ from $\{a_j^+\}$ and $\{a_j^-\}$ by zero terms, they are also divergent by Lemma 3.3.1.

Let $\{\alpha_\ell\}_{\ell \in \mathbb{N}}$ and $\{\beta_\ell\}_{\ell \in \mathbb{N}}$ be real sequences such that $\lim_{\ell \rightarrow \infty} \alpha_\ell = \alpha$ and $\lim_{\ell \rightarrow \infty} \beta_\ell = \beta$. Let $k_1, m_1 \in \mathbb{N}$ be the smallest integers such that

- $S_1 \triangleq P_1 + \cdots + P_{k_1} > \beta_1$ and
- $T_1 \triangleq S_1 - (Q_1 + \cdots + Q_{m_1}) < \alpha_1$.

Inductively, define $\{k_\ell\}_{\ell \in \mathbb{N}}$ and $\{m_\ell\}_{\ell \in \mathbb{N}}$ by

- $k_{\ell+1} \triangleq \min \{ k \in \mathbb{N}_{>k_\ell} \mid T_\ell + \sum_{j=k_\ell+1}^k P_j > \beta_{\ell+1} \}$
- $S_{\ell+1} \triangleq T_\ell + \sum_{j=k_\ell+1}^{k_{\ell+1}} P_j$
- $m_{\ell+1} \triangleq \min \{ m \in \mathbb{N}_{>m_\ell} \mid S_{\ell+1} - \sum_{j=m_\ell+1}^m Q_j < \alpha_{\ell+1} \}$
- $T_{\ell+1} \triangleq S_{\ell+1} - \sum_{j=m_\ell+1}^{m_{\ell+1}} Q_j$

for each $\ell \in \mathbb{N}$. As $k_\ell \rightarrow \infty$ and $m_\ell \rightarrow \infty$ as $\ell \rightarrow \infty$, this construction gives the natural rearrangement $\phi: \mathbb{N} \rightarrow \mathbb{N}$.

By the construction, we have $|S_\ell - \beta_\ell| \leq P_{k_\ell}$ and $|T_\ell - \alpha_\ell| \leq Q_{m_\ell}$ for each $\ell \in \mathbb{N}$. As $P_j, Q_j \rightarrow 0$ as $j \rightarrow \infty$, we have $S_\ell \rightarrow \beta$ and $T_\ell \rightarrow \alpha$ as $\ell \rightarrow \infty$; α and β are cluster points of $\{\sum_{j=1}^k a_{\phi(j)}\}_{k \in \mathbb{N}}$ (as long as they are finite).

Moreover, for every sufficiently large $n \in \mathbb{N}$, we have $k_\ell + m_\ell \leq n < k_{\ell+1} + m_{\ell+1}$ for some $\ell \in \mathbb{N}$, and thus $\min\{T_\ell, T_{\ell+1}\} \leq \sum_{j=1}^n a_{\phi(j)} \leq S_{\ell+1}$. This, or some more rigorous explanation using arbitrary $\varepsilon \in \mathbb{R}_+$, implies that there do not exist cluster points smaller than α or greater than β . \square

3.4 The Cauchy Product

Definition 3.4.1: Cauchy Product

Given two real sequences $\{a_j\}_{j=0}^\infty$ and $\{b_j\}_{j=0}^\infty$, define

$$C_k \triangleq \sum_{j=0}^k a_j b_{k-j}.$$

The series $\sum_{k=1}^\infty C_k$ is called the *Cauchy product* of $\sum_{j=0}^\infty a_j$ and $\sum_{j=0}^\infty b_j$.

Theorem 3.4.1

Let $\{a_j\}_{j=0}^{\infty}$ and $\{b_j\}_{j=0}^{\infty}$ be two real sequences. Let $\sum_{k=0}^{\infty} C_k$ be the Cauchy product of them.

- (i) If $\sum a_j$ converges absolutely, and if $\sum b_j$ converges, then $\sum C_k$ converges to $(\sum a_j)(\sum b_j)$.
- (ii) If both $\sum a_j$ and $\sum b_j$ converge absolutely, $\sum C_k$ converges absolutely as well.

Proof. (ii) directly follows from the inequality $\sum_{k=0}^n |C_k| \leq (\sum_{j=0}^n |a_j|)(\sum_{j=0}^n |b_j|)$ as long as (i) is proven.

Let $S_n \triangleq \sum_{k=0}^n C_k$, $A_n \triangleq \sum_{j=0}^n a_j$, and $B_n \triangleq \sum_{j=0}^n b_j$. Let $B \triangleq \lim_{n \rightarrow \infty} B_n$ and $\mu_n \triangleq B_n - B$. Then,

$$\begin{aligned} S_n &= \sum_{k=0}^n C_k = \sum_{k=0}^n \sum_{j=0}^k b_{k-j} = \sum_{j=0}^n a_j \sum_{k=j}^n b_{k-j} \\ &= \sum_{j=0}^n a_j B_{n-j} = \sum_{j=0}^n a_j (B + \mu_{n-j}) = B \sum_{j=0}^n a_j + \sum_{j=0}^n a_j \mu_{n-j}. \end{aligned}$$

Claim. $\lim_{n \rightarrow \infty} \sum_{j=0}^n a_j \mu_{n-j} = 0$.

Take any $\varepsilon \in \mathbb{R}_+$ so there exists $N \in \mathbb{N}$ such that

- $|\mu_n| < \varepsilon$ for all $n \geq N$ (by $\mu_n \rightarrow 0$) and
- $\sum_{j=n+1}^m |a_j| < \varepsilon$ for all $m > n \geq N$ (by $\sum_{j=0}^k |a_j|$ being Cauchy).

As μ_n converges, there exists $\mu^* \triangleq \sup_{n \in \mathbb{N}} |\mu_n|$. Let $K_n \triangleq \sum_{j=0}^n a_j \mu_{n-j}$. Whenever $n > 2N$,

$$\begin{aligned} |K_n| &\leq \sum_{j=0}^n |a_j| \cdot |\mu_{n-j}| = \sum_{j=0}^{N-1} |a_j| \cdot |\mu_{n-j}| + \sum_{j=N}^n |a_j| \cdot |\mu_{n-j}| \\ &\leq \varepsilon \sum_{j=0}^{N-1} |a_j| + \mu^* \sum_{j=N}^n |a_j| \leq \varepsilon \left[\sum_{j=0}^n |a_j| + \mu^* \right]. \end{aligned}$$

Hence, $\lim_{n \rightarrow \infty} K_n = 0$; thus $\lim_{n \rightarrow \infty} S_n = (\sum a_j)(\sum b_j)$. □

3.5 Series on Infinite Dimensional Banach Spaces

Definition 3.5.1: Uniform Convergence of Series

Fix a domain $\Omega \subseteq \mathbb{R}^n$. Given a sequence $\{f_j : \Omega \rightarrow \mathbb{R}\}_{j \in \mathbb{N}}$, define $F_n : \Omega \rightarrow \mathbb{R}$ by

$$F_n(x) := \sum_{j=1}^n f_j(x)$$

for each $x \in \Omega$ and $n \in \mathbb{N}$.

- (i) If $\lim_{n \rightarrow \infty} F_n(x)$ exists for all $x \in \Omega$, then the series $\sum_{j=1}^{\infty} f_j$ is said to *converge pointwise on Ω* .
- (ii) Suppose $\sum_{j=1}^{\infty} f_j(x)$ converges pointwise on Ω and let $F(x) \triangleq \lim_{n \rightarrow \infty} F_n(x)$. The series $\sum_{j=1}^{\infty} f_j$ is said to *converge uniformly on Ω* if $\{F_n\}_{n=1}^{\infty}$ uniformly converges to F on Ω .

Theorem 3.5.1

If $\{f_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}^\Omega$ is a sequence of continuous functions and converges uniformly, then $\lim_{n \rightarrow \infty} f_n$ is continuous as well.

Proof. MAS241. □

Definition 3.5.2: Uniform Cauchy

A sequence of function $\{f_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}^\Omega$ is said to be *uniformly Cauchy* on Ω if

$$\forall \varepsilon \in \mathbb{R}_+, \exists N_* \in \mathbb{N}, \forall n, m \geq N_*, \forall x \in \Omega, |f_n(x) - f_m(x)| < \varepsilon.$$

Lemma 3.5.1

A sequence of function $\{f_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}^\Omega$ uniformly converges on Ω if and only if $\{f_n\}_{n \in \mathbb{N}}$ is uniformly Cauchy on Ω .

Proof. (\Rightarrow) Let $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. Take any $\varepsilon \in \mathbb{R}_+$. Then, there exists $N_* \in \mathbb{N}$ such that, if $n \geq N_*$, then $|f_n(x) - f(x)| < \varepsilon/2$ for all $x \in \Omega$. Consequently, whenever $n, m \geq N_*$, $|f_n(x) - f_m(x)| \leq |f_n(x) - f(x)| + |f(x) - f_m(x)| < \varepsilon$.

(\Leftarrow) For each $x \in \mathbb{R}$, the sequence $\{f_n(x)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ is Cauchy. As $(\mathbb{R}, |\cdot|)$ is a Banach space, there uniquely exists the limit $f \triangleq \lim_{n \rightarrow \infty} f_n$. Take any $\varepsilon \in \mathbb{R}_+$. Then, there exists $N_* \in \mathbb{N}$ such that $|f_m(x) - f_n(x)| < \varepsilon/2$ for all $n, m \geq N_*$ and $x \in \Omega$. From this, we get $f_n(x) - \varepsilon/2 \leq \lim_{m \rightarrow \infty} f_m(x) = f(x) \leq f_n(x) + \varepsilon/2$. Hence, $|f_n(x) - f(x)| \leq \varepsilon/2 < \varepsilon$ holds for all $n \geq N_*$ and $x \in \Omega$. □

Note:-

Lemma 3.5.1 holds for arbitrary sequence of functions from Ω to any Banach space.

Lemma 3.5.2

Let $\{f_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^\Omega$ be a series of continuous functions. If $\sum_{j=1}^{\infty} f_j$ converges uniformly on Ω , then $\sum_{j=1}^{\infty} f_j$ is continuous on Ω .

Proof. Lemma 3.5.1. □

Chapter 4

Analysis for Series Functions

4.1 Calculus of Series Functions

Theorem 4.1.1

Given a sequence $\{f_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ of differentiable functions, suppose the following.

- (i) $\{f_j(x_0)\}_{j \in \mathbb{N}} \subseteq \mathbb{R}$ converges for some $x_0 \in (a, b)$.
 - (ii) $\{f'_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ uniformly converges on (a, b) .
- Then, $f_j \rightrightarrows f$ for some $f : (a, b) \rightarrow \mathbb{R}$ on (a, b) . Furthermore, f is differentiable on (a, b) and $\forall x \in (a, b)$, $f'(x) = \lim_{j \rightarrow \infty} f'_j(x)$.

Proof. We shall first show the uniform convergence of $\{f_j\}$. Take any $\varepsilon \in \mathbb{R}_+$. Then, there exists $N \in \mathbb{N}$ such that, for all $j, k \geq N$,

$$(|f_j(x_0) - f_k(x_0)| < \varepsilon/2) \wedge (\forall x \in (a, b), |f'_j(x) - f'_k(x)| < \varepsilon/2(b-a)).$$

By MVT, for all $x, \tilde{x} \in (a, b)$ with $x \neq \tilde{x}$, there exists $x_* \in (a, b)$ such that

$$(f_j - f_k)(x) - (f_j - f_k)(\tilde{x}) = (f_j - f_k)'(x_*) \cdot (x - \tilde{x})$$

Hence, $|(f_j - f_k)(x) - (f_j - f_k)(\tilde{x})| < \varepsilon/2$. Therefore, $|(f_j - f_k)(x)| < \varepsilon$ by triangle inequality obtained by setting $\tilde{x} = x_0$. This directly implies that $\{f_j\}$ is uniformly Cauchy and thus uniformly converges by Lemma 3.5.1. ✓

Let $f_j \rightarrow f$. Fixing $x \in (a, b)$, define

$$\psi_j(t) \triangleq \frac{f_j(t) - f_j(x)}{t - x} \quad \text{and} \quad \psi(t) \triangleq \frac{f(t) - f(x)}{t - x}$$

for $t \in (a, b)$ and $t \neq x$. Now, we claim that $\{\psi_j\}_{j \in \mathbb{N}}$ is uniformly Cauchy. Take any $\varepsilon \in \mathbb{R}_+$. Then, for $j, k \geq N$,

$$|\psi_j(t) - \psi_k(t)| = \left| \frac{(f_j - f_k)(t) - (f_j - f_k)(x)}{t - x} \right| < \frac{\varepsilon}{2(b-a)}.$$

Hence, $\{\psi_j\}$ uniformly converges by Lemma 3.5.1, and $\psi_j \rightarrow \psi$ as $f_j \rightarrow f$.

Let $A_j \triangleq \lim_{t \rightarrow x} \psi_j(t) = f'_j(x)$. By the supposition (ii), we have convergence of $\{A_j\}_{j \in \mathbb{N}}$. Now, we claim that $\lim_{t \rightarrow x} \psi(t) = \lim_{j \rightarrow \infty} A_j$. Let $A_j \rightarrow A$. Take any $\varepsilon \in \mathbb{R}_+$. There exists $N' \in \mathbb{N}$ such that, if $j \geq N'$, we have $|\psi(t) - \psi_j(t)| < \varepsilon/3$ for all $t \in (a, b) \setminus \{x\}$ and $|A_j - A| < \varepsilon/3$. In

addition, from the definition of A_j , there exists $\delta \in \mathbb{R}_+$ such that, whenever $0 < |t - x| < \delta$, we have $|\psi_{N'}(t) - A_{N'}| < \varepsilon/3$. Now, we have

$$|\psi(t) - A| \leq |\psi(t) - \psi_{N'}(t)| + |\psi_{N'}(t) - A_{N'}| + |A_{N'} - A| < \varepsilon$$

for $0 < |t - x| < \delta$. Hence, $f'(x) = \lim_{t \rightarrow x} \psi(t) = \lim_{j \rightarrow \infty} f'_j(x)$. \square

Corollary 4.1.1 Term-by-Term Differentiation

Given a sequence $\{f_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ of differentiable functions, let $F_n = \sum_{j=1}^n f_j$. Suppose the following.

- (i) $\{F_n(x_0)\}_{n \in \mathbb{N}} \subseteq \mathbb{R}$ converges for some $x_0 \in (a, b)$.
 - (ii) $\{F'_n\}_{n \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ uniformly converges on (a, b) .
- Then, $\{F_n\}$ converges uniformly to a function $F: (a, b) \rightarrow \mathbb{R}$ on (a, b) . Furthermore, F is differentiable on (a, b) and $\forall x \in (a, b)$, $F'(x) = \sum_{j=1}^{\infty} f'_j(x)$.

Example 4.1.1

Let $f_j(x) = \sin(x/j^2)$ for $-1 < x < 1$ and $F_n = \sum_{j=1}^n f_j$.

For $x_0 = 0$, the sequence $\{F_n(x_0)\}_{n \in \mathbb{N}}$ converges (to zero). Now, we have $F'_n(x) = \sum_{j=1}^n \cos(x/j^2)/j^2$. Then, for $n, m \in \mathbb{N}$ with $m \geq n$, $|F'_m(x) - F'_n(x)| \leq \sum_{j=n+1}^m 1/j^2 \rightarrow 0$ as $n, m \rightarrow \infty$. Hence, $\{F'_n\}$ is uniformly Cauchy; and thus it converges uniformly by Lemma 3.5.1. Hence, Corollary 4.1.1 guarantees the uniform convergence and differentiability of $\sum_{j=1}^{\infty} f_j$.

Theorem 4.1.2

Given a sequence $\{f_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ of functions Riemann integrable on (a, b) , if $f_j \rightrightarrows f$ on (a, b) , then f is Riemann integrable on (a, b) . Furthermore, $\int_a^b f(x) dx = \lim_{j \rightarrow \infty} \int_a^b f_j(x) dx$.

Proof. \square

Corollary 4.1.2 Term-by-Term Integration

Given a sequence $\{f_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}^{(a,b)}$ of functions Riemann integrable on (a, b) , suppose $\sum f_j \rightrightarrows F$ for some $F: (a, b) \rightarrow \mathbb{R}$. Then, $\int_a^b F(x) dx = \lim_{n \rightarrow \infty} \int_a^b \sum_{j=1}^n f_j(x) dx$.

End.