


Indian Institute of Technology Kharagpur
Department of Mathematics
MA11004 - Linear Algebra, Numeircal and Complex Analysis
Problem Sheet - 7
Spring 2021

1. (a) Evaluate $\int_{1-i}^{2+i} (2x + iy + 1)dz$, along the paths
 - (i) $x = t + 1, y = 2t^2 - 1$,
 - (ii) the straight line joining $1 - i$ and $2 + i$.
- (b) Evaluate $\int_0^{1+i} (x - y + ix^2)dz$ along
 - (i) the straight line from $z = 0$ and $z = 1 + i$,
 - (ii) real axis from $z = 0$ and $z = 1$ and then a line parallel to imaginary axis from $z = 1$ to $z = 1 + i$.
- (c) Find the value of the integral $\int_C (z + 1)^2 dz$ where C is the boundary of the rectangle with vertices at the points $1 + i, -1 + i, -1 - i$ and $1 - i$.
- (d) Compute $\int_{\Gamma} |z| dz$ where Γ is the left half of the unit circle $|z| = 1$ from $z = -i$ to $z = i$.
- (e) Find the value of $\int_C (z^2 - iz) dz$ along the curve C which is $y = x^3 - 3x^2 + 4x - 1$ joining points $(1, 1)$ and $(2, 3)$.
2. (a) Verify that the value of the integral $\int_C z^2 dz$ is same for the following cases:
 - (i) C is the straight line joining the points $A(0, 0)$ and $B(1, 2)$.
 - (ii) C is the straight line path from $A(0, 0)$ to $P(1, 0)$ followed by the straight line path from $P(1, 0)$ to $B(1, 2)$.
 - (iii) C is the parabolic path $y = 2x^2$ joining the points $A(0, 0)$ and $B(1, 2)$.
- (b) Integrate the function $f(z) = xz$ along the straight line from $A(1, 1)$ to $B(2, 4)$ in the complex plane. Is the value same if the path of integration from A to B is along the curve $x = t, y = t^2$?
- (c) Evaluate the function f defined by the integral $f(z) = \oint_{|w|=1} \frac{e^{w^2} - 1}{w - z} dw$.
3. $F(a) = \oint_C \frac{(4z^2 + z + 5)}{(z - a)} dz$, where C is the ellipse $(\frac{x}{2})^2 + (\frac{y}{3})^2 = 1$ taken in counter clockwise sense. Find $F(3.5), F(i), F'(-1)$ and $F''(-i)$.
4. (a) Evaluate $\oint_C \frac{3z^2 + z + 1}{(z^2 - 1)(z + 3)} dz$, where C is the circle $|z| = 2$.
- (b) Compute $\oint_C \frac{e^{z^2}}{(z - 2)} dz$ over the contour $C, C: |z - (2 + i)| = 3$.
- (c) Compute $\oint_C \frac{1}{(z^2 + 4)^2} dz$ over the contour $C: |z - i| = \frac{3}{2}$.

- (d) Evaluate $\oint_C \frac{1}{(z-a)^n} dz$, where n is any integer and C is any closed curve containing 'a'.
5. (a) Show that $|\oint_C \frac{e^z}{z^2+1} dz| \leq \frac{4}{3}\pi e^2$ where $C: |z| = 2$.
- (b) Estimate an upper bound for $|\oint_{|z|=3} \frac{\text{Log}(z)}{z-4i} dz|$.
6. (a) Find the value of $\oint_{|z|=1} \frac{4z^2-4z+1}{(z-2)(z^2+4)} dz$.
- (b) Compute $\oint_{|z+1-i|=2} \frac{z+4}{z^2+2z+5} dz$.
- (c) Compute $\oint_{|z|=6} (\frac{e^{2iz}}{z^4} - \frac{z^4}{(z-i)^3}) dz$.
- (d) Evaluate the integral $\oint_{|z|=1} \frac{dz}{2-\bar{z}}$.
7. Evaluate $\oint_C \frac{\cos z}{z(z^2+8)} dz$ over the contour shown below:



8. Evaluate $\oint_C \frac{z^2+2}{z^2-2}$ where C is the circle
- (a) $|z| = \frac{3}{2}$.
- (b) $|z-1| = 1$.
- (c) $|z| = \frac{1}{2}$.
9. Verify the Cauchy's theorem for the function $z^3 - iz^2 - 5z + 2i$ if C is
- (a) the circle $|z| = 1$.
- (b) the circle $|z-1| = 2$.
- (c) the ellipse $|z-3i| + |z+3i| = 20$.
10. If $0 < r < R$, evaluate the integral $I = \oint_C \frac{R+z}{z(R-z)} dz$, where $C: |z| = r$.
Further using this result deduce that
- (a) $\int_0^{2\pi} \frac{d\theta}{R^2 - 2rR \cos \theta + r^2} = \frac{2\pi}{R^2 - r^2}$.
- (b) $\int_0^{2\pi} \frac{\sin \theta d\theta}{R^2 - 2rR \cos \theta + r^2} = 0$.
11. By integrating $f(z) = \frac{1}{(R-z)}$ over $C: |z| = r$, $0 < r < R$ and using Problem 10, show that
- $$\int_0^{2\pi} \frac{R \cos \theta}{R^2 - 2rR \cos \theta + R^2} d\theta = \frac{2\pi r}{R^2 - r^2}.$$