

# ADVANCED CALCULUS

## MA11003

SECTION 11, 12, & 15CD

Dr. Jitendra Kumar

Professor  
Department of Mathematics  
Indian Institute of Technology Kharagpur  
West Bengal 721302, India



Webpage: <http://www.facweb.iitkgp.ac.in/~jkumar/>

# Concepts Covered

## Differential Calculus

### Functions of Several Variables

- ❑ Method of Lagrange's Multiplier

## Method of Lagrange's Multiplier

Find the Maxima/Minima of the function

$$u = f(x, y) \quad \text{with the constraint} \quad \phi(x, y) = 0$$

Using chain rule

$$\frac{du}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx}$$

$$\text{At the point of extrema} \quad \frac{du}{dx} = 0 \quad \Rightarrow \quad \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx} = 0$$

At the point of extrema  $\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx} = 0$

The equation  $\phi(x, y) = 0$  is satisfied at any point and so at the point of extrema

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0$$

$$\phi(x, y) = 0$$

We eliminate  $\frac{dy}{dx}$  from the above equations.

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0$$

$$\left( \frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} \right) \underbrace{\left( -\frac{f_y}{\phi_y} \right)}_{=:\lambda} + \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx} = 0 \quad \Rightarrow \quad \frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0$$

$$\frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0$$

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0$$

$$\phi(x, y) = 0$$

### Method of Lagrange's Multiplier (Working Rule)

max/min  $u = f(x, y)$  with the constraint  $\phi(x, y) = 0$

Define an auxiliary function  $F(x, y, \lambda) = f(x, y) + \lambda \phi(x, y)$

Necessary conditions for extrema of  $F$

$$F_x = 0 \Rightarrow f_x + \lambda \phi_x = 0$$

$$F_y = 0 \Rightarrow f_y + \lambda \phi_y = 0$$

$$F_\lambda = 0 \Rightarrow \phi = 0$$

## REMARK:

Using the method of Lagrange's multiplier, we obtain stationary points (candidates for extrema). We do not determine the nature of the stationary points. In practice we usually are interested in finding max/min value of the function under some given constraint.

Usually there are a few candidates (critical points), so we can evaluate  $f$  at all of them and choose the largest and smallest values.

Hence, no further test is required if we wish to find only absolute maximum and minimum.

**Problem - 1:** Find maximum/minimum of the function  $x^2 - y^2 - 2x$  in the region  $x^2 + y^2 \leq 1$

**I. Local extrema in the interior**  $x^2 + y^2 < 1$

$$\text{Let } f(x, y) = x^2 - y^2 - 2x$$

$$f_x = 0 \implies x = 1$$

Critical Point: (1,0)

$$f_y = 0 \implies y = 0$$

However this point lies on the boundary so **no critical point lies** in the interior.

## II. Local extrema on the boundary $x^2 + y^2 = 1$

**Problem**            max/ min    $x^2 - y^2 - 2x$     subje to    $x^2 + y^2 = 1$

Auxiliary function for the Lagrange multiplier

$$F(x, y, \lambda) = (x^2 - y^2 - 2x) + \lambda(x^2 + y^2 - 1)$$

**Critical Point:**

$$F_x = 0 \implies x(1 + \lambda) = 1$$


$$F_y = 0 \implies y(\lambda - 1) = 0$$

$$F_\lambda = 0 \implies x^2 + y^2 = 1$$



## II. Local extrema on the boundary $x^2 + y^2 = 1$

$$x(1 + \lambda) = 1$$



$$\lambda = 0, -2$$



$$x = \frac{1}{2}$$

Candidates for extrema

$$y(\lambda - 1) = 0$$



$$y = 0 \text{ \& } \lambda = 1$$



$$y = 0$$



$$\lambda = 1$$

$$(\pm 1, 0)$$

$$\left(\frac{1}{2}, \pm \frac{\sqrt{3}}{2}\right)$$

$$x^2 + y^2 = 1$$



$$x = \pm 1$$



$$y = \pm \frac{\sqrt{3}}{2}$$

### III. Function Values:

Candidates for extrema

$$(\pm 1, 0) \quad \left(\frac{1}{2}, \pm \frac{\sqrt{3}}{2}\right)$$

$$f(x, y) = x^2 - y^2 - 2x$$

Points	$(1, 0)$	$(-1, 0)$	$\left(\frac{1}{2}, \pm \frac{\sqrt{3}}{2}\right)$
Function Value	$-1$	$3$	$-\frac{3}{2}$

Maximum value of the function:  $3$

Minimum value of the function:  $-\frac{3}{2}$



**QUIZ QUESTION ?**

LINK FOR RESPONSES: <http://www.facweb.iitkgp.ac.in/~jkumar/teach/MA11003.html>

Consider the problem

Maximize  $f(x, y) = x^2 + y^2$  in the domain

$$(x - 2)^2 + (y - 1)^2 \leq 20$$

Critical points are

<b>(A)</b>	$(0, 0), (2, 1), (6, 3)$
<b>(B)</b>	$(0, 0), (2, -1), (6, 3)$
<b>(C)</b>	$(0, 0), (-2, 1), (6, 3)$
<b>(D)</b>	$(0, 0), (-2, -1), (6, 3)$

**Problem - 2:** Find maximum and minimum values of the function  $f(x, y) = x^2 + y^2$   
in the region  $(x - 2)^2 + (y - 1)^2 \leq 20$

**I. Local extrema in the interior**  $(x - 2)^2 + (y - 1)^2 < 20$

$$f_x = 0 \implies x = 0$$

$$f_y = 0 \implies y = 0$$

Critical Point: (0,0)

**II. Local extrema on the boundary**  $(x - 2)^2 + (y - 1)^2 = 20$

Problem    max/min     $x^2 + y^2$     subject to     $(x - 2)^2 + (y - 1)^2 = 20$

Auxiliary function for the Lagrange multiplier

$$F(x, y, \lambda) = (x^2 + y^2) + \lambda((x - 2)^2 + (y - 1)^2 - 20)$$

$$F(x, y, \lambda) = (x^2 + y^2) + \lambda((x - 2)^2 + (y - 1)^2 - 20)$$

Critical Point:

$$F_x = 0 \Rightarrow x + \lambda(x - 2) = 0 \Rightarrow (x - 2) = -\frac{2}{1 + \lambda} \Rightarrow x = -2, 6$$

$$F_y = 0 \Rightarrow y + \lambda(y - 1) = 0 \Rightarrow (y - 1) = -\frac{1}{1 + \lambda} \Rightarrow y = -1, 3$$

$$F_\lambda = 0 \Rightarrow (x - 2)^2 + (y - 1)^2 = 20 \Rightarrow (1 + \lambda) = \pm \frac{1}{2} \Rightarrow \lambda = -\frac{1}{2}, -\frac{3}{2}$$

Points	$(0, 0)$	$(-2, -1)$	$(6, 3)$
Function Value	0	5	45

Minimum value : 0

Maximum value: 45

# KEY TAKEAWAY

## Method of Lagrange's Multiplier

max/min  $u = f(x, y)$  with the constraint  $\phi(x, y) = 0$

Auxiliary function  $F(x, y, \lambda) = f(x, y) + \lambda \phi(x, y)$

Necessary conditions for extrema of  $F$

$$F_x = 0 \Rightarrow f_x + \lambda \phi_x = 0$$

$$F_y = 0 \Rightarrow f_y + \lambda \phi_y = 0$$

$$F_\lambda = 0 \Rightarrow \phi = 0$$

*Thank You*