







X

You may be eligible for this trial

Get started

# You are not out of options

Learn more about a clinical trial for men with metastatic castrate-resistant

prostate cancer

Get started

The MEVPRO-1 clinical tri for men who have metasta responding to hormone the Zytiga®).

#### We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

Decline All

Accept All

**Cookies Preferences** 

The purpose of MEVPRO-1 is to see if taking the study medicine (mevrometostat) alongside a standard treatment (enzalutamide) is safe and works to slow down or stop

You may be eligible for this trial

Get started

# Watch a short video about the study medicine



#### We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

# Abo

The study medicine is called mev day.

Prostate cancer cells need hormones (such as testosterone) to grow and divide and can often become resistant to hormone therapy and continue to progress and spread to other parts of the body.

You may be eligible for this trial

Get started

cancer to progress.

Mevrometostat, the study medicine, is thought to work by blocking the abnormal EZH2 activity in cells, which may help prevent or delay hormone therapy resistance and cancer growth and progression.

# Who may participate

The MEVPRO-1 clinical trial may be an option for adults with metastatic castrate-resistant prostate cancer that has become resistant to hormone therapy. Specifically, participants must have previously been treated with

abiraterone acetate (Zytiga®) for a

# We Care About Your Privacy

Participants can have received ch chemotherapy since their cancer Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

#### Condition

Metastatic Prostate C

Age

18+ years

#### You may be eligible for this trial

Get started

Male

Each clinical study has its own guidelines for who can participate, called eligibility criteria. However, only the research study staff can determine if you qualify to enroll in the study.

#### View more eligibility criteria >



# Get started – See if you may be eligible

# Answer a 2-minute questionna

A first step as you consider this about your health and medical good fit for you, you may choose Pfizer Clinical Trial Contact Ce

# Speak with a study represental

If you're interested, a Pfizer Cli do a more detailed review of th be able to participate and be re

#### We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

# Confirm your eligibility.

Your answers to these questions will only be linked to you if your responses indicate that you may be eligible to participate in this study and you choose to share your contact information with the study team. Pfizer study team members

You may be eligible for this trial

Get started

Answer a 2-minute questionnaire

If you would like to proceed with taking the questionnaire, you may click 'next'.

Next
------

Continue →

# What to expect



Participants will be ass

Study medicine (mevro

#### OR

• Enzalutamide *OR* docetaxel (de

#### We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

Mevrometostat is taken as tablets by mouth twice-a-day and enzalutamide is taken as capsules by mouth once-a-day.

You may be eligible for this trial

Get started

#### Length of study treatment

(i)

Depends on how the cancer responds and how well the study medicine is tolerated

#### Number of study visits



3 or 4 in the first 2 months, then every 3 or 4 weeks

# Long-term follow up

Every 12 weeks

#### We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

# Frequently asked questions

#### How long do study visits last?

You may be eligible for this trial

Get started

cancer worsens, they have any side effects that become too severe, they or their doctor decide they should stop, or the trial ends. Participants receiving docetaxel may be treated for up to 30 weeks.

Is this helpful?

#### What are the possible benefits of participating?



# What are the possible risks of par

## We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>



Discover lo



# You may be eligible for this trial

Get started



How clinical trials work



Protecting your safety & privacy



Steps to joi clinical trial





Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>

>



**About Pfizer** 

For investigators

The Breakthrough newsl

Site map

Terms of use

#### **Privacy policy**

#### Contact us

# You may be eligible for this trial

Get started







# We Care About Your Privacy

Pfizer uses cookies and similar technologies to enhance and personalize your customer experience. By granting us permission, you enable Pfizer and our analytics and marketing partners to collect, use, and share information about your website interactions to tailor your digital experiences, our services, and content for you. To learn more about how Pfizer uses these technologies, please read our <a href="Privacy Policy">Privacy Policy</a>