Checklist of the Birds of Kenya

>FOURTH EDITION



Bird Committee, Nature Kenya
East Africa Natural History Society

Nairobi, September 2009 + corrections

NatureKenya
The East Africa Natural History Society

6 166000 025765 >

Checklist of the Birds of Kenya

Fourth Edition

Key to the abbreviations used

AM Afrotropical migrant[1]
E endemic species or race
Ex species which are thought to have become extinct in Kenya
IO visitor from Indian Ocean islands
MM migrant[1] from the Malagasy region
OM migrant[1] from the Oriental region
PM migrant[1] from the Palaearctic region
RS visitor from the Red Sea
SO visitor from the southern oceans or the Antarctic
VIO vagrant from the northwest Indian Ocean
VN vagrant from the Nearctic region
VP vagrant from the Palaearctic region
VS vagrant from the southern oceans or the Antarctic
VSA vagrant from southern Africa

When these letters are in lower case, migrants of that category may occur alongside resident, non-migratory individuals or individuals of one of the other migrant categories. When they are in round brackets, some birds in Kenya might be migrants or wanderers from that particular region or area.

If possible, where there are fewer than four records, these are listed under the species name. Where no reference is given, the record is usually listed in Zimmerman *et al.* (1996) or has been accepted by the East Africa Rarities Committee (EARC).

The BirdLife/IUCN criteria for assigning threat status and category (Critically Endangered, Endangered, Vulnerable, Near Threatened, and Data Deficient) are included where appropriate.

[1] 'Migrant', in the accepted ornithological sense, means a bird that moves on a regular, seasonal basis from one, to one or more defined areas; thus, the majority of wader species that breed in the High Arctic leave their breeding areas to spend most of the year on coasts and wetlands throughout the temperate and tropical areas of the world; they are therefore true migrants. On the other hand, Little Grebes that travel, often at night, in search of newly formed ponds, are wanderers, not migrants.

ISBN: 616-6-00-002576-5
© Bird Committee, Nature Kenya—the East Africa Natural History Society, September 2009
The Nature Kenya office is at the National Museums of Kenya, Museum Hill, Nairobi
Postal address: P. O. Box 44486, 00100 Nairobi, Kenya
tel: (+254) (0) 20 374 6090 or 374 9957; fax: (+254) (0) 20 374 1049
e-mail: office@naturekenya.org; website: www.naturekenya.org

Publishing history:
Check-list of the Birds of Kenya, Ornithological Sub-Committee [OS-C], EANHS, March 1981
Check-list of the Birds of Kenya, Second Edition, OS-C, EANHS, May 1986
Check-list of the Birds of Kenya, Third Edition, OS-c, EANHS, June 1996

Printed in Kenya by Colourprint Limited
Industrial Area, Road-C, off Enterprise Road (third right turn), behind Sameer Africa
Postal address: P. O. Box 44466, 00100 Nairobi, Kenya
tel: (+254) (0) 20 556 278/556 611/533 895/531 310; fax: (+254) (0) 20 552 992/554 156
cell: (+254) (0) 722 203 645 *or* (+254) (0) 733 203 645
e-mail: info@colourprint.co.ke

Cover photograph: Sharpe's Longclaw *Macronynx sharpei* by Peter Usher
taken on 30 August 2009 on the North Kinangop
using a Canon 450D camera with a Canon EF100 – 400 mm IS lens;
exposure details: 1/1600 s at f5.6, 400 mm, ISO setting 400



Map of Kenya showing Important Bird Areas (IBAs), National Parks and National Reserves

BirdLife INTERNATIONAL

BirdLife IBA IMPORTANT BIRD AREA

KENYA WILDLIFE SERVICE

RSPB

Checklist

With minor corrections, 13 December 2017

Struthionidae: Ostrich

☐ Ostrich *Struthio camelus*
In the 1996 list, *Struthio (camelus) molybdophanes*, English name the Somali Ostrich, was treated as a separate species (Freitag & Robinson 1993), necessitating the qualifier 'Common' for the species containing the other two races (nominate *camelus* and *massaicus*); with the lumping in the present list no qualifier is required

Numididae: guineafowl

☐ Helmeted Guineafowl *Numida meleagris*
☐ Crested Guineafowl *Guttera pucherani*
The nominate, mainly coastal and lowland race is sometimes regarded as being a species, the Kenya Crested Guineafowl; mainly western and highland forest birds of the race *verreauxi* are then split off as a species, the Crested Guineafowl
☐ Vulturine Guineafowl *Acryllium vulturinum*

Phasianidae: quails, francolins, spurfowl and allies

☐ Stone Partridge *Ptilopachus petrosus*
☐ Coqui Francolin *Francolinus coqui*
☐ Ring-necked Francolin *Francolinus streptophorus* **Near Threatened**
☐ Red-winged Francolin *Francolinus levaillantii*
☐ Shelley's Francolin *Francolinus shelleyi*
Francolinus shelleyi macarthuri **E**
☐ Moorland Francolin *Francolinus psilolaemus*
☐ Orange River Francolin *Francolinus levaillantoides*
One collected, October 1973, Mt Kulal and others seen and tape-recorded, November 1989, Huri Hills
☐ Crested Francolin *Francolinus sephaena*
☐ Scaly Francolin *Francolinus squamatus*
☐ Hildebrandt's Francolin *Francolinus hildebrandti*
Francolinus hildebrandti altumi **E**
☐ Jackson's Francolin *Francolinus jacksoni* **E**
☐ Chestnut-naped Francolin *Francolinus castaneicollis*
Sight record, 21 June 1975, south of Moyale (Britton 1980)
☐ Red-necked Spurfowl *Francolinus afer*
☐ Yellow-necked Spurfowl *Francolinus leucoscepus*
☐ Common Quail *Coturnix coturnix* **pm**
☐ Harlequin Quail *Coturnix delegorguei* **am**
☐ Blue Quail *Coturnix adansonii* **AM**

Anatidae: ducks and geese

☐ White-faced Whistling Duck *Dendrocygna viduata*
☐ Fulvous Whistling Duck *Dendrocygna bicolor* **AM**

- ☐ White-backed Duck *Thalassornis leuconotus*
- ☐ Spur-winged Goose *Plectropterus gambensis*
- ☐ Knob-billed Duck *Sarkidiornis melanotos* **am**
- ☐ Egyptian Goose *Alopochen aegyptiaca*
  It was *Alopochen aegyptiacus* in the 1996 list; gender change follows David & Gosselin (2002b)
- ☐ African Pygmy Goose *Nettapus auritus*
- ☐ Cape Teal *Anas capensis*
- ☐ Gadwall *Anas strepera* **PM**
- ☐ Eurasian Wigeon *Anas penelope* **PM**
- ☐ African Black Duck *Anas sparsa*
- ☐ Mallard *Anas platyrhynchos* **VP**
  The second record since 1929 was of nine, 30 November – 3 December 2002, Marsabit (Borghesio *et al*. 2004)
- ☐ Yellow-billed Duck *Anas undulata* **am**
- ☐ Northern Shoveler *Anas clypeata* **PM**
- ☐ Red-billed Teal *Anas erythrorhyncha*
- ☐ Northern Pintail *Anas acuta* **PM**
- ☐ Garganey *Anas querquedula* **PM**
- ☐ Eurasian Teal *Anas crecca* **PM**
  It was called the Common Teal in the 1996 list
- ☐ Hottentot Teal *Anas hottentota*
- ☐ Southern Pochard *Netta erythrophthalma* **am**
- ☐ Common Pochard *Aythya ferina* **PM**
- ☐ Ferruginous Duck *Aythya nyroca* **PM**
- ☐ Tufted Duck *Aythya fuligula* **PM**
- ☐ Maccoa Duck *Oxyura maccoa* **Near Threatened**

Diomedeidae: albatrosses

Both these species were included in *Diomedea* in the 1996 list

- ☐ Black-browed Albatross *Thalassarche melanophrys* **SO**
  The specific epithet was spelled *melanophris* in the 1996 list, following several earlier authors, e.g., White (1965), Britton (1980) and Harrison (1983)
- ☐ Shy Albatross *Thalassarche cauta* **SO**

Procellariidae: petrels and shearwaters

- ☐ Cape Petrel *Daption capense* **VS**
  The sole record, 26 September 1974, off Mombasa. It was called the Pintado Petrel in the 1996 list
- ☐ Antarctic Prion *Pachyptila desolata* **SO**
  Three records, September 1983, north of Malindi; August 1988, Watamu; and September 1990, Shimoni
- ☐ Slender-billed Prion *Pachyptila belcheri* **VS**
  The sole record, one found dead, 16 August 1984, Watamu
- ☐ White-chinned Petrel *Procellaria aequinoctialis* **VS**
  The sole records are of individuals photographed, 9 August and 15 September 1990, off Shimoni
- ☐ Wedge-tailed Shearwater *Puffinus pacificus* **IO**

☐ Audubon's Shearwater *Puffinus lherminieri* **IO**

☐ Sooty Shearwater *Puffinus griseus* **VS**
First Kenya record, one found dead, 30 May 2004, Watamu (Jackson & Valle 2007)

☐ Jouanin's Petrel *Bulweria fallax* **VIO; Near Threatened**
All specimens: 13 December 1953 and 9 December 1985, off Malindi; 3 December 1991, off Mtwapa

Hydrobatidae: storm-petrels

☐ Wilson's Storm-petrel *Oceanites oceanicus* **SO**

☐ Black-bellied Storm-petrel *Fregetta tropica* **VS**
The only records are of two beached specimens, June 1988, Watamu and June 1994, near Malindi

☐ Leach's Storm-petrel *Oceanodroma leucorhoa* **VP**
A specimen washed up, 8 February 1969 (not 1967 as stated in Zimmerman *et al.*), Tiwi River mouth, south of Mombasa (Parsons 1969), and three further sight records (see Zimmerman *et al.*)

☐ Matsudaira's Storm-petrel *Oceanodroma matsudairae* **VP; Data Deficient**
The only records are of two sightings, July and August 1981, off Mombasa

Podicipedidae: grebes

☐ Little Grebe *Tachybaptus ruficollis*

☐ Great Crested Grebe *Podiceps cristatus*

☐ Black-necked Grebe *Podiceps nigricollis*

Phoenicopteridae: flamingos

☐ Greater Flamingo *Phoenicopterus roseus* **am, pm**

☐ Lesser Flamingo *Phoeniconaias minor* **am; Near Threatened**

Ciconiidae: storks

☐ Yellow-billed Stork *Mycteria ibis* **(am)**

☐ African Open-billed Stork *Anastomus lamelligerus* **am**

☐ Black Stork *Ciconia nigra* **PM**

☐ Abdim's Stork *Ciconia abdimii* **AM**

☐ Woolly-necked Stork *Ciconia episcopus*

☐ White Stork *Ciconia ciconia* **PM**

☐ Saddle-billed Stork *Ephippiorhynchus senegalensis*

☐ Marabou Stork *Leptoptilos crumeniferus*

Threskiornithidae: ibises and spoonbills

☐ Sacred Ibis *Threskiornis aethiopicus*

☐ Olive Ibis *Bostrychia olivacea*
It was called the African Green Ibis in the 1996 list

☐ Hadada Ibis *Bostrychia hagedash*

☐ Glossy Ibis *Plegadis falcinellus* **am, pm**

☐ Eurasian Spoonbill *Platalea leucorodia* **PM**

☐ African Spoonbill *Platalea alba*

Ardeidae: herons, egrets and bitterns

- ☐ Little Bittern *Ixobrychus minutus* **am, pm**
- ☐ Dwarf Bittern *Ixobrychus sturmii* **am**
- ☐ White-backed Night Heron *Gorsachius leuconotus*
- ☐ Black-crowned Night Heron *Nycticorax nycticorax* **am, pm**
- ☐ Striated Heron *Butorides striata*
  It was called the Green-backed Heron *B. striatus* in the 1996 list; gender change follows David & Gosselin (2002b)
- ☐ Squacco Heron *Ardeola ralloides* **am, pm**
  It was called the Common Squacco Heron in the 1996 list
- ☐ Madagascar Pond Heron *Ardeola idae* **MM; Endangered**
  It was called the Madagascar Squacco Heron in the 1996 list
- ☐ Rufous-bellied Heron *Ardeola rufiventris*
- ☐ Cattle Egret *Bubulcus ibis* **am**
- ☐ Grey Heron *Ardea cinerea* **am, pm**
- ☐ Black-headed Heron *Ardea melanocephala*
- ☐ Goliath Heron *Ardea goliath*
- ☐ Purple Heron *Ardea purpurea* **(pm)**
- ☐ Great White Egret *Ardea alba*
  It was called *Casmerodius albus* and Great Egret in the 1996 list
- ☐ Yellow-billed Egret *Egretta intermedia*
  It was called *Mesophoyx intermedia* in the 1996 list
- ☐ Black Heron *Egretta ardesiaca*
- ☐ Little Egret *Egretta garzetta*
- ☐ Western Reef Heron *Egretta gularis* **AM**
- ☐ Dimorphic Egret *Egretta dimorpha*
  It was called *Egretta (garzetta) dimorpha* in the 1996 list

Phaethontidae: tropicbirds

- ☐ White-tailed Tropicbird *Phaethon lepturus* **IO**

Fregatidae: frigatebirds

- ☐ Greater Frigatebird *Fregata minor* **IO**
- ☐ Lesser Frigatebird *Fregata ariel* **IO**
  Three confirmed records only: January 1980, December 1994 and March 2003, all Malindi–Watamu area

Scopidae: Hamerkop

- ☐ Hamerkop *Scopus umbretta*

Balaenicipitidae: Shoebill

- ☐ Shoebill *Balaeniceps rex* **Vulnerable**
  Two modern records: September–November 1994, Masai Mara, and December 1994–mid 1995, Amboseli

Pelecanidae: pelicans

- ☐ Great White Pelican *Pelecanus onocrotalus*
- ☐ Pink-backed Pelican *Pelecanus rufescens*

Sulidae: boobies

- ☐ Masked Booby *Sula dactylatra* **IO**
- ☐ Red-footed Booby *Sula sula* **IO**
- ☐ Brown Booby *Sula leucogaster* **IO**

Phalacrocoracidae: cormorants

- ☐ Reed Cormorant *Phalacrocorax africanus*
  It was called the Long-tailed Cormorant in the 1996 list
- ☐ Great Cormorant *Phalacrocorax carbo*

Anhingidae: darters

- ☐ African Darter *Anhinga rufa*

Falconidae: falcons

- ☐ Pygmy Falcon *Polihierax semitorquatus*
- ☐ Lesser Kestrel *Falco naumanni* **PM; Vulnerable**
- ☐ Common Kestrel *Falco tinnunculus* **pm**
  The resident race *rufescens* is often called the Rock Kestrel
- ☐ Greater Kestrel *Falco rupicoloides*
- ☐ Fox Kestrel *Falco alopex*
- ☐ Grey Kestrel *Falco ardosiaceus*
  Specific epithet spelled *ardosiacus* in the 1996 list
- ☐ Dickinson's Kestrel *Falco dickinsoni*
- ☐ Red-necked Falcon *Falco chiquera*
- ☐ Red-footed Falcon *Falco vespertinus* **PM; Near Threatened**
- ☐ Amur Falcon *Falco amurensis* **PM**
- ☐ Eleonora's Falcon *Falco eleonorae* **PM**
- ☐ Sooty Falcon *Falco concolor* **PM; Near Threatened**
- ☐ Eurasian Hobby *Falco subbuteo* **PM**
- ☐ African Hobby *Falco cuvieri*
- ☐ Lanner Falcon *Falco biarmicus*
- ☐ Saker Falcon *Falco cherrug* **PM; Endangered**
- ☐ Peregrine Falcon *Falco peregrinus* **pm**
- ☐ Barbary Falcon *Falco pelegrinoides* **PM**
- ☐ Taita Falcon *Falco fasciinucha* **Near Threatened**

Accipitridae: diurnal birds of prey other than falcons

- ☐ Secretarybird *Sagittarius serpentarius*
  It was included in its own family, Sagittariidae, in the 1996 list and called the Secretary Bird

- ☐ Osprey *Pandion haliaetus* **PM**
- ☐ African Cuckoo Hawk *Aviceda cuculoides* **am**
- ☐ European Honey Buzzard *Pernis apivorus* **PM**
  It was called the Eurasian Honey Buzzard in the 1996 list
- ☐ Bat Hawk *Macheiramphus alcinus*
- ☐ African Black-shouldered Kite *Elanus caeruleus* [**am**]
  It was called simply Black-shouldered Kite in the 1996 list. There is a ringing recovery, from near Pretoria to eastern Uganda, of this sometimes dispersive species (Oatley *et al.* 1998)
- ☐ African Swallow-tailed Kite *Chelictinia riocourii*
- ☐ Black Kite *Milvus migrans* **am, pm**
  The Afrotropical race *parasitus* is often called the Yellow-billed Kite
- ☐ African Fish Eagle *Haliaeetus vocifer*
- ☐ Palm-nut Vulture *Gypohierax angolensis*
- ☐ Lammergeier *Gypaetus barbatus*
  Specific epithet was incorrectly spelled *barbarus* in the 1996 list
- ☐ Egyptian Vulture *Neophron percnopterus* **Endangered**
- ☐ Hooded Vulture *Necrosyrtes monachus*
- ☐ White-backed Vulture *Gyps africanus* **Near Threatened**
  It was called the African White-backed Vulture in the 1996 list
- ☐ Rüppell's Vulture *Gyps rueppellii* **Near Threatened**
  It was called Rüppell's Griffon Vulture in the 1996 list
- ☐ White-headed Vulture *Trigonoceps occipitalis* **Vulnerable**
- ☐ Lappet-faced Vulture *Torgos tracheliotus* **Vulnerable**
  It was called *Torgos tracheliotos* in the 1996 list
- ☐ Short-toed Snake Eagle *Circaetus gallicus* **PM**
  The only confirmed record is of one seen on 2 February 2001 in Tsavo East National Park (EARC 2005), all previous claimed records having been rejected
- ☐ Beaudouin's Snake Eagle *Circaetus beaudouini* **Vulnerable**
  Recently raised to species level; the only confirmed record is of one seen on 21 January 2001 at Mungatsi, western Kenya (EARC 2005)
- ☐ Black-chested Snake Eagle *Circaetus pectoralis*
- ☐ Brown Snake Eagle *Circaetus cinereus*
- ☐ Southern Banded Snake Eagle *Circaetus fasciolatus* **Near Threatened**
- ☐ Western Banded Snake Eagle *Circaetus cinerascens*
  It was called simply the Banded Snake Eagle in the 1996 list
- ☐ Bateleur *Terathopius ecaudatus* **Near Threatened**
- ☐ Western Marsh Harrier *Circus aeruginosus* **PM**
  It was called the Eurasian Marsh Harrier in the 1996 list
- ☐ African Marsh Harrier *Circus ranivorus*
- ☐ Pallid Harrier *Circus macrourus* **PM; Near Threatened**
- ☐ Montagu's Harrier *Circus pygargus* **PM**
- ☐ African Harrier Hawk *Polyboroides typus*
  Often called by the Afrikaans name Gymnogene
- ☐ Dark Chanting Goshawk *Melierax metabates*

- ☐ Eastern Chanting Goshawk *Melierax poliopterus*
  It was called the Eastern Pale Chanting Goshawk in the 1996 list
- ☐ Gabar Goshawk *Micronisus gabar*
- ☐ African Goshawk *Accipiter tachiro*
- ☐ Shikra *Accipiter badius*
- ☐ Levant Sparrowhawk *Accipiter brevipes* **PM**
- ☐ Little Sparrowhawk *Accipiter minullus*
- ☐ Ovambo Sparrowhawk *Accipiter ovampensis*
- ☐ Eurasian Sparrowhawk *Accipiter nisus* **PM**
- ☐ Rufous-breasted Sparrowhawk *Accipiter rufiventris*
- ☐ Great Sparrowhawk *Accipiter melanoleucus*
  Sometimes called the Black Sparrowhawk (as it is in southern Africa)
- ☐ Lizard Buzzard *Kaupifalco monogrammicus*
- ☐ Grasshopper Buzzard *Butastur rufipennis* **AM**
- ☐ Common Buzzard *Buteo buteo* **PM**
  The race that occurs here, *vulpinus*, is known as the Steppe Buzzard
- ☐ Mountain Buzzard *Buteo oreophilus*
- ☐ Long-legged Buzzard *Buteo rufinus* **PM**
- ☐ Augur Buzzard *Buteo augur*
- ☐ Lesser Spotted Eagle *Aquila pomarina* **PM**
- ☐ Greater Spotted Eagle *Aquila clanga* **PM; Vulnerable**
- ☐ Tawny Eagle *Aquila rapax*
- ☐ Steppe Eagle *Aquila nipalensis* **PM**
- ☐ Eastern Imperial Eagle *Aquila heliaca* **PM; Vulnerable**
  In the 1996 list it was called simply the Imperial Eagle; the qualifier is now required as the Spanish form has been given specific rank (*Aquila adalberti*)
- ☐ Verreaux's Eagle *Aquila verreauxii*
- ☐ Wahlberg's Eagle *Aquila wahlbergi* **am**
- ☐ African Hawk Eagle *Aquila spilogaster*
  This and the next two species were included in *Hieraaetus* in the 1996 list
- ☐ Booted Eagle *Aquila pennata* **PM**
- ☐ Ayres's Hawk Eagle *Aquila ayresii*
- ☐ Martial Eagle *Polemaetus bellicosus* **Near Threatened**
- ☐ Long-crested Eagle *Lophaetus occipitalis*
- ☐ Cassin's Hawk Eagle *Spizaetus africanus*
  One juvenile collected, 17 April 1926 on Mt Elgon (see Clark & Edelstam 2001) and an adult photographed, 16 May 2006 in the Imenti Forest, Meru District. It was called Cassin's Hawk-Eagle by Gill & Wright (2006)
- ☐ Crowned Eagle *Stephanoaetus coronatus*
  It was called the African Crowned Eagle in the 1996 list

Otididae: bustards

- ☐ Arabian Bustard *Ardeotis arabs*
  A pair collected, January 1932, northern Turkana, and one photographed, January 1989 near Ileret

- ☐ Kori Bustard *Ardeotis kori*
- ☐ Denham's Bustard *Neotis denhami* **Near Threatened**
- ☐ Heuglin's Bustard *Neotis heuglinii*
- ☐ White-bellied Bustard *Eupodotis senegalensis*
- ☐ Buff-crested Bustard *Lophotis gindiana*
  It was included in *Eupodotis ruficrista* in the 1996 list
- ☐ Black-bellied Bustard *Lissotis melanogaster*
  This and the next species were in *Eupodotis* in the 1996 list
- ☐ Hartlaub's Bustard *Lissotis hartlaubii*

Rallidae: rails and relatives

- ☐ White-spotted Flufftail *Sarothrura pulchra*
- ☐ Buff-spotted Flufftail *Sarothrura elegans*
- ☐ Red-chested Flufftail *Sarothrura rufa*
- ☐ Streaky-breasted Flufftail *Sarothrura boehmi* **AM**
- ☐ Striped Flufftail *Sarothrura affinis*
  One caught and ringed on 19 June 1977 in the Ngurumans (Zimmerman *et al.* 1996) may be the sole confirmed record as those from the 1920s and 1930s are disputed
- ☐ African Water Rail *Rallus caerulescens*
- ☐ African Crake *Crex egregia* **am**
- ☐ Corncrake *Crex crex* **PM; Near Threatened**
- ☐ Black Crake *Amaurornis flavirostra*
  Specific epithet was *flavirostris* in the 1996 list; the change follows David & Gosselin (2002a)
- ☐ Baillon's Crake *Porzana pusilla* **am, (pm)**
- ☐ Spotted Crake *Porzana porzana* **PM**
- ☐ Striped Crake *Aenigmatolimnas marginalis* **AM**
- ☐ Purple Swamphen *Porphyrio porphyrio*
- ☐ Allen's Gallinule *Porphyrio alleni*
- ☐ Common Moorhen *Gallinula chloropus*
- ☐ Lesser Moorhen *Gallinula angulata* **AM**
- ☐ Red-knobbed Coot *Fulica cristata* **am**

Heliornithidae: finfoots

- ☐ African Finfoot *Podica senegalensis*

Gruidae: cranes

- ☐ Grey Crowned Crane *Balearica regulorum* **Vulnerable**
- ☐ Black Crowned Crane *Balearica pavonina* **Near Threatened**
- ☐ Demoiselle Crane *Anthropoides virgo* **PM**
  The sole record is of nine photographed on 15 January 1986, north of Malindi
- ☐ Common Crane *Grus grus* **PM**
  The first Kenya record was of one northeast of Eldoret on 2 October 1999 (EARC 2004); two seen and photographed by many people, January–February 2008, Lake Naivasha, has not yet been published

Turnicidae: buttonquails and Quailplover

☐ Common Buttonquail *Turnix sylvaticus*
It was called the Common Button-quail in the 1996 list

☐ Black-rumped Buttonquail *Turnix hottentottus*
Only three confirmed records since 1955: February 1958, August–November 1988 and October 1993, all in the Kitale area. In the 1996 list it was called *Turnix hottentotta*; the gender is changed now following David & Gosselin (2002b); also, the English name was Black-rumped Button-quail

☐ Quailplover *Ortyxelos meiffrenii*
It was called Quail-plover in the 1996 list; Gill & Wright (2006) also use Quail-plover

Burhinidae: thick-knees

☐ Stone-curlew *Burhinus oedicnemus* **PM**
It was called the Eurasian Thick-knee in the 1996 list

☐ Senegal Thick-knee *Burhinus senegalensis*
☐ Water Thick-knee *Burhinus vermiculatus*
☐ Spotted Thick-knee *Burhinus capensis*

Haematopodidae: oystercatchers

☐ Eurasian Oystercatcher *Haematopus ostralegus* **PM**

Dromadidae: Crab-plover

☐ Crab-plover *Dromas ardeola* **AM**

Recurvirostridae: stilts and avocets

☐ Black-winged Stilt *Himantopus himantopus* **am, (pm)**
☐ Pied Avocet *Recurvirostra avosetta* **am**

Charadriidae: plovers

☐ Northern Lapwing *Vanellus vanellus* **PM**
The only record is of one, February 1995–August 1996, Sabaki estuary, north of Malindi (Jackson 1997)

In the following nine *Vanellus* species, the word 'Lapwing' is sometimes used instead of 'Plover'

☐ Long-toed Plover *Vanellus crassirostris*
☐ Blacksmith Plover *Vanellus armatus*
☐ Spur-winged Plover *Vanellus spinosus*
☐ Black-headed Plover *Vanellus tectus*
☐ Senegal Plover *Vanellus lugubris*
☐ Black-winged Plover *Vanellus melanopterus*
☐ Crowned Plover *Vanellus coronatus*
☐ African Wattled Plover *Vanellus senegallus*
☐ Brown-chested Plover *Vanellus superciliosus*
☐ Pacific Golden Plover *Pluvialis fulva* **PM, (NM)**
☐ Grey Plover *Pluvialis squatarola* **PM**
☐ Common Ringed Plover *Charadrius hiaticula* **PM**
It was called simply the Ringed Plover in the 1996 list

☐ Little Ringed Plover *Charadrius dubius* **PM**

- ☐ Kittlitz's Plover *Charadrius pecuarius* **am**
- ☐ Three-banded Plover *Charadrius tricollaris*
- ☐ White-fronted Plover *Charadrius marginatus*
- ☐ Kentish Plover *Charadrius alexandrinus* **PM**
- ☐ Chestnut-banded Plover *Charadrius pallidus* **Near Threatened**
- ☐ Lesser Sand Plover *Charadrius mongolus* **PM**
  It was called the Lesser Sandplover in the 1996 list
- ☐ Greater Sand Plover *Charadrius leschenaultii* **PM**
  It was called the Greater Sandplover in the 1996 list
- ☐ Caspian Plover *Charadrius asiaticus* **PM**

Rostratulidae: painted-snipes

- ☐ Greater Painted-snipe *Rostratula benghalensis*

Jacanidae: jacanas

- ☐ Lesser Jacana *Microparra capensis*
- ☐ African Jacana *Actophilornis africanus*

Scolopacidae: sandpipers and relatives

- ☐ Jack Snipe *Lymnocryptes minimus* **PM**
- ☐ Pintail Snipe *Gallinago stenura* **PM**
  Three records, January 1969 Lake Naivasha, October–December 1981 near Mombasa and January 1982 Lake Naivasha
- ☐ African Snipe *Gallinago nigripennis* **am**
- ☐ Great Snipe *Gallinago media* **PM; Near Threatened**
- ☐ Common Snipe *Gallinago gallinago* **PM**
- ☐ Black-tailed Godwit *Limosa limosa* **PM; Near Threatened**
- ☐ Bar-tailed Godwit *Limosa lapponica* **PM**
- ☐ Whimbrel *Numenius phaeopus* **PM**
- ☐ Eurasian Curlew *Numenius arquata* **PM; Near Threatened**
- ☐ Spotted Redshank *Tringa erythropus* **PM**
- ☐ Common Redshank *Tringa totanus* **PM**
- ☐ Marsh Sandpiper *Tringa stagnatilis* **PM**
- ☐ Common Greenshank *Tringa nebularia* **PM**
- ☐ Green Sandpiper *Tringa ochropus* **PM**
- ☐ Wood Sandpiper *Tringa glareola* **PM**
- ☐ Terek Sandpiper *Xenus cinereus* **PM**
- ☐ Common Sandpiper *Actitis hypoleucos* **PM**
- ☐ Spotted Sandpiper *Actitis macularius* **VN**
  The only record is of one, 5 September 1999, Mountain Lodge, Mt Kenya (Bishop & Bishop 2001)
- ☐ Ruddy Turnstone *Arenaria interpres* **PM**
- ☐ Red Knot *Calidris canutus* **PM**
- ☐ Sanderling *Calidris alba* **PM**

☐ Red-necked Stint *Calidris ruficollis* **PM**
Three records: May 1981, north of Malindi; July 1987, Lake Magadi; and August 1989, Mida Creek

☐ Little Stint *Calidris minuta* **PM**

☐ Temminck's Stint *Calidris temminckii* **PM**

☐ Long-toed Stint *Calidris subminuta* **PM**

☐ Pectoral Sandpiper *Calidris melanotos* **VP, VN**
One collected May 1952, Lake Naivasha and one seen, September 1981, Mombasa

☐ Curlew Sandpiper *Calidris ferruginea* **PM**

☐ Dunlin *Calidris alpina* **PM**

☐ Broad-billed Sandpiper *Limicola falcinellus* **PM**

☐ Buff-breasted Sandpiper *Tryngites subruficollis* **VN; Near Threatened**
One seen at Kerio Bay, Lake Turkana on 8 December 1973. The genus name was incorrectly spelled in the 1996 list

☐ Ruff *Philomachus pugnax* **PM**

☐ Red-necked Phalarope *Phalaropus lobatus* **PM**

☐ Grey Phalarope *Phalaropus fulicarius* **PM**
The sole confirmed record is of one on 1 September 1979, Nairobi National Park (OS-c 1981)

Glareolidae: Egyptian Plover, coursers and pratincoles

☐ Egyptian Plover *Pluvianus aegyptius*
Two records, August 1971 and August 2003, near Todenyang, Lake Turkana. It was called the Egyptian-plover in the 1996 list

☐ Cream-coloured Courser *Cursorius cursor* **VP**
The sole record is of several birds seen, January–February 1987, on the east side of Lake Turkana

☐ Somali Courser *Cursorius somalensis*

☐ Temminck's Courser *Cursorius temminckii*

☐ Double-banded Courser *Rhinoptilus africanus*
It was called the Two-banded Courser in the 1996 list

☐ Heuglin's Courser *Rhinoptilus cinctus*

☐ Bronze-winged Courser *Rhinoptilus chalcopterus* **AM**
It was called the Violet-tipped Courser in the 1996 list

☐ Collared Pratincole *Glareola pratincola* **am**
The Palaearctic nominate race winters in Africa but there is no suggestion that any reach Kenya

☐ Black-winged Pratincole *Glareola nordmanni* **PM; Near Threatened**

☐ Madagascar Pratincole *Glareola ocularis* **MM; Vulnerable**

☐ Rock Pratincole *Glareola nuchalis*

Laridae: gulls, terns and skimmers

The genera and nomenclature used follow Sangster *et al.* (2007)

☐ Sooty Gull *Larus hemprichii* **am**

☐ Common Gull *Larus canus* **VP**
The sole record is of an adult, probably of the race *heinei*, seen and photographed, 1 December 2005, Sabaki River Mouth. It was called the Mew Gull by Gill & Wright (2006) and Sangster *et al.* (2007)

☐ Kelp Gull *Larus dominicanus* **VS**
The sole record is of an adult photographed, 2 January 1984, Malindi

- ☐ Lesser Black-backed Gull *Larus fuscus* **PM**
  Following Sangster *et al.* (2007), includes the taxa *heuglini*, *taimyrensis* and *barabensis* which were included under Heuglin's Gull in the 1996 list; Sangster *et al.* no longer recognize Heuglin's Gull as a species
- ☐ Pallas's Gull *Larus ichthyaetus* **PM**
  It was called the Great Black-headed Gull in the 1996 list and by Gill & Wright (2006)
- ☐ Grey-headed Gull *Chroicocephalus cirrocephalus*
  This and the next two species were included in *Larus* in the 1996 list
- ☐ Black-headed Gull *Chroicocephalus ridibundus* **PM**
- ☐ Slender-billed Gull *Chroicocephalus genei* **PM**
- ☐ Gull-billed Tern *Gelochelidon nilotica* **PM**
  It was included in *Sterna* in the 1996 list
- ☐ Caspian Tern *Hydroprogne caspia* **PM**
  It was included in *Sterna* in the 1996 list
- ☐ Lesser Crested Tern *Sterna bengalensis* **am**
- ☐ Sandwich Tern *Sterna sandvicensis* **PM**
- ☐ Swift Tern *Sterna bergii* **am, io**
  It was called the Greater Crested Tern in the 1996 list
- ☐ Roseate Tern *Sterna dougallii*
- ☐ Common Tern *Sterna hirundo* **PM**
- ☐ White-cheeked Tern *Sterna repressa*
- ☐ Little Tern *Sternula albifrons* **PM**
  This and the next species were included in *Sterna* in the 1996 list
- ☐ Saunders's Tern *Sternula saundersi* **RS**
- ☐ Bridled Tern *Onychoprion anaethetus*
  It was included in *Sterna* in the 1996 list
- ☐ Sooty Tern *Onychoprion fuscata*
  It was included in *Sterna* in the 1996 list
- ☐ Whiskered Tern *Chlidonias hybrida* **(pm)**
  It was *hybridus* in the 1996 list; *hybrida* is correct (David & Gosselin 2002a). Palaearctic birds might occur
- ☐ White-winged Black Tern *Chlidonias leucopterus* **PM**
  It was called by the unhelpful name White-winged Tern in the 1996 list and by Gill & Wright (2006)
- ☐ Black Tern *Chlidonias niger* **VP**
  Only three confirmed records: October 1980, Dandora; February 1983, Thika; and February 1985, Ferguson's Gulf, Lake Turkana
- ☐ Brown Noddy *Anous stolidus*
- ☐ Lesser Noddy *Anous tenuirostris* **IO**
- ☐ African Skimmer *Rynchops flavirostris* **Near Threatened**
  It was included in a family of its own (Rynchopidae) in the 1996 list

Stercorariidae: skuas

- ☐ Pomarine Skua *Stercorarius pomarinus* **PM**
- ☐ Arctic Skua *Stercorarius parasiticus* **VP**
- ☐ Long-tailed Skua *Stercorarius longicaudus* **VP**
  The only record is of an adult photographed on 26 August 1961, at Ferguson's Gulf, Lake Turkana

Pteroclidae: sandgrouse

- ☐ Chestnut-bellied Sandgrouse *Pterocles exustus*
- ☐ Yellow-throated Sandgrouse *Pterocles gutturalis*
- ☐ Black-faced Sandgrouse *Pterocles decoratus*
- ☐ Lichtenstein's Sandgrouse *Pterocles lichtensteinii*
- ☐ Four-banded Sandgrouse *Pterocles quadricinctus*

Columbidae: pigeons and doves

- ☐ Feral Pigeon *Columba livia*
- ☐ Speckled Pigeon *Columba guinea*
- ☐ African Olive Pigeon *Columba arquatrix*
  It was called the Olive Pigeon in the 1996 list
- ☐ Eastern Bronze-naped Pigeon *Columba delegorguei*
- ☐ Lemon Dove *Aplopelia larvata*
- ☐ European Turtle Dove *Streptopelia turtur* **VP**
- ☐ Dusky Turtle Dove *Streptopelia lugens*
- ☐ African White-winged Dove *Streptopelia reichenowi* **Near Threatened**
- ☐ African Mourning Dove *Streptopelia decipiens*
- ☐ Red-eyed Dove *Streptopelia semitorquata*
- ☐ Ring-necked Dove *Streptopelia capicola*
- ☐ Laughing Dove *Streptopelia senegalensis*
- ☐ Emerald-spotted Wood Dove *Turtur chalcospilos*
- ☐ Blue-spotted Wood Dove *Turtur afer*
- ☐ Tambourine Dove *Turtur tympanistria*
- ☐ Namaqua Dove *Oena capensis*
- ☐ Bruce's Green Pigeon *Treron waalia*
- ☐ African Green Pigeon *Treron calvus*
  Following David & Gosselin (2002b), *calvus* is correct (it was *calva* in the 1996 list)

Psittacidae: lovebirds and parrots

- ☐ Red-headed Lovebird *Agapornis pullarius*
- ☐ Yellow-collared Lovebird *Agapornis personatus*
  Feral birds continue to breed around Mombasa
- ☐ Fischer's Lovebird *Agapornis fischeri* **Near Threatened**
  Not included in the 1996 list; feral birds have recently been reported in Nairobi National Park
- ☐ Hybrid Lovebird *Agapornis fischeri* × *personatus*
  It was called Yellow-collared × Fischer's Lovebird hybrid in the 1996 list
- ☐ Grey Parrot *Psittacus erithacus* **Near Threatened**
- ☐ Red-fronted Parrot *Poicephalus gulielmi*
- ☐ Meyer's Parrot *Poicephalus meyeri*
  It was called the Brown Parrot in the 1996 list
- ☐ Brown-headed Parrot *Poicephalus cryptoxanthus*
- ☐ African Orange-bellied Parrot *Poicephalus rufiventris*

Musophagidae: turacos

- ☐ Great Blue Turaco *Corythaeola cristata*
- ☐ Schalow's Turaco *Tauraco schalowi*
- ☐ Black-billed Turaco *Tauraco schuetti*
- ☐ Fischer's Turaco *Tauraco fischeri* **Near Threatened**
- ☐ White-crested Turaco *Tauraco leucolophus*
- ☐ Hartlaub's Turaco *Tauraco hartlaubi*
- ☐ Purple-crested Turaco *Tauraco porphyreolophus*
- ☐ Ross's Turaco *Musophaga rossae*
- ☐ Bare-faced Go-away-bird *Corythaixoides personatus*
- ☐ White-bellied Go-away-bird *Corythaixoides leucogaster*
- ☐ Eastern Grey Plantain-eater *Crinifer zonurus*

Cuculidae: cuckoos and coucals

- ☐ Jacobin Cuckoo *Clamator jacobinus* **am, pm, om**
  It was called the Black-and-white Cuckoo *Oxylophus jacobinus* in the 1996 list
- ☐ Levaillant's Cuckoo *Clamator levaillantii* **am**
  It was included in the genus *Oxylophus* in the 1996 list
- ☐ Great Spotted Cuckoo *Clamator glandarius* **am, (pm)**
  Palaearctic migrants may occur
- ☐ Thick-billed Cuckoo *Pachycoccyx audeberti*
- ☐ Red-chested Cuckoo *Cuculus solitarius* **am**
- ☐ Black Cuckoo *Cuculus clamosus* **am**
- ☐ Common Cuckoo *Cuculus canorus* **PM**
  It was called the Eurasian Cuckoo in the 1996 list
- ☐ African Cuckoo *Cuculus gularis* **AM**
- ☐ Lesser Cuckoo *Cuculus poliocephalus* **PM, OM**
  It was called the Asian Lesser Cuckoo in the 1996 list; the present name follows Gill & Wright (2006)
- ☐ Madagascar Cuckoo *Cuculus rochii* **MM**
  It was called the Madagascar Lesser Cuckoo in the 1996 list; the present name follows Gill & Wright (2006)
- ☐ Barred Long-tailed Cuckoo *Cercococcyx montanus*
- ☐ Klaas's Cuckoo *Chrysococcyx klaas*
- ☐ African Emerald Cuckoo *Chrysococcyx cupreus*
- ☐ Diederik Cuckoo *Chrysococcyx caprius* **am**
- ☐ Yellowbill *Ceuthmochares aereus* **am**
- ☐ Black Coucal *Centropus grillii*
- ☐ Blue-headed Coucal *Centropus monachus*
- ☐ Senegal Coucal *Centropus senegalensis*
- ☐ White-browed Coucal *Centropus superciliosus*

Tytonidae: barn owls

- ☐ Grass Owl *Tyto capensis*
  It was called the African Grass Owl in the 1996 list
- ☐ Barn Owl *Tyto alba*

Strigidae: typical owls

- ☐ Sokoke Scops Owl *Otus ireneae* **Endangered**
- ☐ African Scops Owl *Otus senegalensis*
- ☐ Eurasian Scops Owl *Otus scops* **PM**
- ☐ Northern White-faced Scops Owl *Ptilopsis leucotis*
  The 1996 list included this and the next species in *Otus leucotis*
- ☐ Southern White-faced Scops Owl *Ptilopsis granti*
  See note above
- ☐ Cape Eagle Owl *Bubo capensis*
- ☐ Spotted Eagle Owl *Bubo africanus*
- ☐ Greyish Eagle Owl *Bubo cinerascens*
  It was included as a race of *Bubo africanus* in the 1996 list
- ☐ Verreaux's Eagle Owl *Bubo lacteus*
- ☐ Pel's Fishing Owl *Scotopelia peli*
- ☐ African Wood Owl *Strix woodfordii*
- ☐ Pearl-spotted Owlet *Glaucidium perlatum*
- ☐ Red-chested Owlet *Glaucidium tephronotum*
- ☐ African Barred Owlet *Glaucidium capense*
- ☐ Abyssinian Owl *Asio abyssinicus*
  The sole confirmed record is of one collected on 10 September 1961 at 3350 m, on Mt Kenya. It was called the African Long-eared Owl in the 1996 list
- ☐ Short-eared Owl *Asio flammeus* **VP**
- ☐ Marsh Owl *Asio capensis*

Caprimulgidae: nightjars

- ☐ Eurasian Nightjar *Caprimulgus europaeus* **PM**
- ☐ Dusky Nightjar *Caprimulgus fraenatus*
- ☐ Nubian Nightjar *Caprimulgus nubicus* **AM**
- ☐ Donaldson Smith's Nightjar *Caprimulgus donaldsoni*
  The American zoologist and collector Dr Arthur Donaldson Smith (1864–1939) had no hyphen in his name
- ☐ Fiery-necked Nightjar *Caprimulgus pectoralis*
  The taxonomy of this and the next species is complex and controversial. The race *C. p. nigriscapularis* is called the Black-shouldered Nightjar and is considered a separate species by some
- ☐ Montane Nightjar *Caprimulgus poliocephalus*
- ☐ Swamp Nightjar *Caprimulgus natalensis*
  It was called the African White-tailed Nightjar in the 1996 list
- ☐ Plain Nightjar *Caprimulgus inornatus* **AM, (pm)**
- ☐ Star-spotted Nightjar *Caprimulgus stellatus*
- ☐ Freckled Nightjar *Caprimulgus tristigma*

- ☐ Long-tailed Nightjar *Caprimulgus climacurus* **am**
- ☐ Slender-tailed Nightjar *Caprimulgus clarus*
- ☐ Gabon Nightjar *Caprimulgus fossii* **am**
  It was called the Square-tailed Nightjar or Mozambique Nightjar by some authors
- ☐ Standard-winged Nightjar *Macrodipteryx longipennis* **AM**
- ☐ Pennant-winged Nightjar *Macrodipteryx vexillarius* **AM**

Apodidae: swifts

- ☐ Scarce Swift *Schoutedenapus myoptilus*
- ☐ Böhm's Spinetail *Neafrapus boehmi*
- ☐ Mottled Spinetail *Telacanthura ussheri* — was *Telecanthura* in the original edition
- ☐ Sabine's Spinetail *Rhaphidura sabini* **Ex**
- ☐ African Palm Swift *Cypsiurus parvus*
- ☐ Alpine Swift *Tachymarptis melba*
  This and the next species were included in *Apus* in the 1996 list. There is no evidence that Palaearctic birds reach Kenya
- ☐ Mottled Swift *Tachymarptis aequatorialis*
- ☐ Common Swift *Apus apus* **PM**
  It was called the Eurasian Swift in the 1996 list
- ☐ Nyanza Swift *Apus niansae*
- ☐ African Black Swift *Apus barbatus*
- ☐ Forbes-Watson's Swift *Apus berliozi* **AM**
- ☐ Little Swift *Apus affinis*
- ☐ Horus Swift *Apus horus* **am**
- ☐ White-rumped Swift *Apus caffer*

Coliidae: mousebirds

- ☐ Speckled Mousebird *Colius striatus*
  Colius striatus kikuyuensis **E**
- ☐ White-headed Mousebird *Colius leucocephalus*
  Colius leucocephalus turneri **E**
- ☐ Blue-naped Mousebird *Urocolius macrourus*

Trogonidae: trogons

- ☐ Narina Trogon *Apaloderma narina*
- ☐ Bar-tailed Trogon *Apaloderma vittatum*

Coraciidae: rollers

- ☐ Rufous-crowned Roller *Coracias naevius* **am**
  The specific epithets changed to *-us* (from *-a*) endings following David & Gosselin (2002b) for this and the next two species
- ☐ Lilac-breasted Roller *Coracias caudatus* **am**
- ☐ Abyssinian Roller *Coracias abyssinicus*

- ☐ Eurasian Roller *Coracias garrulus* **PM; Near Threatened**
- ☐ Broad-billed Roller *Eurystomus glaucurus* **am, mm**

Alcedinidae: kingfishers

- ☐ Grey-headed Kingfisher *Halcyon leucocephala* **am**
- ☐ Brown-hooded Kingfisher *Halcyon albiventris*
- ☐ Striped Kingfisher *Halcyon chelicuti*
- ☐ Blue-breasted Kingfisher *Halcyon malimbica*
  The only records are of 2–3 birds, 29 November 1993 – mid July 1996, Busia District, western Kenya
- ☐ Woodland Kingfisher *Halcyon senegalensis* **am**
- ☐ Mangrove Kingfisher *Halcyon senegaloides*
- ☐ African Pygmy Kingfisher *Ceyx pictus* **am**
  It was called *Ispidina picta* in the 1996 list
- ☐ Malachite Kingfisher *Alcedo cristata*
- ☐ Shining-blue Kingfisher *Alcedo quadribrachys*
- ☐ Semi-collared Kingfisher *Alcedo semitorquata*
  It was called the Half-collared Kingfisher in the 1996 list
- ☐ Giant Kingfisher *Megaceryle maxima*
- ☐ Pied Kingfisher *Ceryle rudis*

Meropidae: bee-eaters

- ☐ Blue-headed Bee-eater *Merops muelleri*
- ☐ Swallow-tailed Bee-eater *Merops hirundineus*
  The sole record is of one seen, 21 May 2000, Arabuko-Sokoke Forest
- ☐ Little Bee-eater *Merops pusillus*
- ☐ Blue-breasted Bee-eater *Merops variegatus*
- ☐ Cinnamon-chested Bee-eater *Merops oreobates*
- ☐ White-fronted Bee-eater *Merops bullockoides*
- ☐ Somali Bee-eater *Merops revoilii*
- ☐ White-throated Bee-eater *Merops albicollis* **AM**
- ☐ Blue-cheeked Bee-eater *Merops persicus* **PM**
- ☐ Madagascar Bee-eater *Merops superciliosus* **am, mm**
- ☐ Eurasian Bee-eater *Merops apiaster* **PM**
- ☐ Northern Carmine Bee-eater *Merops nubicus* **AM**
- ☐ Southern Carmine Bee-eater *Merops nubicoides* **AM**
  Included in *Merops nubicus* in the 1996 list and by Dickinson (2003); Gill & Wright (2006) adopt the split

Upupidae: Hoopoe

- ☐ Hoopoe *Upupa epops* **am, pm**
  Includes the African Hoopoe, comprising the races *africana*, *waibeli* and *senegalensis*; the nominate race is a Palaearctic migrant

Phoeniculidae: wood-hoopoes

- ☐ Forest Wood-hoopoe *Phoeniculus castaneiceps* **Ex**
  Last recorded pre-1915; see Zimmerman *et al.* (1996) for an explanation

- ☐ White-headed Wood-hoopoe *Phoeniculus bollei*
- ☐ Green Wood-hoopoe *Phoeniculus purpureus*
- ☐ Black-billed Wood-hoopoe *Phoeniculus somaliensis*
- ☐ Violet Wood-hoopoe *Phoeniculus damarensis*
- ☐ Common Scimitarbill *Rhinopomastus cyanomelas*
- ☐ Abyssinian Scimitarbill *Rhinopomastus minor*

Bucerotidae: hornbills

- ☐ Crowned Hornbill *Tockus alboterminatus*
- ☐ Hemprich's Hornbill *Tockus hemprichii*
- ☐ African Grey Hornbill *Tockus nasutus*
- ☐ Red-billed Hornbill *Tockus erythrorhynchus*
- ☐ Eastern Yellow-billed Hornbill *Tockus flavirostris*
- ☐ Von der Decken's Hornbill *Tockus deckeni*
- ☐ Jackson's Hornbill *Tockus jacksoni*
  Considered a race (*jacksoni*) of Von der Decken's Hornbill by several authors
- ☐ Trumpeter Hornbill *Bycanistes bucinator*
- ☐ Black-and-white Casqued Hornbill *Bycanistes subcylindricus*
  It was called the Black-and-white-casqued Hornbill in the 1996 list
- ☐ Silvery-cheeked Hornbill *Bycanistes brevis* **am**

Bucorvidae: ground hornbills

This family was included in the Bucerotidae in the 1996 list

- ☐ Abyssinian Ground Hornbill *Bucorvus abyssinicus*
- ☐ Southern Ground Hornbill *Bucorvus leadbeateri*

Capitonidae: barbets and tinkerbirds

- ☐ Grey-throated Barbet *Gymnobucco bonapartei*
- ☐ White-eared Barbet *Stactolaema leucotis*
- ☐ Green Barbet *Stactolaema olivacea*
- ☐ Speckled Tinkerbird *Pogoniulus scolopaceus* **Ex**
  Last recorded pre- 1939; see Zimmerman *et al.* (1996) for an explanation
- ☐ Green Tinkerbird *Pogoniulus simplex*
  It was called the Eastern Green Tinkerbird in the 1996 list
- ☐ Moustached Tinkerbird *Pogoniulus leucomystax*
  It was called the Moustached Green Tinkerbird in the 1996 list
- ☐ Yellow-rumped Tinkerbird *Pogoniulus bilineatus*
- ☐ Red-fronted Tinkerbird *Pogoniulus pusillus*
- ☐ Yellow-fronted Tinkerbird *Pogoniulus chrysoconus*
- ☐ Yellow-spotted Barbet *Buccanodon duchaillui*
- ☐ Hairy-breasted Barbet *Tricholaema hirsuta*
- ☐ Red-fronted Barbet *Tricholaema diadematа*
- ☐ Spot-flanked Barbet *Tricholaema lacrymosa*

- ☐ Black-throated Barbet *Tricholaema melanocephala*
- ☐ White-headed Barbet *Lybius leucocephalus*
  Lybius leucocephalus senex **E**
- ☐ Black-billed Barbet *Lybius guifsobalito*
- ☐ Black-collared Barbet *Lybius torquatus*
- ☐ Brown-breasted Barbet *Lybius melanopterus*
- ☐ Double-toothed Barbet *Lybius bidentatus*
- ☐ Yellow-billed Barbet *Trachylaemus purpuratus*
- ☐ Red-and-yellow Barbet *Trachyphonus erythrocephalus*
- ☐ D'Arnaud's Barbet *Trachyphonus darnaudii*
  Formerly, some authors considered the race *usambiro* to be a separate species, the Usambiro Barbet

Indicatoridae: honeyguides

- ☐ Cassin's Honeybird *Prodotiscus insignis*
- ☐ Green-backed Honeybird *Prodotiscus zambesiae*
  It was called the Eastern Honeybird in the 1996 list
- ☐ Wahlberg's Honeybird *Prodotiscus regulus*
- ☐ Pallid Honeyguide *Indicator meliphilus*
- ☐ Least Honeyguide *Indicator exilis*
- ☐ Lesser Honeyguide *Indicator minor*
  Includes the Thick-billed Honeyguide *Indicator minor conirostris* which was treated as a separate species in the 1996 list
- ☐ Scaly-throated Honeyguide *Indicator variegatus*
- ☐ Greater Honeyguide *Indicator indicator*

Picidae: wrynecks and woodpeckers

- ☐ Eurasian Wryneck *Jynx torquilla* **PM**
- ☐ Red-throated Wryneck *Jynx ruficollis*
- ☐ Nubian Woodpecker *Campethera nubica*
- ☐ Golden-tailed Woodpecker *Campethera abingoni*
- ☐ Mombasa Woodpecker *Campethera mombassica*
- ☐ Green-backed Woodpecker *Campethera cailliautii*
- ☐ Bennett's Woodpecker *Campethera bennettii* **Ex**
  The sole record is of an adult male (specimen), 20 July 1918, from near Mombasa
- ☐ Fine-banded Woodpecker *Campethera tullbergi*
  Campethera tullbergi hausburgi **E**
- ☐ Buff-spotted Woodpecker *Campethera nivosa*
- ☐ Brown-eared Woodpecker *Campethera caroli*
- ☐ Speckle-breasted Woodpecker *Dendropicos poecilolaemus*
- ☐ Cardinal Woodpecker *Dendropicos fuscescens*
- ☐ Bearded Woodpecker *Dendropicos namaquus*
- ☐ Yellow-crested Woodpecker *Dendropicos xantholophus*
- ☐ African Grey Woodpecker *Dendropicos goertae*
  Called the Grey Woodpecker in the 1996 list. The race *rhodeogaster* is considered a separate species by some
- ☐ Brown-backed Woodpecker *Picoides obsoletus*

Eurylaimidae: broadbills

☐ African Broadbill *Smithornis capensis*

Pittidae: pittas

☐ African Pitta *Pitta angolensis* **AM**

Platysteiridae: batises, wattle-eyes and relatives

☐ Shrike Flycatcher *Megabyas flammulatus*
It was included in *Bias* and called the African Shrike-Flycatcher in the 1996 list

☐ Black-and-white Flycatcher *Bias musicus*

☐ Chestnut Wattle-eye *Dyaphorophyia castanea*

☐ Jameson's Wattle-eye *Dyaphorophyia jamesoni*

☐ Yellow-bellied Wattle-eye *Dyaphorophyia concreta*

☐ Forest Batis *Batis mixta*
Batis mixta ultima **E**

☐ Chin-spot Batis *Batis molitor*

☐ Grey-headed Batis *Batis orientalis*
The sole record is of two specimens taken on 24 October 1910, at Moyale

☐ Pale Batis *Batis soror*

☐ Black-headed Batis *Batis minor*

☐ Pygmy Batis *Batis perkeo*

☐ Brown-throated Wattle-eye *Platysteira cyanea*
It was called the Common Wattle-eye in the 1996 list

☐ Black-throated Wattle-eye *Platysteira peltata*

Malaconotidae: helmetshrikes, bushshrikes, tchagras and puffbacks

The first four species were included in the family Prionopidae (and called helmet-shrikes) in the 1996 list

☐ White-crested Helmetshrike *Prionops plumatus* **am**

☐ Grey-crested Helmetshrike *Prionops poliolophus* **Near Threatened**

☐ Retz's Helmetshrike *Prionops retzii*

☐ Chestnut-fronted Helmetshrike *Prionops scopifrons*
Prionops scopifrons keniensis **E**

The following seven species had Bush-Shrike in their English names in the 1996 list

☐ Grey-headed Bushshrike *Malaconotus blanchoti*

☐ Black-fronted Bushshrike *Chlorophoneus nigrifrons*
This and the next four species were included in *Malaconotus* in the 1996 list

☐ Bocage's Bushshrike *Chlorophoneus bocagei*

☐ Sulphur-breasted Bushshrike *Chlorophoneus sulfureopectus*

☐ Gorgeous Bushshrike *Chlorophoneus viridis*
This was called the Four-coloured Bush-Shrike *Malaconotus quadricolor* in the 1996 list but with the shift to *Chlorophoneus*, the specific epithet *viridis* has precedence

☐ Doherty's Bushshrike *Chlorophoneus dohertyi*

☐ Rosy-patched Bushshrike *Rhodophoneus cruentus*

- ☐ Marsh Tchagra *Tchagra minutus*
  It was called *Bocagia minuta* by some authors
- ☐ Brown-crowned Tchagra *Tchagra australis*
- ☐ Three-streaked Tchagra *Tchagra jamesi*
  Tchagra jamesi mandanus **E**
- ☐ Black-crowned Tchagra *Tchagra senegalus*
  The specific epithet's gender change (from *-a*) follows David & Gosselin (2002b)
- ☐ Pink-footed Puffback *Dryoscopus angolensis*
- ☐ Black-backed Puffback *Dryoscopus cubla*
- ☐ Pringle's Puffback *Dryoscopus pringlii*
- ☐ Northern Puffback *Dryoscopus gambensis*
- ☐ Sooty Boubou *Laniarius leucorhynchus* **Ex**
  The sole record is of one collected in April 1931, in the Kaimosi Forest, western Kenya
- ☐ Slate-coloured Boubou *Laniarius funebris*
- ☐ Lühder's Bushshrike *Laniarius luehderi*
  This and the next species had Bush-Shrike in their English names in the 1996 list
- ☐ Red-naped Bushshrike *Laniarius ruficeps*
- ☐ Tropical Boubou *Laniarius aethiopicus*
- ☐ Papyrus Gonolek *Laniarius mufumbiri* **Near Threatened**
- ☐ Black-headed Gonolek *Laniarius erythrogaster*
- ☐ Brubru *Nilaus afer*

Campephagidae: cuckooshrikes

These were called cuckoo-shrikes in the 1996 list

- ☐ Grey Cuckooshrike *Coracina caesia*
- ☐ White-breasted Cuckooshrike *Coracina pectoralis*
  The only records in the last fifty years are of two pairs, September–October 1993, at 1800m, northeast Mt Elgon, alongside the Uganda border
- ☐ Black Cuckooshrike *Campephaga flava* **am**
- ☐ Red-shouldered Cuckooshrike *Campephaga phoenicea*
- ☐ Petit's Cuckooshrike *Campephaga petiti*
- ☐ Purple-throated Cuckooshrike *Campephaga quiscalina*

Laniidae: shrikes

- ☐ Yellow-billed Shrike *Corvinella corvina*
- ☐ Magpie Shrike *Urolestes melanoleucus*
- ☐ Northern White-crowned Shrike *Eurocephalus rueppelli*
  It was included in the helmetshrikes (Prionopidae) in the 1996 list
- ☐ Red-backed Shrike *Lanius collurio* **PM**
- ☐ Isabelline Shrike *Lanius isabellinus* **PM**
  It was called the Red-tailed Shrike in the 1996 list
- ☐ Mackinnon's Shrike *Lanius mackinnoni*
  It was called Mackinnon's Fiscal in the 1996 list
- ☐ Lesser Grey Shrike *Lanius minor* **PM**

- ☐ Southern Grey Shrike *Lanius meridionalis* **PM**
  The only record is of one seen on 17 February 1988, from near Kibish in the Ilemi Triangle (Pearson *et al.* 1989). It was called the Great Grey Shrike *Lanius excubitor* in the 1996 list
- ☐ Grey-backed Fiscal *Lanius excubitoroides*
- ☐ Long-tailed Fiscal *Lanius cabanisi*
- ☐ Taita Fiscal *Lanius dorsalis*
- ☐ Somali Fiscal *Lanius somalicus*
- ☐ Common Fiscal *Lanius collaris*
- ☐ Woodchat Shrike *Lanius senator* **PM**
- ☐ Masked Shrike *Lanius nubicus* **PM**

Oriolidae: orioles

- ☐ Eurasian Golden Oriole *Oriolus oriolus* **PM**
- ☐ African Golden Oriole *Oriolus auratus* **AM**
- ☐ Green-headed Oriole *Oriolus chlorocephalus*
- ☐ Western Oriole *Oriolus brachyrhynchus*
  It was called the Western Black-headed Oriole in the 1996 list
- ☐ Montane Oriole *Oriolus percivali*
- ☐ Black-headed Oriole *Oriolus larvatus*

Dicruridae: drongos

- ☐ Square-tailed Drongo *Dicrurus ludwigii*
- ☐ Common Drongo *Dicrurus adsimilis*
- ☐ Velvet-mantled Drongo *Dicrurus modestus*

Monarchidae: monarch flycatchers

- ☐ Blue-mantled Crested Flycatcher *Trochocercus cyanomelas*
- ☐ Little Yellow Flycatcher *Erythrocercus holochlorus*
- ☐ Red-bellied Paradise Flycatcher *Terpsiphone rufiventer* **Ex**
  Now only hybrid *rufiventer* × *viridis* occur
- ☐ African Paradise Flycatcher *Terpsiphone viridis* **am**
- ☐ African Blue Flycatcher *Elminia longicauda*
- ☐ Dusky Crested Flycatcher *Elminia nigromitrata*
  It was called *Trochocercus nigromitratus* in the 1996 list
- ☐ White-tailed Crested Flycatcher *Elminia albonotata*
  It was called *Trochocercus albonotatus* in the 1996 list

Corvidae: crows and allies

- ☐ Piapiac *Ptilostomus afer*
- ☐ House Crow *Corvus splendens*
- ☐ Cape Rook *Corvus capensis*
  Some favour the name Cape Crow
- ☐ Pied Crow *Corvus albus*

☐ Dwarf Raven *Corvus edithae*
It was called the Brown-necked Raven *Corvus (ruficollis) edithae* in the 1996 list; called the Somali Crow by some

☐ Fan-tailed Raven *Corvus rhipidurus*

☐ White-necked Raven *Corvus albicollis*
It was called the White-naped Raven in the 1996 list

Paridae: tits

☐ Dusky Tit *Parus funereus*

☐ Northern Black Tit *Parus leucomelas*

☐ White-bellied Tit *Parus albiventris*

☐ Red-throated Tit *Parus fringillinus*

☐ Northern Grey Tit *Parus thruppi*

Remizidae: penduline tits

☐ Mouse-coloured Penduline Tit *Anthoscopus musculus*

☐ Grey Penduline Tit *Anthoscopus caroli*
It was called the African Penduline Tit in the 1996 list

Hirundinidae: saw-wings, swallows and martins

☐ White-headed Saw-wing *Psalidoprocne albiceps*

☐ Black Saw-wing *Psalidoprocne pristoptera*
It was called *Psalidoprocne holomelas* in the 1996 list; *pristoptera* has priority (Sheldon & Winkler (1993)

☐ Grey-rumped Swallow *Pseudhirundo griseopyga* **am**

☐ Mascarene Martin *Phedina borbonica* **MM**
Four records only: June 1978 and June 1993, Lake Jipe; and June 1980 and August 1989 from near Malindi

☐ Plain Martin *Riparia paludicola* **am**

☐ Sand Martin *Riparia riparia* **PM**

☐ Banded Martin *Riparia cincta* **am**

☐ Barn Swallow *Hirundo rustica* **PM**

☐ Angola Swallow *Hirundo angolensis*

☐ Ethiopian Swallow *Hirundo aethiopica*

☐ Wire-tailed Swallow *Hirundo smithii*

☐ Blue Swallow *Hirundo atrocaerulea* **AM; Vulnerable**

☐ Rock Martin *Ptyonoprogne fuligula*
It was included in *Hirundo* in the 1996 list

☐ Common House Martin *Delichon urbicum* **PM**
The species epithet was *urbica* in the 1996 list but was changed to the neuter form following David & Gosselin (2002b)

☐ Lesser Striped Swallow *Cecropis abyssinica*
This and the next three species were included in *Hirundo* in the 1996 list

☐ Rufous-chested Swallow *Cecropis semirufa*

☐ Mosque Swallow *Cecropis senegalensis*

☐ Red-rumped Swallow *Cecropis daurica*

☐ South African Cliff Swallow *Petrochelidon spilodera* **VSA**
An adult seen and photographed on 22 July 2000 at Aruba Dam, Tsavo East National Park is the sole record

Alaudidae: larks

☐ Singing Bush Lark *Mirafra cantillans*
☐ White-tailed Lark *Mirafra albicauda*
☐ Williams's Lark *Mirafra williamsi* **E; Data Deficient**
☐ Friedmann's Lark *Mirafra pulpa* **AM; Data Deficient**
☐ Rufous-naped Lark *Mirafra africana*
Mirafra africana harterti **E**
☐ Red-winged Lark *Mirafra hypermetra*
☐ Flappet Lark *Mirafra rufocinnamomea*
☐ Fawn-coloured Lark *Mirafra africanoides*
☐ Collared Lark *Mirafra collaris*
The only records in the last fifty years are of two specimens, 14 August 1962, near Habaswein; a few seen in August 1970, near the Bura Irrigation Scheme, Tana River; and one seen on 2 November 1974, *c.* 50 km south of Mado Gashe
☐ Gillett's Lark *Mirafra gilletti*
Several specimens, May 1901 from near Mandera and El Wak; and two seen on 6 September 2000 between Mandera and Ramu have been accepted by the East African Rarities Committee (EARC)
☐ Pink-breasted Lark *Mirafra poecilosterna*
☐ Greater Short-toed Lark *Calandrella brachydactyla* **PM**
The sole record is of a specimen, 19 December 1964, from Diani. An earlier specimen (18 November 1899, Athi River) has been re-identified as *Calandrella cinerea saturatior*
☐ Red-capped Lark *Calandrella cinerea*
Calandrella cinerea williamsi **E**
☐ Somali Short-toed Lark *Calandrella somalica*
☐ Masked Lark *Spizocorys personata*
Spizocorys personata intensa **E**, *Spizocorys personata mcchesneyi* **E**
☐ Short-tailed Lark *Pseudalaemon fremantlii*
☐ Crested Lark *Galerida cristata*
☐ Thekla Lark *Galerida theklae*
☐ Chestnut-backed Sparrow Lark *Eremopterix leucotis*
This and the next two species had Sparrow-Lark in their English names in the 1996 list
☐ Chestnut-headed Sparrow Lark *Eremopterix signatus*
The genus is masculine (David & Gosselin 2002b), hence the *-us* ending for the adjective *signatus*
☐ Fischer's Sparrow Lark *Eremopterix leucopareia*
Although the genus is masculine (David & Gosselin 2002b), the species epithet *leucopareia*, not being an adjective, must retain the feminine *-a* ending

Cisticolidae: cisticolas and allies

The following 53 species were included in the Sylviidae in the 1996 list
☐ Red-faced Cisticola *Cisticola erythrops*
☐ Singing Cisticola *Cisticola cantans*
☐ Whistling Cisticola *Cisticola lateralis*

- ☐ Trilling Cisticola *Cisticola woosnami*
- ☐ Chubb's Cisticola *Cisticola chubbi*
- ☐ Hunter's Cisticola *Cisticola hunteri*
- ☐ Rock-loving Cisticola *Cisticola aberrans*
  It was called the Rock Cisticola in the 1996 list
- ☐ Rattling Cisticola *Cisticola chiniana*
- ☐ Boran Cisticola *Cisticola bodessa*
- ☐ Ashy Cisticola *Cisticola cinereolus*
  Within the genus *Cisticola*, many specific epithets have been changed from the feminine *-a* ending to the masculine *-us*, following David & Gosselin (2002b) who show that *Cisticola* is masculine; however, those still retaining the *-a* ending in the present list are indeed correct
- ☐ Tana River Cisticola *Cisticola restrictus* **E; Ex; Data Deficient**
  Known only from seven variable specimens, the last one collected on 4 September 1972, from the Lower Tana River. There is currently considerable doubt concerning the validity of this taxon
- ☐ Wailing Cisticola *Cisticola lais*
  It was called Lynes's Cisticola *Cisticola (lais) distinctus* in the 1996 list
- ☐ Winding Cisticola *Cisticola galactotes*
- ☐ Coastal Cisticola *Cisticola haematocephalus*
  This distinctive bird, originally described as a species by Cabanis (1868), was considered a race of *Cisticola galactotes* in the 1996 list, but here, in line with Gill & Wright (2006), we give it specific status
- ☐ Carruthers's Cisticola *Cisticola carruthersi*
- ☐ Levaillant's Cisticola *Cisticola tinniens*
  Cisticola tinniens oreophilus **E**
- ☐ Stout Cisticola *Cisticola robustus*
- ☐ Aberdare Cisticola *Cisticola aberdare* **E; Endangered**
- ☐ Croaking Cisticola *Cisticola natalensis*
- ☐ Red-pate Cisticola *Cisticola ruficeps*
  The sole record is of two seen on 13 August 1990, near Lokichokio, northwestern Kenya
- ☐ Tiny Cisticola *Cisticola nanus*
- ☐ Siffling Cisticola *Cisticola brachypterus*
  Cisticola brachypterus kericho **E**
- ☐ Foxy Cisticola *Cisticola troglodytes*
- ☐ Long-tailed Cisticola *Cisticola angusticauda*
- ☐ Zitting Cisticola *Cisticola juncidis*
- ☐ Desert Cisticola *Cisticola aridulus*
- ☐ Black-backed Cisticola *Cisticola eximius*
- ☐ Pectoral-patch Cisticola *Cisticola brunnescens*
- ☐ Wing-snapping Cisticola *Cisticola ayresii*
  Cisticola ayresii mauensis **E**
- ☐ Tawny-flanked Prinia *Prinia subflava*
- ☐ Pale Prinia *Prinia somalica*
- ☐ River Prinia *Prinia fluviatilis*
  Not in the 1996 list. The first and only record was of several birds, 8 August 2001, near Lokichokio, northwestern Kenya (Finch 2005)

- ☐ Banded Prinia *Prinia bairdii*
  Prinia bairdii melanops **E**
- ☐ White-chinned Prinia *Schistolais leucopogon*
  It was included in *Prinia* in the 1996 list; 'Prinia' is kept in its English name following Gill & Wright (2006)
- ☐ Red-winged Warbler *Heliolais erythropterus*
  It was *Heliolais erythroptera* in the 1996 list, gender change follows David & Gosselin (2002b)
- ☐ Buff-bellied Warbler *Phyllolais pulchella*
- ☐ Bar-throated Apalis *Apalis thoracica*
  Apalis thoracica fuscigularis **E; Critically Endangered;** the Taita Apalis is regarded as a separate species by some
- ☐ Black-collared Apalis *Apalis pulchra*
- ☐ Yellow-breasted Apalis *Apalis flavida*
  Apalis flavida pugnax **E**
  Apalis flavida flavocincta (the Brown-tailed Apalis) is considered a separate species by some
- ☐ Black-throated Apalis *Apalis jacksoni*
- ☐ White-winged Apalis *Apalis chariessa* **Vulnerable**
  Apalis chariessa chariessa **E; Ex.** Known only from the type, 13 September 1878, and another specimen, August 1961, both from the lower Tana River
- ☐ Black-headed Apalis *Apalis melanocephala*
  Apalis melanocephala nigrodorsalis **E**
- ☐ Chestnut-throated Apalis *Apalis porphyrolaema*
- ☐ Buff-throated Apalis *Apalis rufogularis*
- ☐ Karamoja Apalis *Apalis karamojae* **Vulnerable**
  Added to the Kenya list since the 3rd edition. From 2004, a small population has been resident in the eastern Masai Mara (EARC 2007)
- ☐ Grey Apalis *Apalis cinerea*
- ☐ Brown-headed Apalis *Apalis alticola*
- ☐ Red-fronted Warbler *Urorhipis rufifrons*
  It was called *Spiloptila rufifrons* in the 1996 list*; Urorhipis rufifrons rufidorsalis* **E**
- ☐ Grey-capped Warbler *Eminia lepida*
- ☐ Grey-backed Camaroptera *Camaroptera brachyura*
  The grey-backed subspecies are sometimes regarded as the species *C. brevicaudata*, while those races with green backs are restricted to *C. brachyura* (the Green-backed Camaroptera)
- ☐ Olive-green Camaroptera *Camaroptera chloronota*
- ☐ Grey Wren Warbler *Calamonastes simplex*
- ☐ Pale Wren Warbler *Calamonastes undosus*

Pycnonotidae: bulbuls

- ☐ Common Bulbul *Pycnonotus barbatus*
  The small race *dodsoni* is sometimes considered as a separate species, the White-eared Bulbul
- ☐ Shelley's Greenbul *Andropadus masukuensis*
- ☐ Mountain Greenbul *Andropadus nigriceps*
- ☐ Stripe-cheeked Greenbul *Andropadus milanjensis*
- ☐ Little Greenbul *Andropadus virens*
- ☐ Little Grey Greenbul *Andropadus gracilis*
  No documented records since 1967

☐ Ansorge's Greenbul *Andropadus ansorgei*
Andropadus ansorgei kavirondensis **E**

☐ Plain Greenbul *Andropadus curvirostris*
It was called the Cameroon Sombre Greenbul in the 1996 list

☐ Yellow-whiskered Greenbul *Andropadus latirostris*

☐ Slender-billed Greenbul *Andropadus gracilirostris*
Andropadus gracilirostris percivali **E**

☐ Zanzibar Greenbul *Andropadus importunus*
It was called the Zanzibar Sombre Greenbul in the 1996 list

☐ Honeyguide Greenbul *Baeopogon indicator*

☐ Joyful Greenbul *Chlorocichla laetissima*

☐ Yellow-bellied Greenbul *Chlorocichla flaviventris*

☐ Yellow-throated Leaflove *Chlorocichla flavicollis*

☐ Terrestrial Brownbul *Phyllastrephus terrestris*

☐ Northern Brownbul *Phyllastrephus strepitans*

☐ Grey-olive Greenbul *Phyllastrephus cerviniventris*

☐ Toro Olive Greenbul *Phyllastrephus hypochloris*

☐ Fischer's Greenbul *Phyllastrephus fischeri*

☐ Cabanis's Greenbul *Phyllastrephus cabanisi*

☐ Yellow-streaked Greenbul *Phyllastrephus flavostriatus* **Ex**
The only records are of two specimens, 18 November 1917 from near Murang'a, and November 1938 from Mt Kasigau, south of Voi

☐ Tiny Greenbul *Phyllastrephus debilis*

☐ Red-tailed Bristlebill *Bleda syndactyla*

☐ Eastern Nicator *Nicator gularis*

Sylviidae: Old World warblers

☐ Fan-tailed Grassbird *Schoenicola brevirostris*
It was called the Broad-tailed Warbler in the 1996 list

☐ Little Rush Warbler *Bradypterus baboecala*

☐ White-winged Swamp Warbler *Bradypterus carpalis*
It was called the White-winged Warbler in the 1996 list

☐ Evergreen Forest Warbler *Bradypterus lopezi*

☐ Cinnamon Bracken Warbler *Bradypterus cinnamomeus*

☐ Black-faced Rufous Warbler *Bathmocercus rufus*

☐ Moustached Grass Warbler *Melocichla mentalis*
It was called the African Moustached Warbler in the 1996 list

☐ Common Grasshopper Warbler *Locustella naevia* **VP**
It was called simply Grasshopper Warbler in the 1996 list. Two records, one ringed on 19 June [*sic*] 1977, Nguruman (Fayad & Fayad 1977) and another ringed on 8 March 2000, Marsabit (Borghesio *et al*. 2004)

☐ River Warbler *Locustella fluviatilis* **PM**

☐ Savi's Warbler *Locustella luscinioides* **VP**
Two records, of single birds caught, photographed and ringed at Ngulia, 6 December 1975 and 20 December 1987 (Backhurst & Pearson 1976, Pearson *et al.* 1988)

- ☐ Basra Reed Warbler *Acrocephalus griseldis* **PM; Endangered**
- ☐ Greater Swamp Warbler *Acrocephalus rufescens*
- ☐ Lesser Swamp Warbler *Acrocephalus gracilirostris*
- ☐ Great Reed Warbler *Acrocephalus arundinaceus* **PM**
- ☐ Sedge Warbler *Acrocephalus schoenobaenus* **PM**
- ☐ African Reed Warbler *Acrocephalus baeticatus* **am**
- ☐ Eurasian Reed Warbler *Acrocephalus scirpaceus* **PM**
- ☐ Marsh Warbler *Acrocephalus palustris* **PM**
- ☐ Eastern Olivaceous Warbler *Hippolais pallida* **PM**
  The qualifier 'Eastern' is needed now that the hitherto included race *opaca* has been split off as a species, the Western Olivaceous Warbler *Hippolais opaca*
- ☐ Upcher's Warbler *Hippolais languida* **PM**
- ☐ Olive-tree Warbler *Hippolais olivetorum* **PM**
- ☐ Icterine Warbler *Hippolais icterina* **PM**
- ☐ Dark-capped Yellow Warbler *Chloropeta natalensis*
- ☐ Mountain Yellow Warbler *Chloropeta similis*
- ☐ Papyrus Yellow Warbler *Chloropeta gracilirostris* **Vulnerable**
- ☐ Kretschmer's Longbill *Macrosphenus kretschmeri* **Ex**
  Last recorded on 17 September 1920 in Kitovu Forest, Taveta District
- ☐ Yellow-bellied Hyliota *Hyliota flavigaster*
- ☐ Southern Hyliota *Hyliota australis*
- ☐ Green Hylia *Hylia prasina*
- ☐ Yellow-throated Woodland Warbler *Phylloscopus ruficapilla*
  It was *Phylloscopus ruficapillus* in the 1996 list; gender change follows David & Gosselin (2002a)
- ☐ Uganda Woodland Warbler *Phylloscopus budongoensis*
- ☐ Brown Woodland Warbler *Phylloscopus umbrovirens*
- ☐ Willow Warbler *Phylloscopus trochilus* **PM**
- ☐ Common Chiffchaff *Phylloscopus collybita* **PM**
  It was called simply the Chiffchaff in the 1996 list
- ☐ Wood Warbler *Phylloscopus sibilatrix* **PM**
- ☐ Yellow-bellied Eremomela *Eremomela icteropygialis*
- ☐ Yellow-vented Eremomela *Eremomela flavicrissalis*
- ☐ Green-backed Eremomela *Eremomela canescens*
  This was included within *Eremomela pusilla* in the 1996 list
- ☐ Green-capped Eremomela *Eremomela scotops*
- ☐ Turner's Eremomela *Eremomela turneri* **Endangered**
  Eremomela turneri turneri **E.** Considered conspecific with the Brown-capped Eremomela *E. badiceps* by some
- ☐ Northern Crombec *Sylvietta brachyura*
- ☐ Red-faced Crombec *Sylvietta whytii*
- ☐ Somali Crombec *Sylvietta isabellina*
  It was called the Somali Long-billed Crombec in the 1996 list
- ☐ Green Crombec *Sylvietta virens*
- ☐ White-browed Crombec *Sylvietta leucophrys*

- ☐ Blackcap *Sylvia atricapilla* **PM**
- ☐ Garden Warbler *Sylvia borin* **PM**
- ☐ Barred Warbler *Sylvia nisoria* **PM**
- ☐ Common Whitethroat *Sylvia communis* **PM**
- ☐ Brown Parisoma *Parisoma lugens*
- ☐ Banded Parisoma *Parisoma boehmi*
  Parisoma boehmi marsabit **E**

Timaliidae: illadopses, babblers and chatterers

- ☐ Scaly-breasted Illadopsis *Illadopsis albipectus*
- ☐ Pale-breasted Illadopsis *Illadopsis rufipennis*
- ☐ Brown Illadopsis *Illadopsis fulvescens*
- ☐ Mountain Illadopsis *Illadopsis pyrrhoptera*
- ☐ Grey-chested Babbler *Kakamega poliothorax*
  It was called the Grey-chested Illadopsis in the 1996 list; no documented records since 1979
- ☐ African Hill Babbler *Pseudoalcippe abyssinica*
- ☐ Scaly Chatterer *Turdoides aylmeri*
- ☐ Rufous Chatterer *Turdoides rubiginosa*
  Species epithet was *rubiginosus* in the 1996 list, changed following David & Gosselin (2002b)
- ☐ Black-lored Babbler *Turdoides sharpei*
  Turdoides sharpei vepres **E**
- ☐ Brown Babbler *Turdoides plebejus*
- ☐ Arrow-marked Babbler *Turdoides jardineii*
- ☐ Scaly Babbler *Turdoides squamulata*
  Species epithet was *squamulatus* in the 1996 list, changed following David & Gosselin (2002b)
- ☐ Hinde's Babbler *Turdoides hindei* **E; Vulnerable**
- ☐ Northern Pied Babbler *Turdoides hypoleuca*
  Species epithet was *hypoleucus* in the 1996 list, changed following David & Gosselin (2002b)

Zosteropidae: white-eyes

- ☐ Abyssinian White-eye *Zosterops abyssinicus*
- ☐ African Yellow White-eye *Zosterops senegalensis*
  It was called the Yellow White-eye in the 1996 list
- ☐ Montane White-eye *Zosterops poliogastrus*
  It was called *Z. poliogaster* in the 1996 list, but *Z. poliogastrus* is spelling and gender correct (David & Gosselin 2002a). Includes the Kulal White-eye *Z. p. kulalensis* **E**, the Kikuyu White-eye *Z. p. kikuyuensis* **E**, and the Taita White-eye *Z. p. silvanus* **E**; these three taxa are sometimes regarded as separate species

Certhiidae: creepers

- ☐ Spotted Creeper *Salpornis spilonotus*
  Now very rare in the Kapenguria area of western Kenya

Sturnidae: starlings and oxpeckers

- ☐ Rose-coloured Starling *Pastor roseus* **VP**
  An adult seen and photographed on 17 July 2003 in the Masai Mara is the sole record

- ☐ Wattled Starling *Creatophora cinerea* **am**
- ☐ Greater Blue-eared Starling *Lamprotornis chalybaeus*
- ☐ Lesser Blue-eared Starling *Lamprotornis chloropterus*
- ☐ Bronze-tailed Starling *Lamprotornis chalcurus*
- ☐ Splendid Starling *Lamprotornis splendidus* **am**
- ☐ Purple Starling *Lamprotornis purpureus*
- ☐ Rüppell's Starling *Lamprotornis purpuroptera*
  It was called Rüppell's Long-tailed Starling *L. purpuropterus* in the 1996 list
- ☐ Golden-breasted Starling *Lamprotornis regius*
  It was *Cosmopsarus regius* in the 1996 list. *Lamprotornis regius magnificus* **E**
- ☐ Black-bellied Starling *Lamprotornis corruscus*
- ☐ Superb Starling *Lamprotornis superbus*
- ☐ Hildebrandt's Starling *Lamprotornis hildebrandti*
- ☐ Shelley's Starling *Lamprotornis shelleyi* **am**
- ☐ Fischer's Starling *Lamprotornis fischeri*
  It was *Spreo fischeri* in the 1996 list
- ☐ Abbott's Starling *Cinnyricinclus femoralis* **Vulnerable**
- ☐ Violet-backed Starling *Cinnyricinclus leucogaster* **AM**
- ☐ White-crowned Starling *Spreo albicapillus*
  Spreo albicapillus horrensis **E**
- ☐ Magpie Starling *Speculipastor bicolor*
- ☐ Red-winged Starling *Onychognathus morio*
- ☐ Slender-billed Starling *Onychognathus tenuirostris*
- ☐ Waller's Starling *Onychognathus walleri*
- ☐ Bristle-crowned Starling *Onychognathus salvadorii*
- ☐ Stuhlmann's Starling *Poeoptera stuhlmanni*
- ☐ Kenrick's Starling *Poeoptera kenricki*
  Poeoptera kenricki bensoni **E**
- ☐ Sharpe's Starling *Pholia sharpii*
  It was called *Cinnyricinclus sharpii* in the 1996 list
- ☐ Red-billed Oxpecker *Buphagus erythrorhynchus*
- ☐ Yellow-billed Oxpecker *Buphagus africanus*

Turdidae: thrushes

- ☐ Red-tailed Ant Thrush *Neocossyphus rufus*
- ☐ White-tailed Ant Thrush *Neocossyphus poensis*
- ☐ Abyssinian Ground Thrush *Zoothera piaggiae*
- ☐ Orange Ground Thrush *Zoothera gurneyi*
  Zoothera gurneyi chuka **E**, *Zoothera gurneyi raineyi* **E**
- ☐ Spotted Ground Thrush *Zoothera guttata* **AM; Endangered**
- ☐ African Thrush *Turdus pelios*
- ☐ African Bare-eyed Thrush *Turdus tephronotus*
  It was just the Bare-eyed Thrush in the 1996 list

☐ Olive Thrush *Turdus olivaceus*

☐ Taita Thrush *Turdus helleri* **E; Critically Endangered**
It was listed as the species aggregate *Turdus (olivaceus) helleri* in the 1996 list. However, Bowie *et al.* 2005 and Gill & Wright (2006) accord it full specific status

☐ Brown-chested Alethe *Alethe poliocephala*
Alethe poliocephala akelyae **E**

Muscicapidae: chats, wheatears and Old World flycatchers

The following 38 species were included in the Turdidae, 'thrushes, chats and relatives', in the 1996 list

☐ White-starred Robin *Pogonocichla stellata*
Pogonocichla stellata macarthuri **E**

☐ Equatorial Akalat *Sheppardia aequatorialis*

☐ East Coast Akalat *Sheppardia gunningi* **Near Threatened**

☐ Grey-winged Robin *Sheppardia polioptera*

☐ Thrush Nightingale *Luscinia luscinia* **PM**
It was called by the German name Sprosser in the 1996 list

☐ Common Nightingale *Luscinia megarhynchos* **PM**
It was called simply Nightingale in the 1996 list

☐ Irania *Irania gutturalis* **PM**
Often called the White-throated Robin in Europe but in South Africa this is the English name always used for *Cossypha humeralis*; *C. humeralis* is called the White-throated Robin-Chat by Gill & Wright (2006)

☐ Cape Robin Chat *Cossypha caffra*

☐ Blue-shouldered Robin Chat *Cossypha cyanocampter*

☐ Rüppell's Robin Chat *Cossypha semirufa*

☐ White-browed Robin Chat *Cossypha heuglini*

☐ Red-capped Robin Chat *Cossypha natalensis* **am**

☐ Snowy-headed Robin Chat *Cossypha niveicapilla*

☐ Collared Palm Thrush *Cichladusa arquata*

☐ Spotted Palm Thrush *Cichladusa guttata*
It was called the Spotted Morning Thrush in the 1996 list

☐ Bearded Scrub Robin *Cercotrichas quadrivirgata*
It was called the Eastern Bearded Scrub Robin in the 1996 list

☐ Brown-backed Scrub Robin *Cercotrichas hartlaubi*

☐ White-browed Scrub Robin *Cercotrichas leucophrys*
The races *vulpina*, *leucoptera* and *eluta* are sometimes called the White-winged Scrub Robin

☐ Rufous Bush Chat *Cercotrichas galactotes* **PM**

☐ Common Redstart *Phoenicurus phoenicurus* **PM**

☐ Whinchat *Saxicola rubetra* **PM**

☐ Common Stonechat *Saxicola torquatus*
It was called *Saxicola torquata* in the 1996 list; the *-us* ending was changed to *-a* following David & Gosselin (2002b); but *rubetra* is correct for the Whinchat

☐ Capped Wheatear *Oenanthe pileata*

☐ Heuglin's Wheatear *Oenanthe heuglini*
It was *Oenanthe (bottae) heuglini* in the 1996 list

- ☐ Isabelline Wheatear *Oenanthe isabellina* **PM**
- ☐ Northern Wheatear *Oenanthe oenanthe* **PM**
- ☐ Pied Wheatear *Oenanthe pleschanka* **PM**
- ☐ Black-eared Wheatear *Oenanthe hispanica* **PM**
  Two records: one caught, ringed and photographed, 23 March–5 April 1984, Athi River, near Nairobi; and one seen, 2 December 1994, Lake Baringo
- ☐ Desert Wheatear *Oenanthe deserti* **PM**
  Two seen, 17 February 1984, Kiunga (Backhurst 1985); one, 18 October 1996, Kerio Valley (EARC 1998)
- ☐ Abyssinian Wheatear *Oenanthe lugubris*
  It was called the Abyssinian Black Wheatear in the 1996 list; Schalow's Wheatear is an alternative name for Kenyan birds of the race *schalowi*
- ☐ Familiar Chat *Cercomela familiaris*
  It was called the Red-tailed Chat in the 1996 list
- ☐ Brown-tailed Rock Chat *Cercomela scotocerca*
- ☐ Moorland Chat *Cercomela sordida*
  It was called the Alpine Chat in the 1996 list
- ☐ Northern Anteater Chat *Myrmecocichla aethiops*
- ☐ Sooty Chat *Myrmecocichla nigra*
- ☐ Mocking Cliff Chat *Thamnolaea cinnamomeiventris*
  It was called the White-shouldered Cliff Chat in the 1996 list
- ☐ Common Rock Thrush *Monticola saxatilis* **PM**
- ☐ Little Rock Thrush *Monticola rufocinereus*

The following six genera were included in the (restricted) Muscicapidae, Old World flycatchers, in the 1996 list

- ☐ White-eyed Slaty Flycatcher *Melaenornis fischeri*
- ☐ Northern Black Flycatcher *Melaenornis edolioides*
- ☐ Southern Black Flycatcher *Melaenornis pammelaina*
- ☐ Pale Flycatcher *Bradornis pallidus*
- ☐ African Grey Flycatcher *Bradornis microrhynchus*
  Bradornis microrhynchus taruensis **E**
- ☐ Silverbird *Empidornis semipartitus*
- ☐ Spotted Flycatcher *Muscicapa striata* **PM**
- ☐ Gambaga Flycatcher *Muscicapa gambagae* **am**
- ☐ Ashy Flycatcher *Muscicapa caerulescens*
- ☐ Swamp Flycatcher *Muscicapa aquatica*
- ☐ Chapin's Flycatcher *Muscicapa lendu* **Vulnerable**
- ☐ African Dusky Flycatcher *Muscicapa adusta*
  Muscicapa adusta marsabit **E**
- ☐ Lead-coloured Flycatcher *Myioparus plumbeus*
- ☐ Pied Flycatcher *Ficedula hypoleuca* **VP**
  Not included in the 1996 list; the sole record is of a first-winter male (collected), 8 December 1965, at Kakamega Forest (Pearson 1998)
- ☐ Collared Flycatcher *Ficedula albicollis* **VP**
  One collected on 2 October 1972 at Ng'iya, western Kenya (Britton 1980)
- ☐ Semi-collared Flycatcher *Ficedula semitorquata* **PM; Near Threatened**

Nectariniidae: sunbirds

We follow Irwin (1999) in recognizing seven genera of sunbirds, whereas the 1996 list used only two; the first eight species below were included in *Anthreptes* in 1996

- ☐ Plain-backed Sunbird *Anthreptes reichenowi* **Near Threatened**
- ☐ Western Violet-backed Sunbird *Anthreptes longuemarei*
- ☐ Eastern Violet-backed Sunbird *Anthreptes orientalis*
- ☐ Uluguru Violet-backed Sunbird *Anthreptes neglectus*
- ☐ Grey-chinned Sunbird *Anthreptes rectirostris*
  It was called the Green Sunbird in the 1996 list
- ☐ Collared Sunbird *Hedydipna collaris* — *Anthodiaeta* (Mann & Cheke 2006)
- ☐ Pygmy Sunbird *Hedydipna platyura* **am**
  Only three documented records in the last fifty years: November 1976, Baringo; November 1979, Kapedo; October 1990, Kerio Valley
- ☐ Amani Sunbird *Hedydipna pallidigaster* **Endangered**

The following 27 species were included in *Nectarinia* in the 1996 list; some species epithet changes from feminine to masculine result from this

- ☐ Green-headed Sunbird *Cyanomitra verticalis*
- ☐ Olive Sunbird *Cyanomitra olivacea*
- ☐ Mouse-coloured Sunbird *Cyanomitra veroxii*
- ☐ Green-throated Sunbird *Chalcomitra rubescens*
- ☐ Amethyst Sunbird *Chalcomitra amethystina*
- ☐ Scarlet-chested Sunbird *Chalcomitra senegalensis*
- ☐ Hunter's Sunbird *Chalcomitra hunteri*
- ☐ Tacazze Sunbird *Nectarinia tacazze*
- ☐ Bronze Sunbird *Nectarinia kilimensis*
- ☐ Malachite Sunbird *Nectarinia famosa*
- ☐ Scarlet-tufted Sunbird *Nectarinia johnstoni*
  It was called the Scarlet-tufted Malachite Sunbird in the 1996 list
- ☐ Golden-winged Sunbird *Drepanorhynchus reichenowi*
  Drepanorhynchus reichenowi lathburyi **E**
- ☐ Olive-bellied Sunbird *Cinnyris chloropygius*
- ☐ Northern Double-collared Sunbird *Cinnyris reichenowi*
  It was called *Nectarinia preussi* in the 1996 list, but with its inclusion in *Cinnyris* here, the ICZN (1999) requires the species epithet *reichenowi* to be used
- ☐ Eastern Double-collared Sunbird *Cinnyris mediocris*
- ☐ Usambara Double-collared Sunbird *Cinnyris usambaricus*
  This species is added to the Kenya list following Bowie *et al.* (2004)
- ☐ Beautiful Sunbird *Cinnyris pulchellus*
- ☐ Marico Sunbird *Cinnyris mariquensis*
- ☐ Red-chested Sunbird *Cinnyris erythrocercus*
- ☐ Black-bellied Sunbird *Cinnyris nectarinioides*
- ☐ Purple-banded Sunbird *Cinnyris bifasciatus*
- ☐ Tsavo Sunbird *Cinnyris tsavoensis*
  It was included (as the race *tsavoensis*) with the Purple-banded Sunbird in the 1996 list

- ☐ Violet-breasted Sunbird *Cinnyris chalcomelas*
- ☐ Orange-tufted Sunbird *Cinnyris bouvieri*
- ☐ Shining Sunbird *Cinnyris habessinicus*
- ☐ Variable Sunbird *Cinnyris venustus*
- ☐ Superb Sunbird *Cinnyris superbus*
- ☐ Copper Sunbird *Cinnyris cupreus*

Passeridae: sparrow weavers, Old World sparrows and petronias

The first four species below were included in the Ploceidae in the 1996 list

- ☐ White-browed Sparrow Weaver *Plocepasser mahali*
  The name Sparrow-Weaver was used in the 1996 list for this and the next two species
- ☐ Chestnut-crowned Sparrow Weaver *Plocepasser superciliosus*
- ☐ Donaldson Smith's Sparrow Weaver *Plocepasser donaldsoni*
  The American zoologist and collector Dr Arthur Donaldson Smith (1864–1939) had no hyphen in his name
- ☐ Rufous-tailed Weaver *Histurgops ruficauda*
  While not included in the 1996 list, this hitherto Tanzanian endemic has bred in small numbers since July 2000 in the Masai Mara

The next two species were included in the Ploceidae in the 1996 list

- ☐ Grey-capped Social Weaver *Pseudonigrita arnaudi*
- ☐ Black-capped Social Weaver *Pseudonigrita cabanisi*
- ☐ House Sparrow *Passer domesticus*
- ☐ Somali Sparrow *Passer castanopterus*
- ☐ Kenya Rufous Sparrow *Passer rufocinctus*
  It was called simply the Rufous Sparrow in the 1996 list and the Kenya Sparrow by Gill & Wright (2006). The race *shelleyi* (Shelley's Rufous Sparrow) is sometimes regarded as a separate species
- ☐ Grey-headed Sparrow *Passer griseus*
  The races *gongonensis*, *suahelicus* and *swainsonii* are frequently regarded as separate species with the English names Parrot-billed Sparrow, Swahili Sparrow and Swainson's Sparrow, respectively (see, for instance Fry & Keith (Eds) 2004, Gill & Wright 2006)
- ☐ Chestnut Sparrow *Passer eminibey*
- ☐ Yellow-spotted Petronia *Petronia pyrgita*

Ploceidae: weavers, bishops and widowbirds

The first three species were called Buffalo-Weavers in the 1996 list

- ☐ White-billed Buffalo Weaver *Bubalornis albirostris*
- ☐ Red-billed Buffalo Weaver *Bubalornis niger*
- ☐ White-headed Buffalo Weaver *Dinemellia dinemelli*
- ☐ Speckle-fronted Weaver *Sporopipes frontalis*
- ☐ Grosbeak Weaver *Amblyospiza albifrons*
- ☐ Baglafecht Weaver *Ploceus baglafecht*
  The races *emini*, *reichenowi* and *stuhlmanni* have often borne the English names Emin's Weaver, Reichenow's Weaver and Stuhlmann's Weaver, respectively
- ☐ Slender-billed Weaver *Ploceus pelzelni*

- ☐ Little Weaver *Ploceus luteolus*
- ☐ Spectacled Weaver *Ploceus ocularis*
- ☐ Black-necked Weaver *Ploceus nigricollis*
- ☐ Black-billed Weaver *Ploceus melanogaster*
- ☐ Eastern Golden Weaver *Ploceus subaureus*
  It was called the African Golden Weaver in the 1996 list
- ☐ Holub's Golden Weaver *Ploceus xanthops*
- ☐ Orange Weaver *Ploceus aurantius*
  No documented records since December 1984
- ☐ Heuglin's Masked Weaver *Ploceus heuglini*
- ☐ Golden Palm Weaver *Ploceus bojeri*
- ☐ Taveta Golden Weaver *Ploceus castaneiceps*
- ☐ Northern Brown-throated Weaver *Ploceus castanops*
- ☐ Northern Masked Weaver *Ploceus taeniopterus*
  The first record was of one collected in January 1959 at Ileret, near the northeastern shore of Lake Turkana but since the 1970s it has become well established at Lake Baringo (Zimmerman *et al.* 1996)
- ☐ Lesser Masked Weaver *Ploceus intermedius*
- ☐ Vitelline Masked Weaver *Ploceus vitellinus*
  It was included in *Ploceus velatus* in the 1996 list
- ☐ Speke's Weaver *Ploceus spekei*
- ☐ Village Weaver *Ploceus cucullatus*
  It was called the Black-headed Weaver in the 1996 list
- ☐ Vieillot's Black Weaver *Ploceus nigerrimus*
- ☐ Clarke's Weaver *Ploceus golandi* **E; Endangered**
- ☐ Juba Weaver *Ploceus dichrocephalus*
- ☐ Yellow-backed Weaver *Ploceus melanocephalus*
  Gill & Wright (2006) rather confusingly call this species the Black-headed Weaver (a direct translation of the specific epithet) but this is the name quite often used for *Ploceus cucullatus*, here called the Village Weaver
- ☐ Golden-backed Weaver *Ploceus jacksoni*
  It was called Jackson's Golden-backed Weaver in the 1996 list
- ☐ Chestnut Weaver *Ploceus rubiginosus* **am**
- ☐ Yellow-mantled Weaver *Ploceus tricolor* **Ex**
  Last recorded in July 1972 in Kakamega Forest
- ☐ Compact Weaver *Ploceus superciliosus*
- ☐ Dark-backed Weaver *Ploceus bicolor*
- ☐ Brown-capped Weaver *Ploceus insignis*
- ☐ Red-headed Malimbe *Malimbus rubricollis*
- ☐ Red-headed Weaver *Anaplectes melanotis*
  It was called *Anaplectes rubriceps* in the 1996 list — *rubriceps* now correct, Oschadleus 2009
- ☐ Cardinal Quelea *Quelea cardinalis*
- ☐ Red-headed Quelea *Quelea erythrops* **am**
  Few documented records in the last fifteen years
- ☐ Red-billed Quelea *Quelea quelea* **am**
- ☐ Yellow-crowned Bishop *Euplectes afer*

- ☐ Fire-fronted Bishop *Euplectes diadematus*
- ☐ Black Bishop *Euplectes gierowii*
- ☐ Zanzibar Red Bishop *Euplectes nigroventris*
- ☐ Black-winged Red Bishop *Euplectes hordeaceus*
- ☐ Southern Red Bishop *Euplectes orix*
- ☐ Northern Red Bishop *Euplectes franciscanus*
- ☐ Yellow Bishop *Euplectes capensis*
- ☐ Fan-tailed Widowbird *Euplectes axillaris*
- ☐ Yellow-mantled Widowbird *Euplectes macroura*
  The race *macrocerca* is sometimes given the English name the Yellow-shouldered Widowbird
- ☐ Marsh Widowbird *Euplectes hartlaubi*
  It was called Hartlaub's Marsh Widowbird in the 1996 list
- ☐ White-winged Widowbird *Euplectes albonotatus*
- ☐ Red-collared Widowbird *Euplectes ardens*
- ☐ Long-tailed Widowbird *Euplectes progne*
  Euplectes progne delamerei **E**
- ☐ Jackson's Widowbird *Euplectes jacksoni* **Near Threatened**

Estrildidae: waxbills

- ☐ White-breasted Negrofinch *Nigrita fusconotus*
  This and the next species take the masculine *-us* ending following David & Gosselin (2002b)
- ☐ Grey-headed Negrofinch *Nigrita canicapillus*
- ☐ Yellow-bellied Waxbill *Coccopygia quartinia*
  It was included in *Estrilda* in the 1996 list
- ☐ Green-backed Twinspot *Mandingoa nitidula*
- ☐ Abyssinian Crimsonwing *Cryptospiza salvadorii*
- ☐ Fawn-breasted Waxbill *Estrilda paludicola*
- ☐ Crimson-rumped Waxbill *Estrilda rhodopyga*
- ☐ Black-rumped Waxbill *Estrilda troglodytes*
- ☐ Common Waxbill *Estrilda astrild*
- ☐ Black-crowned Waxbill *Estrilda nonnula*
- ☐ Kandt's Waxbill *Estrilda kandti*
  Listed as the Black-headed Waxbill *Estrilda atricapilla* in the 1996 list (but see Fry & Keith (Eds) 2004)
- ☐ Black-faced Waxbill *Estrilda erythronotus*
- ☐ Black-cheeked Waxbill *Estrilda charmosyna*
- ☐ Red-headed Bluebill *Spermophaga ruficapilla*
- ☐ Black-bellied Seed-cracker *Pyrenestes ostrinus*
- ☐ Red-cheeked Cordon-bleu *Uraeginthus bengalus*
  Uraeginthus bengalus brunneigularis **E**
- ☐ Blue-capped Cordon-bleu *Uraeginthus cyanocephalus*
- ☐ Purple Grenadier *Granatina ianthinogaster*
  It was included in *Uraeginthus* in the 1996 list
- ☐ Peters's Twinspot *Hypargos niveoguttatus*

- ☐ Brown Twinspot *Clytospiza monteiri*
- ☐ Orange-winged Pytilia *Pytilia afra*
- ☐ Green-winged Pytilia *Pytilia melba*
- ☐ Black-bellied Firefinch *Lagonosticta rara*
- ☐ Bar-breasted Firefinch *Lagonosticta rufopicta*
- ☐ Red-billed Firefinch *Lagonosticta senegala*
- ☐ African Firefinch *Lagonosticta rubricata*
- ☐ Jameson's Firefinch *Lagonosticta rhodopareia*
- ☐ Cut-throat Finch *Amadina fasciata*
- ☐ African Quailfinch *Ortygospiza fuscocrissa*
  The English name was the Quail-Finch in the 1996 list and its scientific name was *O. atricollis*
- ☐ Orange-breasted Waxbill *Amandava subflava*
  It was called the Zebra Waxbill in the 1996 list
- ☐ Locust Finch *Paludipasser locustella*
  It was included in *Ortygospiza* in the 1996 list
- ☐ Grey-headed Silverbill *Odontospiza griseicapilla*
  This and the next four species were included in *Lonchura* in the 1996 list
- ☐ Bronze Mannikin *Spermestes cucullatus*
- ☐ Black-and-white Mannikin *Spermestes bicolor*
  The race *nigriceps*, the Rufous-backed Mannikin, has often been regarded as a separate species in the past
- ☐ Magpie Mannikin *Spermestes fringilloides*
- ☐ African Silverbill *Euodice cantans*

Viduidae: Parasitic Weaver, indigobirds and whydahs

This family was treated as a subfamily, Viduinae, within the Estrildidae in the 1996 list

- ☐ Parasitic Weaver *Anomalospiza imberbis*
  It was included in the Ploceidae in the 1996 list
- ☐ Pin-tailed Whydah *Vidua macroura*
- ☐ Broad-tailed Paradise Whydah *Vidua obtusa*
  The last documented record was of a male collected at Chuka in February 1947
- ☐ Eastern Paradise Whydah *Vidua paradisaea*
  It was called simply the Paradise Whydah in the 1996 list
- ☐ Steel-blue Whydah *Vidua hypocherina*
- ☐ Straw-tailed Whydah *Vidua fischeri*
- ☐ Village Indigobird *Vidua chalybeata*
- ☐ Dusky Indigobird *Vidua funerea*
  Tape recorded and collected in the early 1990s in the Kerio Valley alongside the other two Kenya indigobird species (Payne *et al.* 1992). It was called the Variable Indigobird in the 1996 list
- ☐ Purple Indigobird *Vidua purpurascens*

Motacillidae: wagtails, longclaws and pipits

- ☐ Yellow Wagtail *Motacilla flava* **PM**
  It is called the Western Yellow Wagtail by Gill & Wright (2006) who split off the Alaskan and northeastern Siberian races as the species *M. tschutschensis*, the Eastern Yellow Wagtail

- ☐ Cape Wagtail *Motacilla capensis*
- ☐ Grey Wagtail *Motacilla cinerea* **PM**
- ☐ Mountain Wagtail *Motacilla clara*
- ☐ White Wagtail *Motacilla alba* **PM**
- ☐ African Pied Wagtail *Motacilla aguimp*
- ☐ Golden Pipit *Tmetothylacus tenellus* **AM**
- ☐ Sharpe's Longclaw *Macronyx sharpei* **E; Endangered**
- ☐ Yellow-throated Longclaw *Macronyx croceus*
- ☐ Pangani Longclaw *Macronyx aurantiigula*
- ☐ Rosy-breasted Longclaw *Macronyx ameliae*
- ☐ Grassland Pipit *Anthus cinnamomeus*
  The taxon *latistriatus*, described from Mumias in 1899, and not recorded in Kenya since, is sometimes considered a race of *A. cinnamomeus*, and sometimes (e.g., by Clancey 1990) as a full species, *A. latistriatus*, Jackson's Pipit
- ☐ Tawny Pipit *Anthus campestris* **PM**
- ☐ Long-billed Pipit *Anthus similis*
- ☐ Plain-backed Pipit *Anthus leucophrys*
- ☐ Buffy Pipit *Anthus vaalensis*
  The Kenyan race *goodsoni* was formerly treated within *Anthus leucophrys* in the 1996 list
- ☐ Tree Pipit *Anthus trivialis* **PM**
- ☐ Red-throated Pipit *Anthus cervinus* **PM**
- ☐ Striped Pipit *Anthus lineiventris*
- ☐ Bush Pipit *Anthus caffer*
- ☐ Sokoke Pipit *Anthus sokokensis* **Endangered**
- ☐ Malindi Pipit *Anthus melindae* **Near Threatened**

Fringillidae: canaries, citrils, seedeaters and relatives

The current arrangement of genera and species within this family follows Gill & Wright (2006), itself largely based on that of van den Elzen (1985)

- ☐ Yellow-crowned Canary *Serinus flavivertex*
  It was included within *Serinus canicollis* in the 1996 list

The following fourteen *Crithagra* species were assigned to *Serinus* in the 1996 list where four species (*mozambica*, *dorsostriata*, *sulphurata* and *striolata*) had the masculine ending *-us*

- ☐ African Citril *Crithagra citrinelloides*
  Crithagra citrinelloides kikuyuensis **E**
- ☐ Southern Citril *Crithagra hyposticta*
  It was included in the African Citril in the 1996 list
- ☐ Papyrus Canary *Crithagra koliensis*
- ☐ Reichenow's Seedeater *Crithagra reichenowi*
  It was called the Yellow-rumped Seedeater in the 1996 list; however, that English name has now been given to the Ethiopian and Eritrean species *C. xanthopygia* by Gill & Wright (2006)
- ☐ Black-throated Seedeater *Crithagra atrogularis*
  The most recent documented record was of a pair seen on 26 September 1991 near Sioport, western Kenya; this was the first record for over fifty years
- ☐ Yellow-fronted Canary *Crithagra mozambica*

- ☐ Northern Grosbeak Canary *Crithagra donaldsoni*
- ☐ Southern Grosbeak Canary *Crithagra buchanani*
- ☐ White-bellied Canary *Crithagra dorsostriata*
- ☐ Brimstone Canary *Crithagra sulphurata*
- ☐ Stripe-breasted Seedeater *Crithagra reichardi*
  It was called *Serinus (reichardi) striatipectus* in the 1996 list
- ☐ Streaky-headed Seedeater *Crithagra gularis*
  Only known from three records: the type, 14 June 1900, southern slopes of Mt Elgon; July 1973, Suam River; September 1993, south of Kapenguria. It was called *Serinus (gularis) elgonensis* in the 1996 list
- ☐ Streaky Seedeater *Crithagra striolata*
- ☐ Thick-billed Seedeater *Crithagra burtoni*
  Crithagra burtoni albifrons **E**
- ☐ Oriole Finch *Linurgus olivaceus*
  It was called the Oriole-Finch in the 1996 list

Emberizidae: Old World buntings

- ☐ Ortolan Bunting *Emberiza hortulana* **PM**
- ☐ House Bunting *Emberiza striolata*
- ☐ Cinnamon-breasted Bunting *Emberiza tahapisi*
  It was called the Cinnamon-breasted Rock Bunting in the 1996 list
- ☐ Somali Bunting *Emberiza poliopleura*
  It was called the Somali Golden-breasted Bunting in the 1996 list
- ☐ Golden-breasted Bunting *Emberiza flaviventris*
- ☐ Brown-rumped Bunting *Emberiza affinis*
  No documented records since 1988

About this list

With 1100 species, a rich variety of habitats, and relative ease of travel and access to its many National Parks and reserves, Kenya is one of the main countries in Africa for bird watching. For its many visitors and for increasing numbers of local enthusiasts, an updated version of the country's bird list has become essential.

More than a decade has passed since the Bird Committee (or Ornithological Sub-committee as it was then called) last revised this list. In the meantime, avian systematics and taxonomy have moved on rapidly, and species and genus changes are constantly being suggested from increasing numbers of molecular studies. Some surprising results are emerging from the DNA sequencing of more and more groups of birds. There will never be a perfect time for a review of the list, and the present offering is an attempt to keep up with the more generally accepted changes, bearing in mind that further revision will be needed in a few years' time.

The starting point for species and genera was our own 1996 list. This was reviewed in the light of decisions made in the later volumes of *Birds of Africa* (e.g., Fry & Keith (Eds) 2004) and those of the third edition of 'Howard & Moore' (Dickinson 2003). We have adopted a number of more recently suggested amendments, but have tended to take a cautious approach to new species-splits and generic arrangements. We have avoided changes where questions clearly remain or where we feel that wider acceptance is still needed. Our judgement of species limits has continued to be based on the biological species concept. The current total of 1100 accepted species may well be fewer than in some lists used by today's birders.

The order of families differs from the traditional one used for the previous list. It follows the 'gamebirds-and-ducks-first' sequence of Howard & Moore, which was informed by up-to-date genetic studies, and is now being used increasingly. But even this order is bound to be modified as a result of the recent DNA work by Hackett *et al.* (2008).

Our approach to English names has been conservative. We have been reluctant to move away from well-established names, and have done so only in favour of an alternative which we regard as being clearly better or more appropriate, or where a species-split has made a new name necessary. In an attempt to standardize, the International Ornithological Congress has published a recommended list of world English bird names (Gill & Wright 2006). We have changed some of our names to conform with suggested ones in this list, but in other cases we have preferred to retain our own name; in some, but not all, cases, this has been noted. We have generally used as few hyphens in group names as possible.

Omitted species

The following species are not included in the checklist for the reasons given.

Mascarene Shearwater *Puffinus atrodorsalis*

It was included in the 1996 list in square brackets, i.e., there was insufficient evidence. It is now considered a race of Audubon's Shearwater (Bretagnolle *et al.* 2000).

Red-tailed Tropicbird *Phaethon rubricauda*

It was included in the 1996 list in square brackets. Although there have been unsubstantiated sightings from Kenyan waters, there is no confirmed record to date.

Eurasian Griffon *Gyps fulvus*

Although details of a January 2000 sighting from Amboseli National Park have appeared in the literature (*Bulletin of the African Bird Club* 8: 59), the details have not yet been submitted to the EARC. It was called the Griffon Vulture by Gill & Wright (2006).

Little Gull *Hydrocoloeus minutus*

The record of *c.* 60 at Lake Turkana in January 1979 (Groh 1979) has been rejected (see Zimmerman *et al.* (1996) — where it is called *Larus minutus* — for more detail).

Mediterranean Gull *Larus melanocephalus*

Lake Turkana and Lamu sightings are considered dubious, see Zimmerman *et al.* (1996).

White-eyed Gull *Larus leucophthalmus*

It was included in the 1996 list in square brackets; the claimed record from Lake Turkana remains unsubstantiated; see Zimmerman *et al.* (1996) for more detail.

Arctic Tern *Sterna paradisaea*

Although details and photographs of one from the Sabaki River Mouth (6 July 2002) have appeared in the literature (Bisschop 2002 and [Bisschop 2003]) this observation has not been submitted to the EARC.

Catharacta skuas

There have been several coastal sightings of large brown skuas belonging to this genus, but none has been substantiated (Zimmerman *et al.* 1996). Gill & Wright (2006) do not admit this genus.

Pallid Swift *Apus pallidus*

The EARC's decision on a claimed record has been put on hold pending "expert opinion".

Boubous *Laniarius* spp.

The Tropical Boubou *Laniarius aethiopicus* has recently been split by Nguembock *et al.* (2008) into several taxa, including what they treat as three full species which occur in Kenya. It is possible that these three species will find acceptance by the authors of the next edition of Dickinson (2003) due to be published in 2010. The most widespread of these new taxa is the Tropical Boubou *Laniarius major*; in the far north of the country, around Moyale, the Ethiopian Boubou *L. aethiopicaus* occurs, while the coastal lowlands are occupied by the Dimorphic Boubou *L. sublacteus*. In the present list we remain conservative, retaining *Laniarius aethiopicus*, the Tropical Boubou.

Melodious Warbler *Hippolais polyglotta*

A bird identified as this species was caught, photographed and ringed at Ngulia, Tsavo on 19 November 1995 (Pearson *et al.* 1998). It was determined on plumage and biometric characters and, in addition, an analysis of mitrochondrial DNA carried out in the laboratory of Dr Andreas Helbig confirmed its maternal identity. After the death of Dr Helbig, information has come to hand that the information sent to David Pearson was incorrect (Gerhard Nikolaus *in litt.*). The species was not included in the 1996 list.

Forest Robin *Stiphrornis erythrothorax*

It was included in the 1996 list in square brackets but the specimen labelled "Kipkabus, nr Eldoret 16 April 1966" collected by Brian Heath, is considered to have been labelled erroneously. The specimen is suspected to have originated in western Uganda.

Rüppell's Weaver *Ploceus galbula*

Inclusion in the 1996 list was based on a 1901 female specimen from the Kenya–Ethiopian border area (Ash 1988). However, John Miskell (pers. comm.) considers the record dubious as he believes Erlanger (1907) to have been confused over the females of Rüppell's and Juba Weavers, and further, that his specimens from Southern Somalia and the Kenya–Ethiopian border may have been mis-identified.

'Nairobi Pipit'

From 1994, Brian Finch has noted pipits in Nairobi National Park which he, and others, believe to be an undescribed species, the 'Nairobi Pipit'. The bird is frequently mentioned in websites and an article on it has been published (Finch 2002). However, this bird has not been scientifically described or named, nor has it been submitted to the EARC.

References

Ash, J.S. 1988. Rüppell's Weaver *Ploceus galbula* new to Kenya. *Scopus* 11: 94–95.

Backhurst, G.C. 1985. Desert Wheatear *Oenanthe deserti* in Kenya. *Scopus* 9: 140–141.

Backhurst, G.C. & Pearson, D.J. 1976. Savi's Warbler in Tsavo, an addition to the Kenya list. *EANHS Bulletin* 1976: 21–22.

Bishop, R. & Bishop, A. 2001. Spotted Sandpiper *Actitis macularia* in Kenya: the first record in sub-Saharan Africa. *Bulletin of the African Bird Club* 8: 48–49.

Bisschop, J. 2002. Arctic Tern in Kenya in July 2002. *Dutch Birding* 24: 358–359.

[Bisschop, J. 2003. Arctic Tern in Kenya in July 2002. *Kenya Birds* 10: 19. *A shortened version of the original paper in Dutch Birding.*]

Borghesio, L., Ndang'ang'a, K., Nalyanya, N., Kahindi, M. & Laiolo, P. 2004. Some interesting bird observations from northern Kenya. *Scopus* 24: 33–39.

Bowie, R.C.K., Fjeldså, J., Hackett, S.J. & Crowe, T.M. 2004. Systematics and biogeography of Double-collared Sunbirds from the Eastern Arc Mountains, Tanzania. *Auk* 121: 660–681.

Bowie, R.C.K., Voelker, G., Fjeldså, J., Lens., L., Hackett, S.J. & Crowe, T.M. 2005. Systematics of the Olive Thrush *Turdus olivaceus* species complex with reference to the taxonomic status of the endangered Taita Thrush *T. helleri. Journal of Avian Biology* 36: 391–404.

Bretagnolle, V., Attié, C. & Mougeot, F. 2000. Audubon's Shearwaters *Puffinus lherminieri* on Réunion Island, Indian Ocean: behaviour, census, distribution, biometrics and breeding biology. *Ibis* 142: 399–412.

Britton, P.L. (Ed.) 1980. *Birds of East Africa.* Nairobi: EANHS.

Cabanis, J. 1868. [Neue ostafrikanische Vögel des Baron von der Decken]. *Journal für Ornithologie* 16: 411–415.

Clancey, P.A. 1990. A review of the indigenous pipits of the Afrotropics. *Durban Museum Novitates.* 15: 42–72.

Clark, W.S. & Edelstam, C. 2001. First record of Cassin's Hawk-Eagle [*sic*] *Spizaetus africanus* for Kenya. *Bulletin of the African Bird Club* 8: 138–139.

Clark, W.S. 2001. First record of European Griffon *Gyps fulvus* for Kenya. *Bulletin of the African Bird Club* 8: 59–60.

David, N. & Gosselin, M. 2002a. Gender agreement of avian species names. *Bulletin of the British Ornithologists' Club* 122: 14–49.

David, N. & Gosselin, M. 2002b. The grammatical gender of avian genera. *Bulletin of the British Ornithologists' Club* 122: 257–282.

Dickinson, E.C. (ed.) 2003. *The Howard & Moore Complete Checklist of the Birds of the World. Revised & enlarged 3rd edition.* London: A. & C. Black/Croom Helm.

EARC [East African Rarities Committee] 2004. East African Rarities Committee [*title*]. *Scopus* 24: 53.

EARC [East African Rarities Committee] 2005. East African Rarities Committee report. *Scopus* 25: 69.

EARC [East African Rarities Committee] 2007. East African Bird Rarities Committee report. *Scopus* 26: 50.

Elzen, R. van den 1985. Systematics and evolution of African canaries and seedeaters (Aves: Carduelidae). *Proceedings of the International Symposium on African Vertebrates, Bonn 1984*. pp. 435–451.

Erlanger, C.F. 1907. Beiträge zur Vogelfauna Nordostafrikas. *Journal für Ornithologie* 1907: 1–158.

Fayad, V.C. & Fayad, C.C. 1977. A Grasshopper Warbler *Locustella naevia* from Kenya. *Scopus* 1: 84–86.

Finch, B. 2002. A 'Nairobi Pipit'. *Swara* 25 (3): 18–20.

Finch, B.W. 2005. River Prinia *Prinia fluviatilis* near Lokichokio: a new species for Kenya and East Africa. *Scopus* 25: 55–60.

Freitag, S. & Robinson, T.J. 1993. Phylogeographic patterns in mitochondrial DNA of the ostrich (*Struthio camelus*). *Auk* 110: 614–622.

Fry, C.H. & Keith, S. (Eds) 2004. *The Birds of Africa* Vol VII. London: Christopher Helm.

Gill, F. & Wright, M. 2006. *Birds of the World: Recommended English Names.* Princeton NJ: Princeton University Press.

Groh, G. 1979. Little Gull *Larus minutus* at Lake Turkana: a new bird for Kenya. *Scopus* 3: 26.

Hackett, S.J., Kimball, R.T., Reddy, S., Bowie, R.C.K., Braun, E.L., Braun, M.J., Chojnowski, J.L., Cox, W.A., Han, K.-L., Harshman, J., Huddleston, C.J., Marks, B.D., Miglia, K.J., Moore, W.S., Sheldon, F.H., Steadman, D.W., Witt, C.C., Yuri, T. 2008. Phylogenomic study of birds reveals their evolutionary history. *Science* 320: 1763–1768.

Harrison, P. 1983. *Seabirds an identification guide.* Beckenham: Croom Helm.

International Commission on Zoological Nomenclature 1999. *International code of zoological nomenclature.* 4th edition. London: ICZN.

Irwin, M.P.S. 1999. The genus *Nectarinia* and the evolution and diversification of sunbirds: an Afrotropical perspective. *Honeyguide* 45: 45–58.

Jackson, C. 1997. First record of Northern Lapwing *Vanellus vanellus* in East Africa. *Scopus* 19: 113–114.

Jackson, C. & Valle, S. 2007. Sooty Shearwater *Puffinus griseus*: a new species for eastern Africa. *Scopus* 26: 31–33.

Oatley, T.B., Oschadleus, H.D., Navarro, R.A. & Underhill, L.G. 1998. *Review of ring recoveries of birds of prey in southern Africa: 1948–1998*. Johannesburg: Endangered Wildlife Trust.

Ornithological Sub-committee, EANHS. 1981. East African Bird Report 1980. *Scopus* 4: 120.

Ornithological Sub-committee, EANHS. 1996. *Check-list of the birds of Kenya.* 3rd edition. Nairobi: OS-c, EANHS.

Nguembok, B., Fjeldså, J., Couloux, A. & Pasquet, E. 2008. Phylogeny of *Laniarius*: Molecular data reveal *L. liberatus* synonomous with *L. erlangeri* and "plumage coloration" as unreliable morphological characters for defining species and species groups. *Molecular Phylogenics and Evolution* 48: 396–407.

Parsons, B.T. 1969. A record of *Oceanodroma leucorhoa* from Kenya. *Bulletin of the British Ornithologists' Club* 89: 120–121.

Payne, R.B., Payne, L.L., Nhlane, M.E.D. & Hustler, K. 1992. Species status and distribution of the parasitic indigo-birds *Vidua* in east and southern Africa. In *Proceedings 8th Pan-African Ornithological Congress* (Ed: R.T. Wilson). Umberleigh: Bartridge Partners.

Pearson, D.J. 1998. Pied Flycatcher *Ficedula hypoleuca* at Kakamega Forest — a readmission to the East African list. *Scopus* 20: 46–48.

Pearson, D.J., Coverdale, M.A.C., Archer, A.L. & Forbes-Watson, A.D. 1989. Great Grey Shrike *Lanius excubitor* in the Ilemi Triangle. *Scopus* 13: 134.

Pearson, D.J., Finch, B.W. & Backhurst, D.E.G. 1988. A second Savi's Warbler *Locustella luscinioides* at Ngulia. *Scopus* 11: 94.

Pearson, D.J., Jackson, C.H. & Backhurst, G.C. 1998. Melodious Warbler *Hippolais polyglotta* at Ngulia, Kenya—first record for East Africa. *Scopus* 20: 43–45.

Sangster, G., Collinson, J.M., Knox, A.G., Parkin, D.T. & Svensson, L. 2007. Taxonomic recommendations for British birds: Fourth report. *Ibis* 149: 853–857.

Sheldon, F.H. & Winkler, D.W. 1993. Intergeneric phylogenetic relationships of swallows estimated by DNA–DNA hybridization. *Auk* 110: 798–824.

White, C.M.N. 1965. *A revised check list of African non-passerine birds.* Lusaka: Government Printer.

Zimmerman, D.A., Turner, D.A. & Pearson, D.J. 1996. *Birds of Kenya and northern Tanzania.* Halfway House: Russel Friedman.

Acknowledgements

The Bird Committee thanks David Pearson and Don Turner for the initial drafting of the list; David Pearson for 'About this list'; Graeme Backhurst for editing, layout and design. Fleur Ng'weno kept everyone connected.

Several other people gave valuable advice and input: Peter Usher for making a special trip to take the cover photo; Darcy Ogada for her meticulous proofreading; Mike Davidson for securing valuable sponsorship; Bob Dowsett for taxonomic advice and help with references; John Miskell and Nigel Redman for advice on names; Gerhard Nikolaus for investigating the 'Melodious Warbler' record; Amedeo Buonajuti for work on the map. Gordon Boy, Brian Finch, David Fisher, Marilda Gascoyne, Colin Jackson, Anthony Kuria, Simon Musila, Oliver Nasirwa, Catherine Ngarachu, Alfred Owino, Shailesh Patel, Itai Shanni, staff of the Ornithology Section of the National Museums of Kenya and the BirdLife International Africa secretariat helped in various ways. In additon, many others, too numerous to mention, gave encouragement and made useful comments.

The Bird Committee thanks the Kenya Wildlife Service for its financial help towards the printing costs and the Nature Kenya management and staff for their steadfast support.

NATIONAL MUSEUMS OF KENYA
WHERE HERITAGE LIVES ON

NatureKenya
The East Africa Natural History Society

Nature Kenya — the East Africa Natural History Society

Nature Kenya is the East Africa Natural History Society (EANHS) in Kenya. It is a membership-based organization, founded in 1909. Nature Kenya invites all who are not members to join and be part of a strong and active society where members are encouraged to participate in the study of nature and help in the implementation of conservation activities.

Connecting nature and people to take action for biodiversity conservation is the Society's mission statement. Nature Kenya strives to:

- Build a strong network of people for conservation across the country
- Enhance knowledge of Kenya's biodiversity
- Advocate policies favourable to biodiversity conservation
- Promote the conservation of key species, sites, and habitats
- Encourage community participation in conservation through the promotion of sustainable benefits.

Index to English group names corrected 13 December

Akalat 31
Albatross 2
Alethe 31
Ant Thrush 30
Apalis 26
Avocet 9

Babbler 29
Barbet 18
Bat Hawk 6
Batis 20
Bee-eater 17
Bishop 35, 36
Bittern 4
Blackcap 28
Bluebill 36
Booby 5
Boubou 21
Bristlebill 27
Broadbill 20
Brownbul 27
Brubru 21
Buffalo Weaver 34
Bulbul 26
Bunting 39
Bush Chat 31
Bushshrike 20, 21
Bustard 7, 8
Buttonquail 9
Buzzard 7

Camaroptera 26
Canary 38
Chat 32
Chatterer 29
Chiffchaff 28
Cisticola 24, 25
Citril 38
Cliff Chat 32
Coot 8
Cordon-bleu 36
Cormorant 5
Corncrake 8
Coucal 14
Courser 11
Crab-plover 9
Crake 8
Crane 8
Creeper 29
Crimsonwing 36
Crombec 28
Crow 22
Cuckoo 14
Cuckoo Hawk 6
Cuckooshrike 21
Curlew 10

Darter 5
Dove 13
Drongo 22
Duck 2

Eagle 6, 7
Egret 4
Eremomela 28

Falcon 5
Finch 37
Finfoot 8
Firefinch 37
Fiscal 21
Flamingo 3
Flufftail 8
Flycatcher 20, 22, 32
Francolin 1, 2
Frigatebird 4

Gadwall 2
Gallinule 8
Garganey 2
Go-away-bird 14
Godwit 10
Gonolek 21
Goose 2
Goshawk 6
Grassbird 27
Grebe 3
Greenbul 26, 27
Greenshank 10
Grenadier 36
Ground Thrush 30
Guineafowl 1
Gull 11, 12

Hamerkop 4
Harrier 6
Harrier Hawk 6
Helmetshrike 20
Heron 4
Hobby 5
Honey Buzzard 6
Honeyguide 19
Hoopoe 17
Hornbill 18
Hyliota 28

Ibis 3
Illadopsis 29
Indigobird 37
Irania 31

Jacana 10

Kestrel 5
Kingfisher 17
Kite 6
Knot 10

Lammergeier 6
Lapwing 9
Lark 24
Leaflove 27
Longbill 28
Longclaw 38
Lovebird 13

Malimbe 35
Mallard 2
Mannikin 37
Martin 23
Moorhen 8
Mousebird 16

Negrofinch 36
Nicator 27
Night Heron 4
Nightingale 31
Nightjar 15, 16
Noddy 12

Oriole 22
Osprey 6
Ostrich 1
Owlet 15
Owl 15
Oxpecker 30
Oystercatcher 9

Painted-snipe 10
Palm Thrush 31
Parisoma 29
Parrot 13
Pelican 5
Petrel 2, 3
Petronia 34
Phalarope 11
Piapiac 22
Pigeon 13
Pintail 2
Pipit 38
Pitta 20
Plantain-eater 14
Plover 9, 10
Pochard 2
Pratincole 11
Prinia 25, 26
Prion 2
Puffback 21
Pygmy Goose 2
Pytilia 37

Quailplover 9
Quailfinch 37
Quail 1
Quelea 35

Rail 8
Raven 23
Redshank 10
Redstart 31
Robin 31
Robin Chat 31
Rock Chat 32
Rock Thrush 32
Roller 16, 17
Rook 22
Ruff 11

Sanderling 10
Sandgrouse 13
Sandpiper 10, 11
Saw-wing 23
Scimitarbill 18
Scrub Robin 31
Secretarybird 5
Seed-cracker 36
Seedeater 38
Shearwater 2, 3
Shikra 7
Shoebill 4
Shoveler 2
Shrike 21, 22
Silverbill 37
Silverbird 32
Skimmer 12
Skua 12
Snipe 10
Social Weaver 34
Sparrow 34
Sparrow Lark 24
Sparrow Weaver 34
Sparrowhawk 7
Spinetail 16
Spoonbill 3
Sprosser 31
Spurfowl 1
Starling, 29, 30
Stilt 9
Stint 11
Stone-curlew 9
Stonechat 31
Stork 3
Storm-petrel 3
Sunbird 33, 34
Swallow 23, 24
Swamphen 8
Swift 16

Tchagra 21
Teal 2
Tern 12
Thick-knee 9
Thrush 30, 31
Tinkerbird 18, 19
Tit 23
Trogon 16
Tropicbird 4
Turaco 14
Turnstone 10
Twinspot 36

Vulture 6

Wagtail 37, 38
Warbler 26, 27, 28, 29
Wattle-eye 20
Waxbill 36, 37
Weaver 34, 35
Wheatear 31, 32
Whimbrel 10
Whinchat 31
Whistling Duck 1
White-eye 29
Whydah 37
Widowbird 36
Wigeon 2
Wood-hoopoe 17, 18
Woodpecker 19
Wren Warbler 26
Wryneck 19

Yellowbill 14