




A beamconv-based LiteBIRD simulation

Marta Monelli

Max Planck Institut für Astrophysik
Garching (Germany)


December 1st, 2021

A sketch of the pipeline



A sketch of the pipeline


beamconv: convolution code simulating TOD for CMB experiments with realistic polarized beams, scanning strategies and HWP.



A sketch of the pipeline


beamconv: convolution code simulating TOD for CMB experiments with realistic polarized beams, scanning strategies and HWP.

DUCC: collection of basic programming tools for numerical computation: fft, sht, healpix, totalconvolve...




Input

Input maps




The input maps are obtained by making use of PySM, including both **CMB signal** and **foreground emission** (thermal dust, synchrotron, AME and free-free).



Single frequency FG maps (140 GHz).



Scanning strategy

In order to simulate LiteBIRD's scanning strategy, some functionalities of `pyScan` have been implemented in `beamconv`.





Scanning strategy

In order to simulate LiteBIRD's scanning strategy, some functionalities of `pyScan` have been implemented in `beamconv`.




Scanning strategy

In order to simulate LiteBIRD's scanning strategy, some functionalities of pyScan have been implemented in beamconv.




Focal plane specifics



The Instrument Model Database (IMO) contains all the relevant information:


```
{  
    'name': 'M02_030_QA_140T',  
    'wafer': 'M02',  
    'pixel': 30,  
    'pixtype': 'MP1',  
    [...]  
    'pol': 'T',  
    'orient': 'Q',  
    'quat': [1, 0, 0, 0]  
}
```

HWP specifics




$$\mathcal{M}_{\text{ideal}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

HWP specifics




$$\mathcal{M}_{\text{ideal}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$




Noise specifics

The IMO contains also the parameters that enter in the noise power spectrum:

$$P(f) = \textcolor{brown}{NET}^2 \left[\frac{f^2 + \textcolor{brown}{f}_{\text{knee}}^2}{f^2 + \textcolor{brown}{f}_{\text{min}}^2} \right]^{\alpha}.$$




Noise specifics



The IMO contains also the parameters that enter in the noise power spectrum:


$$P(f) = \text{NET}^2 \left[\frac{f^2 + f_{\text{knee}}^2}{f^2 + f_{\text{min}}^2} \right]^\alpha.$$

Noise power spectrum (theory vs 300 realizations averaged) [μK^2]





Injection of the CMB dipole

The dipole signal is calculated by following the procedure employed in `litebird_sim`'s dipole module.




Injection of the CMB dipole

The dipole signal is calculated by following the procedure employed in `litebird_sim`'s dipole module.



Output

Time ordered data




This is a day of observation for a single detector.

The signal is dominated by the foreground emission.

The “periodicity” corresponds to a precession period.

Time ordered data



This is a day of observation for a single detector.

The signal is dominated by the foreground emission.

The “periodicity” corresponds to a precession period.

Let's give a closer look:
[demo on GitHub.](#)


A countercheck (expectation)

Data model with ideal HWP:

$$d = \frac{1}{2}m_{ij}I + \frac{1}{2}[m_{qq}c_1c_2 + m_{uu}s_1s_2]Q + \frac{1}{2}[m_{qq}s_1c_2 - m_{uu}c_1s_2]U.$$

Difference between d with and without HWP:


$$\Delta d = \sin 2(\psi - \xi) [-\sin 2(\phi + \psi)Q + \cos 2(\phi + \psi)U],$$



A countercheck (result)

$$\Delta d = \sin 2(\psi - \xi) [-\sin 2(\phi + \psi)Q + \cos 2(\phi + \psi)U] ,$$

This quantity has opposite sign for detectors sensitive to orthogonal direction of the polarization angle ($\xi \rightarrow \xi \pm \pi/2$). This effect is visible in the simulation:



Moving forward

Still to be implemented

Besides smaller adjustments, there are a few major changes that are will be implemented in the near future:

- ☐ include frequency-dependence of HWP non-idealities;
- ☐ use more realistic beam shapes;
- ☐ adapt the pipeline for production purposes.

Still to be implemented

Besides smaller adjustments, there are a few major changes that are will be implemented in the near future:

- ☐ include frequency-dependence of HWP non-idealities;
- ☐ use more realistic **beam shapes**;
- ☐ adapt the pipeline for production purposes.

Still to be implemented

Besides smaller adjustments, there are a few major changes that are will be implemented in the near future:

- ☐ include frequency-dependence of HWP non-idealities;
- ☐ use more realistic beam shapes;
- ☐ adapt the pipeline for **production** purposes.

Still to be implemented

Besides smaller adjustments, there are a few major changes that are will be implemented in the near future:

- ☐ include frequency-dependence of HWP non-idealities;
- ☐ use more realistic beam shapes;
- ☐ adapt the pipeline for production purposes.

Wish list

Once those changes will be implemented, the pipeline could be used to

- ▷ study how the HWP non-idealities affect the measurement of the cosmic birefringence angle β ;
- ▷ determine requirements on non-idealities so that the systematics on β are well below 0.1° ;
- ▷ study the impact of non-idealities on the EB angle calibration;
- ▷ study the impact of HWP non-idealities on Q/U maps of Tau A.

Wish list

Once those changes will be implemented, the pipeline could be used to

- ▷ study how the HWP non-idealities affect the measurement of the cosmic birefringence angle β ;
- ▷ determine requirements on non-idealities so that the systematics on β are well below 0.1° ;
- ▷ study the impact of non-idealities on the EB angle calibration;
- ▷ study the impact of HWP non-idealities on Q/U maps of Tau A.

Wish list

Once those changes will be implemented, the pipeline could be used to

- ▷ study how the HWP non-idealities affect the measurement of the cosmic birefringence angle β ;
- ▷ determine requirements on non-idealities so that the systematics on β are well below 0.1° ;
- ▷ study the impact of non-idealities on the EB angle calibration;
- ▷ study the impact of HWP non-idealities on Q/U maps of Tau A.

Wish list

Once those changes will be implemented, the pipeline could be used to

- ▷ study how the HWP non-idealities affect the measurement of the cosmic birefringence angle β ;
- ▷ determine requirements on non-idealities so that the systematics on β are well below 0.1° ;
- ▷ study the impact of non-idealities on the EB angle calibration;
- ▷ study the impact of HWP non-idealities on Q/U maps of Tau A.

Wish list

Once those changes will be implemented, the pipeline could be used to

- ▷ study how the HWP non-idealities affect the measurement of the cosmic birefringence angle β ;
- ▷ determine requirements on non-idealities so that the systematics on β are well below 0.1° ;
- ▷ study the impact of non-idealities on the EB angle calibration;
- ▷ study the impact of HWP non-idealities on Q/U maps of Tau A.